

# A $p$ -ADIC ARITHMETIC INNER PRODUCT FORMULA

DANIEL DISEGNI AND YIFENG LIU

**ABSTRACT.** Fix a prime number  $p$  and let  $E/F$  be a CM extension of number fields in which  $p$  splits relatively. Let  $\pi$  be an automorphic representation of a quasi-split unitary group of even rank with respect to  $E/F$  such that  $\pi$  is ordinary above  $p$  with respect to the Siegel parabolic subgroup. We construct the cyclotomic  $p$ -adic  $L$ -function of  $\pi$ , and show, under certain conditions, that if its order of vanishing at the trivial character is 1, then the rank of the Selmer group of the Galois representation of  $E$  associated with  $\pi$  is at least 1. Furthermore, under a certain modularity hypothesis, we use special cycles on unitary Shimura varieties to construct some explicit elements in the Selmer group called Selmer theta lifts; and we prove a precise formula relating their  $p$ -adic heights to the derivative of the  $p$ -adic  $L$ -function. In parallel to Perrin-Riou's  $p$ -adic analogue of the Gross–Zagier formula, our formula is the  $p$ -adic analogue of the arithmetic inner product formula recently established by Chao Li and the second author.

## CONTENTS

1. Introduction	2
1.1. Cyclotomic $p$ -adic $L$ -function	3
1.2. Application to Selmer groups	4
1.3. Application to symmetric power of elliptic curves	5
1.4. A $p$ -adic arithmetic inner product formula	6
1.5. Structure and strategy	6
1.6. Notation and conventions	7
2. Siegel hermitian varieties	8
2.1. Running notation	8
2.2. Siegel hermitian varieties and automorphy line bundles	10
2.3. Relation with PEL type moduli spaces	13
3. Cyclotomic $p$ -adic $L$ -function	16
3.1. Doubling space and degenerate principal series	16
3.2. Siegel hermitian Eisenstein series	21
3.3. Relevant representations	23
3.4. Local doubling zeta integral	27
3.5. Construction of $p$ -adic $L$ -function	29
4. $p$ -adic heights of Selmer theta lifts	33
4.1. The hermitian space	33
4.2. Selmer theta lifts	34
4.3. A $p$ -adic arithmetic inner product formula	38
4.4. Local height away from $p$	41
4.5. Local height above $p$	42
4.6. Proof of main theorems	47
4.7. Generalities on $p$ -adic measures	54
Appendix A. Bi-extensions and $p$ -adic height pairings	55
A.1. Étale correspondences	55
A.2. Remarks on sheaves	55
A.3. Bi-extensions of cycles	58
A.4. Relation with Beilinson's local index	59
A.5. Crystalline property of bi-extensions	61

*Date:* April 29, 2022.

*2010 Mathematics Subject Classification.* 11G18, 11G40, 11G50, 11R34.

A.6. Recollection on $p$ -adic Galois representation	61
A.7. Decomposition of $p$ -adic height pairing	62
Appendix B. Proof of Theorem A.6	63
B.1. Preparation	63
B.2. Rigid de Rham–Witt complexes	64
B.3. Log rigid cohomology with proper support	67
B.4. Abel–Jacobi map via rigid cohomology	68
B.5. Further preparation	73
B.6. Proof of Proposition B.10	75
References	85

## 1. INTRODUCTION

In 1986, Gross and Zagier published a groundbreaking formula relating the heights of Heegner points on modular curves to derivatives of  $L$ -functions, known as the Gross–Zagier formula [GZ86]. For a cuspidal eigenform  $f$  of weight 2, an imaginary quadratic field  $K$  and an unramified Dirichlet character  $\xi$  of  $K$ , the formula shows, under the so-called Heegner condition (which implies that the Rankin–Selberg  $L$ -function  $L(s, f, \xi)$  vanishes at the center 1), that up to some explicit constant,  $L'(1, f, \xi)$  equals the Néron–Tate height of  $H_\xi(f)$  – the  $f$ -isotypic component of the  $K$ -Heegner point weighted by  $\xi$  on a modular curve. Shortly after, Perrin-Riou found the same story in the  $p$ -adic world [PR87]. Namely, she constructed a  $p$ -adic analogue of the (complex)  $L$ -function as a  $p$ -adic measure  $\mathcal{L}_p(f, \xi)$  on the Iwasawa algebra that interpolates  $L(1, f \otimes \chi, \xi)$  where  $\chi$  is a Dirichlet character ramified only at  $p$ , assuming that  $f$  is ordinary at  $p$  and  $p$  splits in  $K$ . She then showed that under the Heegner condition, up to some explicit constant, the derivative of the  $p$ -adic  $L$ -function  $\mathcal{L}_p(f, \xi)$  at the trivial character equals the  $p$ -adic height of  $H_\xi(f)$  – this is referred as the  *$p$ -adic Gross–Zagier formula*.

Since the original work of Gross and Zagier, the Gross–Zagier formula and its  $p$ -adic avatar have been extended to various settings but all (essentially) for curves or fibrations/local systems over curves (see Remark 1.8 below for a brief review of the  $p$ -adic results), until the very recent works by Chao Li and one of us [LL21, LL22]. There, the authors proved a formula computing central  $L$ -derivatives for unitary groups of higher ranks in terms of the Beilinson–Bloch heights of special cycles. This originates from a program initiated by Kudla [Kud02, Kud03, Kud04] and can be regarded as a Gross–Zagier formula in higher dimensions, as well as an arithmetic analogue of Rallis’ inner product formula in the theory of the theta correspondence [Ral82]. The current work contains a  $p$ -adic avatar of the *arithmetic inner product formula* in [LL21, LL22]; this is likewise the first generalization of the  $p$ -adic Gross–Zagier formula to genuinely higher dimensional varieties. A secondary aim of this article is to develop some foundational results in the theory of  $p$ -adic heights of algebraic cycles (in the two appendices); in particular, we prove a crystalline property of bi-extensions, which generalizes the fact that  $p$ -adic regulators take values in Selmer groups.

In the rest of this introduction, we explain our results in more detail. Throughout the article, we fix a prime number  $p$ , an algebraic closure  $\overline{\mathbb{Q}}_p$  of  $\mathbb{Q}_p$ , and a CM extension  $E/F$  of number fields such that every  $p$ -adic place of  $F$  splits in  $E$ . Denote by

- $c \in \text{Gal}(E/F)$  the Galois involution,
- $\mathbb{V}_F^{(\diamond)}$  the set of places of  $F$  above a finite set  $\diamond$  of places of  $\mathbb{Q}$ ,<sup>1</sup>
- $\mathbb{V}_F^{\text{fin}}$  the set of non-archimedean places of  $F$ ,
- $\mathbb{V}_F^{\text{spl}}$ ,  $\mathbb{V}_F^{\text{int}}$  and  $\mathbb{V}_F^{\text{ram}}$  the subsets of  $\mathbb{V}_F^{\text{fin}}$  of those that are split, inert and ramified in  $E$ , respectively.

For every number field  $K$ , we denote by  $\Gamma_{K,p}$  the maximal Hausdorff quotient of

$$K^\times \backslash \mathbb{A}_K^{\infty, \times} \left/ \left( \mathcal{O}_K \otimes \prod_{w \neq p} \mathbb{Z}_w \right)^\times \right.,$$

<sup>1</sup>When  $\diamond = \{w\}$  is a singleton, we simply write  $\mathbb{V}_F^{(w)}$  for  $\mathbb{V}_F^{(\{w\})}$ .

which is naturally a finitely generated  $\mathbb{Z}_p$ -module; and let  $\mathcal{X}_{K,p}$  be the rigid analytic space over  $\mathbb{Q}_p$  such that for every complete topological  $\mathbb{Q}_p$ -ring  $R$ ,  $\mathcal{X}_{K,p}(R)$  is the set of continuous characters from  $\Gamma_{K,p}$  to  $R^\times$ .

**1.1. Cyclotomic  $p$ -adic  $L$ -function.** Take an integer  $r \geq 1$  and put  $n = 2r$ . We equip  $W_r := E^n$  with the skew-hermitian form (with respect to  $\mathfrak{c}$ ) given by the matrix  $w_r := \begin{pmatrix} & 1_r \\ -1_r & \end{pmatrix}$ . Put  $G_r := U(W_r)$ , the unitary group of  $W_r$ , which is a quasi-split reductive group over  $F$ . Denote by  $\dagger$  the involution of  $G_r$  given by the conjugation by the element  $\begin{pmatrix} 1_r & \\ & -1_r \end{pmatrix}$  inside  $\text{Res}_{E/F} \text{GL}_n$ . For  $v \in \mathbb{V}_F^{\text{fin}}$ , let  $K_{r,v} \subseteq G_r(F_v)$  be the stabilizer of the lattice  $\mathcal{O}_{E_v}^n$ .

**Definition 1.1.** Let  $\mathbb{L}$  be a field embeddable into  $\mathbb{C}$ . A *relevant  $\mathbb{L}$ -representation* of  $G_r(\mathbb{A}_F^\infty)$  is a representation  $\pi$  with coefficients in  $\mathbb{L}$  satisfying that for every embedding  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ ,

$$\left( \otimes_{v \in \mathbb{V}_F^{(\infty)}} \pi_v^{[r]} \right) \otimes \iota \pi$$

is a tempered cuspidal automorphic representation of  $G_r(\mathbb{A}_F)$ . Here, for  $v \in \mathbb{V}_F^{(\infty)}$ ,  $\pi_v^{[r]}$  denotes the unique holomorphic discrete series representation of  $G_r(F_v) = G_r(\mathbb{R})$  with the Harish-Chandra parameter  $\{\frac{1-n}{2}, \frac{3-n}{2}, \dots, \frac{n-3}{2}, \frac{n-1}{2}\}$ . In particular,  $\pi$  is admissible and absolutely irreducible.

We consider a finite extension  $\mathbb{L}/\mathbb{Q}_p$  contained in  $\overline{\mathbb{Q}_p}$  and a relevant  $\mathbb{L}$ -representation  $\pi$  of  $G_r(\mathbb{A}_F^\infty)$ . By Lemma 3.13,  $\hat{\pi} := (\pi^\vee)^\dagger$  is a relevant  $\mathbb{L}$ -representation of  $G_r(\mathbb{A}_F^\infty)$  as well.

**Definition 1.2.** For  $v \in \mathbb{V}_F^{(p)}$ , let  $P_v$  be the set (of two elements) of places of  $E$  above  $v$ . For  $u \in P_v$ , we have the representation  $\pi_u$  of  $\text{GL}_n(F_v)$  as a local component of  $\pi$  via the isomorphism  $G_r(F_v) \simeq \text{GL}_n(E_u) = \text{GL}_n(F_v)$ . In particular,  $\pi_u^\vee \simeq \pi_{u^c}$ . We say that  $\pi_u$  is *Panchishkin unramified* if

- (1)  $\pi_u$  is unramified;
- (2) if we write the Satake polynomial of  $\pi_u$ , which makes sense by (1), as

$$P_{\pi_u}(T) = T^n + \beta_{u,1} \cdot T^{n-1} + \beta_{u,2} \cdot q_v \cdot T^{n-2} + \dots + \beta_{u,r} \cdot q_v^{\frac{r(r-1)}{2}} \cdot T^r + \dots + \beta_{u,n} \cdot q_v^{\frac{n(n-1)}{2}} \in \mathbb{L}[T]$$

(see Definition 3.17 for more details), then  $\beta_{u,r} \in O_{\mathbb{L}}^\times$ , where  $q_v$  is the residue cardinality of  $F_v$ .

We collect two important facts about Panchishkin unramified representations:

- The representation  $\pi_u$  is Panchishkin unramified if and only if  $\pi_{u^c}$  is (Lemma 3.21). In particular, it makes sense to say that  $\pi_v$  is Panchishkin unramified.
- If  $\pi_u$  is Panchishkin unramified, then there is a unique polynomial  $Q_{\pi_u}(T) \in \mathbb{L}[T]$  that divides  $P_{\pi_u}(T)$  and has the form

$$Q_{\pi_u}(T) = T^r + \gamma_{u,1} \cdot T^{r-1} + \gamma_{u,2} \cdot q_v \cdot T^{r-2} + \dots + \gamma_{u,r} \cdot q_v^{\frac{r(r-1)}{2}}$$

with  $\gamma_{u,r} \in O_{\mathbb{L}}^\times$  (Proposition 3.24). In particular, we have an unramified principal series  $\underline{\pi}_u$  of  $\text{GL}_r(F_v)$  defined over  $\mathbb{L}$  whose Satake polynomial is  $Q_{\pi_u}(T)$ .

*Remark 1.3.* In fact,  $\pi_v$  is Panchishkin unramified if and only if  $\pi_v$  is unramified and  $\pi$  is ordinary at  $v$  with respect to the standard Siegel parabolic subgroup of  $G_r$  in the sense of Hida [Hid98].

**Theorem 1.4.** *Under the above setup, suppose that  $\pi_v$  is Panchishkin unramified for every  $v \in \mathbb{V}_F^{(p)}$ . For every finite set  $\diamond$  of places of  $\mathbb{Q}$  containing  $\{\infty, p\}$  such that  $\pi_v$  is unramified for every  $v \in \mathbb{V}_F^{\text{fin}} \setminus \mathbb{V}_F^{(\diamond)}$ , there is a unique bounded analytic function  $\mathcal{L}_p^\diamond(\pi)$  on the rigid analytic space  $\mathcal{X}_{F,p} \otimes_{\mathbb{Q}_p} \mathbb{L}$  such that for every finite (continuous) character  $\chi: \Gamma_{F,p} \rightarrow \overline{\mathbb{Q}_p}^\times$  and every isomorphism  $\iota: \overline{\mathbb{Q}_p} \xrightarrow{\sim} \mathbb{C}$ , we have*

$$\iota \mathcal{L}_p^\diamond(\pi)(\chi) = \frac{1}{P_\pi} \cdot \frac{Z_r^{[F:\mathbb{Q}]}}{b_{2r}^\diamond(\mathbf{1})} \cdot \prod_{v \in \mathbb{V}_F^{(p)}} \prod_{u \in P_v} \gamma(\frac{1+r}{2}, \iota(\underline{\pi}_u \otimes \chi_v), \psi_{F,v})^{-1} \cdot L(\frac{1}{2}, \text{BC}(\iota\pi^\diamond) \otimes (\iota\chi^\diamond \circ \text{Nm}_{E/F})),$$

where

- $P_\pi \in \mathbb{C}^\times$  is a certain period for  $\pi$  with respect to  $\iota$  for every embedding  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ ;
- $Z_r := (-1)^r 2^{-r^2-r} \pi^{r^2} \frac{\Gamma(1) \cdots \Gamma(r)}{\Gamma(r+1) \cdots \Gamma(2r)}$  is the value of a certain explicit archimedean local doubling zeta integral;
- $b_{2r}^\diamond(\mathbf{1}) = \prod_{i=1}^{2r} L^\diamond(i, \eta_{E/F}^i)$  is defined in §2.1(F3);

- $\gamma(s, \iota(\pi_u \otimes \chi_v), \psi_{F,v})$  is the gamma factor [Jac79] in which  $\psi_F := \psi_{\mathbb{Q}} \circ \text{Tr}_{F/\mathbb{Q}}$  with  $\psi_{\mathbb{Q}}: \mathbb{A}_{\mathbb{Q}} \rightarrow \mathbb{C}^{\times}$  the standard automorphic additive character.

By definition, for every  $v \in \mathbb{V}_F^{(p)}$  and  $u \in P_v$ ,  $\iota(\pi_u \otimes \chi_v) |_{\mathbb{Z}_p}$  is tempered so that  $\gamma(\frac{1+r}{2}, \iota(\pi_u \otimes \chi_v), \psi_{F,v}) \in \mathbb{C}^{\times}$ .

*Remark 1.5.* We have the following remarks concerning Theorem 1.4.

- (1) A bounded analytic function on the rigid analytic space  $\mathcal{X}_{F,p} \otimes_{\mathbb{Q}_p} \mathbb{L}$  is equivalent to an element in  $\mathbb{Z}_p[[\Gamma_{F,p}]] \otimes_{\mathbb{Z}_p} \mathbb{L}$ , that is, an  $\mathbb{L}$ -valued  $p$ -adic measure on  $\Gamma_{F,p}$ . In particular, the uniqueness of  $\mathcal{L}_p^{\diamond}(\pi)$  is clear.
- (2) The collection of periods  $(P_{\pi}^t)_t$  is only well-defined up to a common factor in  $\mathbb{L}^{\times}$  (see Notation 3.14). In particular, the  $p$ -adic  $L$ -function  $\mathcal{L}_p^{\diamond}(\pi)$  is only well-defined up to a factor in  $\mathbb{L}^{\times}$ .
- (3) The vanishing order of  $\mathcal{L}_p^{\diamond}(\pi)$  at  $\mathbf{1}$  is independent of  $\diamond$ .
- (4) Our  $p$ -adic  $L$ -function is defined over the  $p$ -adic field of definition of the representation and interpolates complex  $L$ -values along *all* isomorphisms  $\overline{\mathbb{Q}_p} \simeq \mathbb{C}$ ; this is a stronger rationality property than the one under a fixed isomorphism  $\overline{\mathbb{Q}_p} \simeq \mathbb{C}$  as in the setup of many previous works in this field.
- (5) Among other technical assumptions, at least when  $\pi$  is ordinary at  $p$  in the usual sense (that is, for every  $u \in P$ ,  $\beta_{u,m} \in O_{\mathbb{L}}^{\times}$  for every  $1 \leq m \leq n$  in the Satake polynomial of  $\pi_u$ ), our  $p$ -adic  $L$ -function has already been constructed in [Ehls20] up to some constant (and with a weaker rationality property). In fact, in [Ehls20] the authors construct more generally a multi-variable  $p$ -adic  $L$ -function in which  $\pi$  is allowed to vary in an ordinary Hida family as well. In this article, we will give a (relatively) self-contained construction of our  $p$ -adic  $L$ -function independent of [Ehls20] since first, the process of the construction itself is an ingredient for the  $p$ -adic height formula; and second, our construction is technically much simpler to follow.

## 1.2. Application to Selmer groups.

**Assumption 1.6.** Suppose that  $F \neq \mathbb{Q}$ , that  $\mathbb{V}_F^{\text{spl}}$  contains all 2-adic (and  $p$ -adic) places, and that every prime in  $\mathbb{V}_F^{\text{ram}}$  is unramified over  $\mathbb{Q}$ . We consider a relevant  $\mathbb{L}$ -representation  $\pi$  of  $G_r(\mathbb{A}_F^{\infty})$  for some finite extension  $\mathbb{L}/\mathbb{Q}_p$  contained in  $\overline{\mathbb{Q}_p}$ , satisfying:

- (1) For every  $v \in \mathbb{V}_F^{\text{ram}}$ ,  $\pi_v$  is spherical with respect to  $K_{r,v}$ , that is,  $\pi_v^{K_{r,v}} \neq \{0\}$ .
- (2) For every  $v \in \mathbb{V}_F^{\text{int}}$ ,  $\pi_v$  is either unramified or almost unramified (see [LL21, Remark 1.4(3)]) with respect to  $K_{r,v}$ ; moreover, if  $\pi_v$  is almost unramified, then  $v$  is unramified over  $\mathbb{Q}$ .
- (3) We have  $R_{\pi} \cup S_{\pi} \subseteq \mathbb{V}_F^{\diamond}$  (Definition 4.3), where
  - $R_{\pi} \subseteq \mathbb{V}_F^{\text{spl}}$  denotes the (finite) subset for which  $\pi_v$  is ramified,
  - $S_{\pi} \subseteq \mathbb{V}_F^{\text{int}}$  denotes the (finite) subset for which  $\pi_v$  is almost unramified.
- (4) For every  $v \in \mathbb{V}_F^{(p)}$ ,  $\pi_v$  is Panchishkin unramified.

Let  $\pi$  be as in Assumption 1.6. The (doubling) root number of  $\pi$  equals  $(-1)^{|S_{\pi}|}$ . Note that for every  $v \in \mathbb{V}_F^{(\infty)}$ , the root number of  $\pi_v^{[r]}$  equals  $(-1)^r$ . It follows that if  $r[F : \mathbb{Q}] + |S_{\pi}|$  is odd, then the  $p$ -adic  $L$ -function  $\mathcal{L}_p^{\diamond}(\pi)$  constructed in Theorem 1.4 vanishes at the trivial character  $\mathbf{1}$ . Thus, it motivates us to consider the derivative  $\partial \mathcal{L}_p^{\diamond}(\pi)(\mathbf{1})$  as an element in the cotangent space of  $\Gamma_{F,p}$  at  $\mathbf{1}$ , which is canonically  $\Gamma_{F,p} \otimes_{\mathbb{Z}_p} \mathbb{L}$ .

Associated with  $\pi$ , we have a semisimple continuous representation  $\rho_{\pi}$  of  $\text{Gal}(\overline{E}/E)$  of dimension  $n$  with coefficients in  $\overline{\mathbb{Q}_p}$ , which satisfies  $\rho_{\pi}^c \simeq \rho_{\pi}^{\vee}(1-n)$  (Lemma 4.8). Then in the interpolation property of  $\mathcal{L}_p^{\diamond}(\pi)$  in Theorem 1.4, we have

$$L(\frac{1}{2}, \text{BC}(\iota\pi^{\diamond}) \otimes (\iota\chi^{\diamond} \circ \text{Nm}_{E/F})) = L^{\diamond}(0, \iota(\rho_{\pi}(r) \otimes \chi|_{\text{Gal}(\overline{E}/E)})),$$

where on the right-hand side we view  $\chi$  as a  $\overline{\mathbb{Q}_p}$ -valued character of  $\text{Gal}(\overline{E}/E)$  via the global class field theory. In particular, the theorem below provides evidence toward the  $p$ -adic Beilinson–Bloch–Kato conjecture for (genuinely) higher-dimensional motives.

**Theorem 1.7.** *Suppose that  $n < p$ . Let  $\pi$  be as in Assumption 1.6 with  $r[F : \mathbb{Q}] + |S_{\pi}|$  odd (so that  $\mathcal{L}_p^{\diamond}(\pi)(\mathbf{1}) = 0$ ). If  $\partial \mathcal{L}_p^{\diamond}(\pi)(\mathbf{1})$  is nonvanishing, then*

$$\dim_{\overline{\mathbb{Q}_p}} H_f^1(E, \rho_{\pi}(r)) \geq 1,$$

where  $H_f^1$  denotes the Bloch–Kato Selmer group [BK90].

*Remark 1.8.* When  $n = 2$ , this result is a variant of the main application of the  $p$ -adic Gross–Zagier formula of [PR87], as generalized to totally real fields by one of us [Dis17] following the development of [GZ86] in [YZZ13]. In different directions, Perrin-Riou’s results had been generalized to the case of higher-weight modular forms by Nekovář [Nek95] and further to the case with twists by higher-weight Hecke characters by Shnidman [Shn16], to the supersingular case by Kobayashi [Kob13], and to the case where  $p$  is not necessarily relative split by one of us [Disa].<sup>2</sup> A common generalization of [Nek95, Shn16, Dis17, Disa] was developed in [Disb].

*Remark 1.9.* Strictly speaking, Theorem 1.7, together with Corollary 1.10 and Theorem 1.11 below, relies on a hypothesis on the characterization of the tempered part of the cohomology of certain unitary Shimura varieties (see Hypothesis 4.9 and Remark 4.10), which is expected to be verified in a sequel of the work [KSZ].

**1.3. Application to symmetric power of elliptic curves.** The above results have applications to the motives of symmetric power of elliptic curves. For simplicity, we only describe the case where  $E$  contains an imaginary quadratic field. More precisely, we consider

- an imaginary quadratic field  $E_0$  in which  $p$  splits,
- a solvable totally real field  $F$ ,
- an elliptic curve  $A$  over  $\mathbb{Q}$  without complex multiplication that has ordinary good reduction at  $p$ .

Denote by  $\diamond_{E_0}$ ,  $\diamond_F$  and  $\diamond_A$  the sets of primes that divide the discriminants of  $E_0$ ,  $F$  and  $A$ , respectively. Put  $E := FE_0$  so that  $E/F$  is a CM extension of number fields such that every  $p$ -adic place of  $F$  splits in  $E$ .

By the modularity of  $A$  and the very recent breakthrough on the automorphy of holomorphic modular forms [NT21a, NT21b], there exists a unique cuspidal automorphic representation  $\Pi(\mathrm{Sym}^{n-1} A)$  of  $\mathrm{GL}_n(\mathbb{A})$  satisfying

- $\Pi(\mathrm{Sym}^{n-1} A)_\infty$  is the principal series representation of characters  $(\arg^{1-n}, \arg^{3-n}, \dots, \arg^{n-3}, \arg^{n-1})$ , where  $\arg: \mathbb{C}^\times \rightarrow \mathbb{C}^\times$  is the character given by  $\arg(z) := z/\sqrt{z\bar{z}}$ ;
- for every prime  $w \notin \diamond_A$ ,  $\Pi(\mathrm{Sym}^{n-1} A)_w$  is unramified with the Satake polynomial

$$\prod_{j=0}^{n-1} (T - \alpha_{w,1}^j \alpha_{w,2}^{n-1-j}) \in \mathbb{Q}[T],$$

where  $\alpha_{w,1}$  and  $\alpha_{w,2}$  are the two roots of the polynomial  $T^2 - a_w(A)T + w$ .

Let  $\Pi(\mathrm{Sym}^{n-1} A_E)$  be the (solvable) base change of  $\Pi(\mathrm{Sym}^{n-1} A)$  to  $E$ , which is a cuspidal automorphic representation of  $\mathrm{GL}_n(\mathbb{A}_E)$ . The representation  $\Pi(\mathrm{Sym}^{n-1} A_E)$  satisfies  $\Pi(\mathrm{Sym}^{n-1} A_E)^\vee \simeq \Pi(\mathrm{Sym}^{n-1} A_E) \simeq \Pi(\mathrm{Sym}^{n-1} A_E)^c$ , hence is a relevant representation in the sense of [LTX<sup>+</sup>22, Definition 1.1.3]. By [LTX<sup>+</sup>22, Remark 1.1.4] and the endoscopic classification for quasi-split unitary groups [Mok15], there exists a cuspidal automorphic representation  $\pi(\mathrm{Sym}^{n-1} A_E)$  of  $G_r(\mathbb{A}_F)$  satisfying

- for every  $v \in \mathbb{V}_F^{(\infty)}$ ,  $\pi(\mathrm{Sym}^{n-1} A_E)_v$  is isomorphic to  $\pi_v^{[r]}$ ;
- for every  $v \in \mathbb{V}_F^{\mathrm{fin}} \setminus \mathbb{V}_F^{(\diamond_A)}$ ,  $\pi(\mathrm{Sym}^{n-1} A_E)_v$  is spherical with respect to  $K_{r,v}$  and its base change to  $\mathrm{GL}_n(E_v)$  is isomorphic to  $\Pi(\mathrm{Sym}^{n-1} A_E)_v$ .

In particular, there exists a relevant  $\mathbb{Q}$ -representation  $\pi$  in the sense of Definition 1.1 such that  $\pi(\mathrm{Sym}^{n-1} A_E)^\infty \simeq \pi \otimes_{\mathbb{Q}} \mathbb{C}$ . Moreover, for every  $v \in \mathbb{V}_F^{(p)}$ ,  $\pi_v \otimes_{\mathbb{Q}} \mathbb{Q}_p$  is Panchishkin unramified. Applying Theorem 1.4 to  $\pi$  (or rather  $\pi \otimes_{\mathbb{Q}} \mathbb{Q}_p$ ), we obtain a bounded analytic function  $\mathcal{L}_p^\diamond(\pi)$  on  $\mathcal{X}_{F,p}$  for every finite set  $\diamond$  of places of  $\mathbb{Q}$  containing  $\diamond_{E_0} \cup \diamond_A \cup \{\infty, p\}$ . For every  $v \in \mathbb{V}_F^{(p)}$  and  $u \in \mathbb{P}_v$ , the unramified representation  $\underline{\pi}_u$  of  $\mathrm{GL}_r(F_v)$  is the one with the Satake polynomial

$$\prod_{j=r}^{n-1} \left( T - \left( \alpha_{p,1}^j \alpha_{p,2}^{n-1-j} \right)^{f_u} \right) \in \mathbb{Q}_p[T],$$

where we have ordered  $\alpha_{p,1}, \alpha_{p,2} \in \mathbb{Q}_p^\times$  in the way that  $\alpha_{p,i} \in p^{i-1} \mathbb{Z}_p^\times$  and  $f_u$  is the residue extension degree of  $E_u/\mathbb{Q}_p$ . The following is an immediate corollary of Theorem 1.7 in which  $S_\pi = \emptyset$ .

**Corollary 1.10.** *Under the above setup, we further assume that*

<sup>2</sup>In fact, in [Kob13], a formula in the nonsplit case is deduced from the split case by making use of some special features of the setup under consideration.

- $n < p$ ,
- $[F : \mathbb{Q}] > 1$ ,
- $r[F : \mathbb{Q}]$  is odd,
- $\diamond_{E_0} \cap \diamond_F = \emptyset$ ,
- every prime in  $\diamond_A \cup \{2\}$  splits in  $E_0$ .

Then  $\mathcal{L}_p(\pi)(\mathbf{1}) = 0$ . Moreover, if  $\partial \mathcal{L}_p(\pi)(\mathbf{1})$  is nonvanishing, then

$$\dim_{\mathbb{Q}_p} H_f^1(E, \text{Sym}^{n-1} H_{\text{ét}}^1(A_{\overline{E}}, \mathbb{Q}_p)(r)) \geq 1.$$

**1.4. A  $p$ -adic arithmetic inner product formula.** Consider  $E/F$  and  $\pi$  as in Assumption 1.6 with  $r[F : \mathbb{Q}] + |\mathbb{S}_\pi|$  odd. We fix an embedding  $E \hookrightarrow \mathbb{C}$  and regard  $E$  as a subfield of  $\mathbb{C}$ . By local theta dichotomy,  $\pi$  determines a hermitian space  $V$  over  $E$  of rank  $n = 2r$  that has signature  $(n-1, 1)$  along the induced inclusion  $F \subseteq \mathbb{R}$  and signature  $(n, 0)$  at other archimedean places of  $F$ . Put  $H := U(V)$ . We then have a system of Shimura varieties  $\{X_L\}_L$  indexed by neat open compact subgroups  $L$  of  $H(\mathbb{A}_F^\infty)$ , which are smooth projective schemes over  $E$  of dimension  $n-1$ .

For the simplicity of the introduction, we fix an isomorphism  $\overline{\mathbb{Q}_p} \simeq \mathbb{C}$  here. Take a neat open compact subgroup  $L \subseteq H(\mathbb{A}_F^\infty)$ . Let  $V_{\pi,L}$  and  $V_{\hat{\pi},L}$  be the  $\pi$  and  $\hat{\pi}$  nearly isotypic subspaces of  $H^{2r-1}(X_L \otimes_E \overline{E}, \mathbb{L}(r))$ , respectively. For every  $\phi \in \mathcal{S}(V^r \otimes_F \mathbb{A}_F^\infty)^L$  for which we assume a certain modularity hypothesis (Hypothesis 4.4) and every  $\varphi \in \pi$ , we will construct an element

$$\Theta_\phi^{\text{Sel}}(\varphi)_L \in H_f^1(E, V_{\pi,L})$$

(Definition 4.12) that is the cohomological analogue of the arithmetic theta lifts constructed in [Liu11a, LL21]. The Poincaré duality induces a perfect pairing  $V_{\hat{\pi},L} \times V_{\pi,L} \rightarrow \mathbb{L}(1)$ . By Nekovář's theory [Nek93], we have a  $p$ -adic height pairing

$$\langle \cdot, \cdot \rangle_E := H_f^1(E, V_{\hat{\pi},L}) \times H_f^1(E, V_{\pi,L}) \rightarrow \Gamma_{E,p} \otimes_{\mathbb{Z}_p} \mathbb{L}$$

using certain canonical Hodge splitting at  $p$ -adic places (see §4.3 for more details). For every  $\varphi_1 \in \hat{\pi}$ , every  $\varphi_2 \in \pi$  and every pair  $\phi_1, \phi_2 \in \mathcal{S}(V^r \otimes_F \mathbb{A}_F^\infty)^L$  for which Hypothesis 4.4 holds, the height

$$\text{vol}^{\natural}(L) \cdot \langle \Theta_{\phi_1}^{\text{Sel}}(\varphi_1)_L, \Theta_{\phi_2}^{\text{Sel}}(\varphi_2)_L \rangle_E \in \Gamma_{E,p} \otimes_{\mathbb{Z}_p} \mathbb{C}$$

is independent of  $L$ , where  $\text{vol}^{\natural}(L)$  denotes a certain canonical volume of  $L$  introduced in [LL21, Definition 3.8]. We will denote the above canonical value as  $\langle \Theta_{\phi_1}^{\text{Sel}}(\varphi_1), \Theta_{\phi_2}^{\text{Sel}}(\varphi_2) \rangle_{\pi,E}^{\natural}$ .

**Theorem 1.11** (Theorem 4.17). *Suppose that  $n < p$ . Let  $\pi$  be as in Assumption 1.6 with  $r[F : \mathbb{Q}] + |\mathbb{S}_\pi|$  odd. We also assume Hypothesis 4.4 for every element in  $\mathcal{S}(V^r \otimes_F \mathbb{A}_F^\infty)$ . For a finite set  $\diamond$  as in Theorem 1.4 and every choice of elements*

- $\varphi_1 = \otimes_v \varphi_{1,v} \in \hat{\pi}$  and  $\varphi_2 = \otimes_v \varphi_{2,v} \in \pi$  such that for every  $v \notin \mathbb{V}_F^{(\diamond)}$ ,  $\varphi_{1,v}$  and  $\varphi_{2,v}$  are fixed by  $K_{r,v}$  such that  $\langle \varphi_{1,v}, \varphi_{2,v} \rangle_{\pi_v} = 1$ ,
- $\phi_1 = \otimes_v \phi_{1,v}, \phi_2 = \otimes_v \phi_{2,v} \in \mathcal{S}(V^r \otimes_F \mathbb{A}_F^\infty)$  with  $\phi_1^\diamond = \phi_2^\diamond$  being the characteristic function of  $(\Lambda^\diamond)^r$  in which  $\Lambda^\diamond$  is a self-dual lattice of  $V \otimes_F \mathbb{A}_F^\diamond$ ,

the identity

$$\text{Nm}_{E/F} \langle \Theta_{\phi_1}^{\text{Sel}}(\varphi_1), \Theta_{\phi_2}^{\text{Sel}}(\varphi_2) \rangle_{\pi,E}^{\natural} = \partial \mathcal{L}_p(\pi)(\mathbf{1}) \cdot \prod_{v \in \mathbb{V}_F^{(p)}} \prod_{u \in \mathbb{P}_v} \gamma\left(\frac{1+r}{2}, \pi_u, \psi_{F,v}\right) \cdot \prod_{v \in \mathbb{V}_F^{(\diamond) \setminus \{\infty\}}} Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\phi_{1,v} \otimes \phi_{2,v}}^{\text{SW}})$$

holds in  $\Gamma_{F,p} \otimes_{\mathbb{Z}_p} \mathbb{C}$ , where the term  $Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\phi_{1,v} \otimes \phi_{2,v}}^{\text{SW}})$  is the local doubling zeta integral with respect to the Siegel–Weil section  $f_{\phi_{1,v} \otimes \phi_{2,v}}^{\text{SW}}$  associated with  $\phi_{1,v} \otimes \phi_{2,v}$ .

**1.5. Structure and strategy.** We explain the structure of the article and the strategy for the proofs. Before that, we point out that throughout the article, we have restricted ourselves to only use  $p$ -adic measures valued in finite extensions of  $\mathbb{Q}_p$  to reduce the technical burden such as infinite dimensional  $p$ -adic Banach spaces.

In Section 2, we make preparation for proving the rationality property of our  $p$ -adic  $L$ -function. In §2.1, we collect two sets of more specialized notation that will be used throughout the main part of the article. In §2.2, we introduce the notion of Siegel hermitian varieties which are over  $\mathbb{Q}_p$  and are the main stage to characterize the rationality of automorphic forms on the unitary group  $G_r$ . In §2.3, we review the construction of an auxiliary

Shimura variety over  $\mathbb{Q}$  that is of PEL type in the sense of Kottwitz, which is needed to prove the rationality of certain Eisenstein series used in the doubling method. The main reason we pass to this auxiliary one is that the theory of algebraic  $q$ -expansions is only available for such Shimura varieties. However, if the reader is satisfied with fixing an isomorphism  $\overline{\mathbb{Q}}_p \simeq \mathbb{C}$  from the beginning and does not care about the field of definition of the  $p$ -adic  $L$ -function, then there is no need to use those parts of §2.2 that are related to Shimura varieties and the entire §2.3.

In Section 3, we construct the  $p$ -adic  $L$ -function. The main strategy is to use the doubling method for an “analytic” family of sections in the degenerate principal series of the doubling unitary group  $G_{2r}$ , similar to [Ehls20]. However, it is worth pointing out that our computation makes no use of Weil representations (or their twisted versions). In particular, we do not need any explicit Schwartz functions on hermitian spaces. In fact, we do not even need an explicit formula for the sections in the degenerate principal series at  $p$ -adic places – what we need is just their Fourier transforms, which have very simple forms. The main reason we can simplify the computation is a formula obtained in the previous work [LL21] for computing the local doubling zeta integral (see Lemma 3.25). Using this formula, the gamma factor in Theorem 1.4 appears naturally and immediately. In §3.1, we review the doubling degenerate principal series and collect some facts on their Siegel–Fourier coefficients. In §3.2, we review the doubling Eisenstein series and prove a certain rationality property of their pullbacks to the diagonal block. In §3.3, we make all the representational-theoretical preparations; in particular, we study Panchishkin unramified representations. In §3.4, we prove several formulae for local doubling zeta integrals. In §3.5, we complete the construction of the  $p$ -adic  $L$ -function by defining it as an inner product of a specific element in  $\hat{\pi} \boxtimes \pi$  and the pullback of the family of doubling Eisenstein series with respect to a careful choice of sections in degenerate principal series.

In Section 4, we construct the so-called Selmer theta lifts under a certain modularity hypothesis, which are Selmer group analogues of the classical theta lifts, and study their  $p$ -adic heights. Before further discussion, we point out that such modularity hypothesis is automatic after certain Hecke actions if we assume the converse of Theorem 1.7, that is,  $H_f^1(E, \rho_\pi(r))$  vanishes. This is why we do not need such modularity hypothesis in the statement of Theorem 1.7. In §4.1, we introduce further notation for the whole section. In §4.2, we construct Selmer theta lifts  $\Theta_\phi^{\text{Sel}}(\varphi)_L$  as the inner product of a form  $\varphi$  of  $\pi$  (or  $\hat{\pi}$ ) and the cycle class of Kudla’s generating function  $Z_{\phi,L}$  for  $\phi \in \mathcal{S}(V^r \otimes_F \mathbb{A}_F^\infty)^L$ , assuming the modularity hypothesis. We show that  $\Theta_\phi^{\text{Sel}}(\varphi)_L$  is naturally an element in the Selmer group  $H^1(E, V_{\pi,L})$  (or  $H^1(E, V_{\hat{\pi},L})$ ). In §4.3, we state the precise version of our  $p$ -adic arithmetic inner product formula, which is slightly stronger than Theorem 1.11 by taking rationality into account. We explain that it makes sense and suffices to consider the  $p$ -adic height pairing  $\langle Z_{T_1}(\phi_1), Z_{T_2}(\phi_2) \rangle_E$  between (weighted) special cycles for a certain pool of Schwartz functions, together with a formula decomposing the (global)  $p$ -adic height pairing into local ones. In §4.4, we compute local  $p$ -adic height pairings between special cycles at (nonarchimedean) places of  $E$  not above  $p$ , based on Theorem A.4 that relates local  $p$ -adic heights to Beilinson’s local indices and the formulae for the latter from previous works [LL21, LL22]. In §4.5, we study local  $p$ -adic height pairings between special cycles at  $p$ -adic places of  $E$ . With a crucial ingredient (Theorem A.6) on the crystalline property of the corresponding bi-extensions, we show that the local  $p$ -adic heights approach to 0  $p$ -adically when one repeatedly applies a certain operator  $U_p$  to the Schwartz functions. In §4.6, we finish the proof of Theorem 1.7 and Theorem 4.17 by using the previous formulae on local  $p$ -adic heights and a limiting argument. In §4.7, we collect some basic facts about  $p$ -adic measures that are used in the previous subsection.

The article has two appendices. In Appendix A, we develop further the theory of  $p$ -adic heights on *general varieties*, after Nekovář. In particular, we prove a comparison result between local  $p$ -adic heights and Beilinson’s local indices. For local  $p$ -adic heights above  $p$ , we prove a key theorem (Theorem A.6) on the crystalline property for certain bi-extensions, whose proof occupies the entire Appendix B.

## 1.6. Notation and conventions.

- We denote  $\mathbb{N} := \{0, 1, 2, \dots\}$ .
- We write  $\pi = 3.1415926\dots$ , to be distinguished from the representation  $\pi$ . We also write  $i$  for the imaginary unit in  $\mathbb{C}$ , to be distinguished from the commonly used index  $i$ .
- When we have a function  $f$  on a product set  $A_1 \times \dots \times A_s$ , we will write  $f(a_1, \dots, a_s)$  instead of  $f((a_1, \dots, a_s))$  for its value at an element  $(a_1, \dots, a_s) \in A_1 \times \dots \times A_s$ .
- For a set  $S$ , we denote by  $\mathbf{1}_S$  the characteristic function of  $S$ .

- All rings are commutative and unital; and ring homomorphisms preserve units. However, we use the word *algebra* in the general sense, which is not necessarily commutative or unital.
- If a base ring is not specified in the tensor operation  $\otimes$ , then it is  $\mathbb{Z}$ .
- For an abelian group  $A$  and a ring  $R$ , we put  $A_R := A \otimes R$  as an  $R$ -module.
- For an abelian group  $A$ , we denote by  $A^{\text{fr}}$  its free quotient.
- For a ring  $R$ , we denote by  $\text{Sch}'_R$  the category of locally Noetherian schemes over  $R$ .
- We denote by  $\mathbf{G}$  the multiplicative group scheme, that is,  $\text{Spec } \mathbb{Z}[X, X^{-1}]$ .
- For an integer  $m \geq 0$ , we denote by  $0_m$  and  $1_m$  the null and identity matrices of rank  $m$ , respectively, and by  $w_m$  the matrix  $\begin{pmatrix} & & & \\ & & & \\ & & & \\ -1_m & & & 1_m \end{pmatrix}$ .
- Let  $\psi_{\mathbb{Q}}: \mathbb{Q} \setminus \mathbb{A}_{\mathbb{Q}} \rightarrow \mathbb{C}^{\times}$  be the standard automorphic additive character that sends  $w^{-1}$  at a prime  $w$  to  $\exp(-2\pi i/w)$ , and put  $\psi_K := \psi_{\mathbb{Q}} \circ \text{Tr}_{K/\mathbb{Q}}$  for every number field  $K$ .
- For a subring  $R \subseteq \mathbb{C}$  and a positive integer  $\Delta$ , we denote by  $R\langle \Delta \rangle \subseteq \mathbb{C}$  the subring generated by  $\Delta^l$ -th roots of unity for all  $l \geq 0$ .
- For a locally compact totally disconnected space  $X$  and a ring  $R$ , we denote by  $\mathcal{S}(X, R)$  the  $R$ -module of  $R$ -valued locally constant compactly supported functions on  $X$ . We omit  $R$  from the notation when  $R = \mathbb{C}$ .

*Acknowledgements.* D. D. would like to thank Ellen Eischen and Zheng Liu for correspondence on  $p$ -adic  $L$ -functions. Y. L. would like to thank Yichao Tian for some general discussion related to Theorem A.6. We would like to thank Marc-Hubert Nicole for useful comments.

## 2. SIEGEL HERMITIAN VARIETIES

Recall that we have fixed the CM extension  $E/F$  of number fields with the Galois involution  $c$ , such that every  $p$ -adic place of  $F$  splits in  $E$ .

**2.1. Running notation.** We introduce two sets of more specialized notation that will be used throughout the main part of the article.

(F1) We denote by

- $\mathbf{V}_F$  and  $\mathbf{V}_F^{\text{fin}}$  the set of all places and non-archimedean places of  $F$ , respectively;
- $\mathbf{V}_F^{\text{spl}}$ ,  $\mathbf{V}_F^{\text{int}}$  and  $\mathbf{V}_F^{\text{ram}}$  the subsets of  $\mathbf{V}_F^{\text{fin}}$  of those that are split, inert and ramified in  $E$ , respectively;
- $\mathbf{V}_F^{(\diamond)}$  the subset of  $\mathbf{V}_F$  of places above a finite set  $\diamond$  of places of  $\mathbb{Q}$ .

Moreover,

- for every  $v \in \mathbf{V}_F$ , we put  $E_v := E \otimes_F F_v$ ;
- for every  $v \in \mathbf{V}_F^{\text{fin}}$ , we denote by  $p_v$  the underlying rational prime of  $v$  and by  $\mathfrak{p}_v$  the maximal ideal of  $\mathcal{O}_{F_v}$ , put  $q_v := |\mathcal{O}_{F_v}/\mathfrak{p}_v|$  which is a power of  $p_v$ , and let  $d_v \geq 0$  be the integer such that  $\mathfrak{p}_v^{d_v}$  generates the different ideal of  $F_v/\mathbb{Q}_{p_v}$ .

(F2) For every  $v \in \mathbf{V}_F^{(p)}$ , let  $\mathbf{P}_v$  be the set of places of  $E$  above  $v$ . Put  $\mathbf{P} := \bigcup_{v \in \mathbf{V}_F^{(p)}} \mathbf{P}_v$ . We fix a subset  $\mathbf{P}_{\text{CM}}$  of  $\mathbf{P}$  satisfying that  $\mathbf{P}_{\text{CM}} \cap \mathbf{P}_v$  is a singleton for every  $v \in \mathbf{V}_F^{(p)}$ .

(F3) Let  $m \geq 0$  be an integer.

- We denote by  $\text{Herm}_m$  the subscheme of  $\text{Res}_{\mathcal{O}_E/\mathcal{O}_F} \text{Mat}_{m,m}$  of  $m$ -by- $m$  matrices  $b$  satisfying  ${}^t b^c = b$ . Put  $\text{Herm}_m^{\circ} := \text{Herm}_m \cap \text{Res}_{\mathcal{O}_E/\mathcal{O}_F} \text{GL}_m$ .
- For every (ordered) partition  $m = m_1 + \cdots + m_s$  with  $m_i$  a positive integer, we denote by

$$\partial_{m_1, \dots, m_s}: \text{Herm}_m \rightarrow \text{Herm}_{m_1} \times \cdots \times \text{Herm}_{m_s}$$

the morphism that extracts the diagonal blocks with corresponding ranks.

- We denote by  $\text{Herm}_m(F)^+$  (resp.  $\text{Herm}_m^{\circ}(F)^+$ ) the subset of  $\text{Herm}_m(F)$  of elements that are totally semi-positive definite (resp. totally positive definite).

(F4) Let  $\eta_{E/F}: F^{\times} \setminus \mathbb{A}_F^{\times} \rightarrow \mathbb{C}^{\times}$  be the quadratic character associated with  $E/F$ . For every finite character  $\chi: F^{\times} \setminus \mathbb{A}_F^{\times} \rightarrow \mathbb{C}^{\times}$  and every integer  $m \geq 1$ , we put

- for every  $v \in \mathbf{V}_F$ ,

$$b_{m,v}(\chi) := \prod_{i=1}^m L(i, \chi_v \eta_{E/F,v}^{m-i}).$$



- for a finite set  $\diamond$  of places of  $\mathbb{Q}$ ,

$$b_{m,\diamond}(\chi) := \prod_{v \in \mathbb{V}_F^{(\diamond)}} b_{m,v}(\chi), \quad b_m^\diamond(\chi) := \prod_{v \in \mathbb{V}_F \setminus \mathbb{V}_F^{(\diamond)}} b_{m,v}(\chi),$$

in which the second is absolutely convergent when  $m$  is even or  $\chi \neq \mathbf{1}$ .

Let  $m \geq 1$  be an integer. We equip  $W_m = E^{2m}$  and  $\bar{W}_m = E^{2m}$  with the skew-hermitian forms (that are  $E$ -linear in the first variable) given by the matrices  $w_m$  and  $-w_m$ , respectively.

- (G1) Let  $G_m$  be the unitary group of both  $W_m$  and  $\bar{W}_m$ . We write elements of  $W_m$  and  $\bar{W}_m$  in the row form, on which  $G_m$  acts from the right. Denote by  $\dagger$  the involution of  $G_m$  given by the conjugation by the element  $\begin{pmatrix} 1_m & \\ & -1_m \end{pmatrix}$  inside  $\text{Res}_{E/F} \text{GL}_{2m}$ .
- (G2) We denote by  $\{e_1, \dots, e_{2m}\}$  and  $\{\bar{e}_1, \dots, \bar{e}_{2m}\}$  the natural bases of  $W_m$  and  $\bar{W}_m$ , respectively.
- (G3) Let  $P_m \subseteq G_m$  be the parabolic subgroup stabilizing the subspace generated by  $\{e_{m+1}, \dots, e_{2m}\}$ , and  $N_m \subseteq P_m$  its unipotent radical.
- (G4) We have

- a homomorphism  $m: \text{Res}_{E/F} \text{GL}_m \rightarrow P_m$  sending  $a$  to

$$m(a) := \begin{pmatrix} a & \\ & {}_t a^{c,-1} \end{pmatrix},$$

which identifies  $\text{Res}_{E/F} \text{GL}_m$  as a Levi factor of  $P_m$ .

- a homomorphism  $n: \text{Herm}_m \rightarrow N_m$  sending  $b$  to

$$n(b) := \begin{pmatrix} 1_m & b \\ & 1_m \end{pmatrix},$$

which is an isomorphism.

- (G5) We define a maximal compact subgroup  $K_m = \prod_{v \in \mathbb{V}_F} K_{m,v}$  of  $G_m(\mathbb{A}_F)$  in the following way:
- for  $v \in \mathbb{V}_F^{\text{fin}}$ ,  $K_{m,v}$  is the stabilizer of the lattice  $\mathcal{O}_{E_v}^{2m}$ ;
  - for  $v \in \mathbb{V}_F^{(\infty)}$ ,  $K_{m,v}$  is the subgroup of the form

$$[k_1, k_2] := \frac{1}{2} \begin{pmatrix} k_1 + k_2 & -ik_1 + ik_2 \\ ik_1 - ik_2 & k_1 + k_2 \end{pmatrix},$$

in which  $k_i \in \text{GL}_m(\mathbb{C})$  satisfies  $k_i \overline{{}_t k_i} = 1_m$  for  $i = 1, 2$ .<sup>3</sup>

Moreover,

- for every place  $w$  of  $\mathbb{Q}$ , put  $K_{m,w} := \prod_{v \in \mathbb{V}_F^{(w)}} K_{m,v}$ ;
  - for a set  $\diamond$  of places of  $\mathbb{Q}$ , put  $K_m^\diamond := \prod_{w \notin \diamond} K_{m,w}$ .
- (G6) For every  $v \in \mathbb{V}_F^{(\infty)}$ , we have a character  $\kappa_{m,v}: K_{m,v} \rightarrow \mathbb{C}^\times$  that sends  $[k_1, k_2]$  to  $\det k_1 / \det k_2$ .
- (G7) For every  $v \in \mathbb{V}_F$ , we define a Haar measure  $dg_v$  on  $G_m(F_v)$  as follows:
- for  $v \in \mathbb{V}_F^{\text{fin}}$ ,  $dg_v$  is the Haar measure under which  $K_{m,v}$  has volume 1;
  - for  $v \in \mathbb{V}_F^{(\infty)}$ ,  $dg_v$  is the product of the measure on  $K_{m,v}$  of total volume 1 and the standard hyperbolic measure on  $G_m(F_v)/K_{m,v}$  (see, for example, [EL, Section 2.1]).

Put  $dg = \prod_v dg_v$ , which is a Haar measure on  $G_m(\mathbb{A}_F)$ .

- (G8) Let  $m_1, \dots, m_s$  be finitely many positive integers. Put

$$G_{m_1, \dots, m_s} := G_{m_1} \times \cdots \times G_{m_s}.$$

We denote by  $\mathcal{A}_{m_1, \dots, m_s}$  the space of both  $\mathcal{Z}(\mathfrak{g}_{m_1, \dots, m_s, \infty})$ -finite and  $K_{m_1, \infty} \times \cdots \times K_{m_s, \infty}$ -finite automorphic forms on  $G_{m_1, \dots, m_s}(\mathbb{A}_F)$ , where  $\mathcal{Z}(\mathfrak{g}_{m_1, \dots, m_s, \infty})$  denotes the center of the complexified universal enveloping algebra of the Lie algebra  $\mathfrak{g}_{m_1, \dots, m_s, \infty}$  of  $G_{m_1, \dots, m_s} \otimes_{\mathbb{Q}} \mathbb{R}$ . For every integer  $w \geq 0$  (as weight), we denote by

- $\mathcal{A}_{m_1, \dots, m_s}^{[w]}$  the maximal subspace of  $\mathcal{A}_{m_1, \dots, m_s}$  on which for every  $v \in \mathbb{V}_F^{(\infty)}$  and every  $1 \leq j \leq s$ ,  $K_{m_j, v}$  acts by the character  $\kappa_{m_j, v}^w$ ,

<sup>3</sup>Here, we choose a complex embedding of  $E$  above  $v$  to identify  $G_m(F_v)$  as a subgroup of  $\text{GL}_{2m}(\mathbb{C})$ . However, neither  $K_{m,v}$  nor the character  $\kappa_{m,v}$  in (G6) depends on such a choice.

- $\mathcal{A}_{m_1, \dots, m_s, \text{hol}}^{[w]} \subseteq \mathcal{A}_{m_1, \dots, m_s}^{[w]}$  the subspace of holomorphic ones.
- (G9) For every vector space  $\mathcal{H}$  on which  $G_{m_1, \dots, m_s}(\mathbb{A}_F^\infty)$  acts, we put  $\mathcal{H}(K) := \mathcal{H}^K$  for every open compact subgroup  $K \subseteq G_{m_1, \dots, m_s}(\mathbb{A}_F^\infty)$ .

**2.2. Siegel hermitian varieties and automorphy line bundles.** We first recall the construction of a CM moduli problem following [LTX<sup>+</sup>22, Section 3.5]. Let  $T$  be the subtorus of  $\text{Res}_{E/\mathbb{Q}} \mathbf{G}$  that is the inverse image of  $\mathbf{G}_{\mathbb{Q}}$  under the norm map  $\text{Nm}_{E/F}: \text{Res}_{E/\mathbb{Q}} \mathbf{G} \rightarrow \text{Res}_{F/\mathbb{Q}} \mathbf{G}$ .

For every nonzero element  $\delta \in E^{c=-1}$ , we denote by  $W^\delta$  the  $E$ -vector space  $E$  (itself) together with a pairing  $\langle \cdot, \cdot \rangle^\delta: E \times E \rightarrow \mathbb{Q}$  given by  $\langle x, y \rangle^\delta = \text{Tr}_{E/\mathbb{Q}}(\delta xy^c)$ . For every  $\mathbb{Q}$ -ring  $R$ , we have  $T(R) = \{t \in (E \otimes_{\mathbb{Q}} R)^\times \mid \langle tx, ty \rangle^\delta = c(t)\langle x, y \rangle^\delta \text{ for some } c(t) \in R^\times\}$ .

For every neat open compact subgroup  $K_T$  of  $T(\mathbb{A}^\infty)$ , we define a moduli problem  $\Sigma^\delta(K_T)$  on  $\text{Sch}'_{\mathbb{Q}_p}$  as follows: for every  $S \in \text{Sch}'_{\mathbb{Q}_p}$ ,  $\Sigma^\delta(K_T)(S)$  is the set of equivalence classes of quadruples  $(A_0, i_0, \lambda_0, \eta_0)$  in which

- $A_0$  is an abelian scheme over  $S$  of relative dimension  $[F : \mathbb{Q}]$ ,
- $i_0: E \rightarrow \text{End}_S(A_0) \otimes \mathbb{Q}$  is an  $E$ -action such that for every  $x \in E$ ,  $\text{tr}(i_0(x) \mid \text{Lie}_S(A_0)) = \sum_{u \in \text{P}_{\text{CM}}} \text{Tr}_{E_u/\mathbb{Q}_p}(x)$ , where  $\text{P}_{\text{CM}}$  is the fixed subset of  $\text{P}$  (§2.1(F2)),
- $\lambda_0: A_0 \rightarrow A_0^\vee$  is a quasi-polarization under which the Rosati involution coincides with the complex conjugation on  $E$  under  $i_0$ ,
- $\eta_0: W^\delta \otimes_{\mathbb{Q}} \mathbb{A}^\infty \rightarrow \text{H}_1^{\text{ét}}(A_0, \mathbb{A}^\infty)$  is an  $K_T$ -level structure.

It is known that  $\Sigma^\delta(K_T)$  is a nonempty scheme finite étale over  $\mathbb{Q}_p$ , which admits a natural action by the finite group  $T(\mathbb{A}^\infty)/T(\mathbb{Q})K_T$  such that each orbit is Galois over  $\text{Spec } \mathbb{Q}_p$  with the Galois group  $T(\mathbb{A}^\infty)/T(\mathbb{Q})K_T$ . We fix such an orbit  $\Sigma_0^\delta(K_T)$ .

For every neat open compact subgroup  $K \subseteq G_m(\mathbb{A}_F^\infty)$ , we consider the moduli problem  $\Sigma_m^\delta(K, K_T)$  on  $\text{Sch}'_{\mathbb{Q}_p}$  as follows: for every  $S \in \text{Sch}'_{\mathbb{Q}_p}$ ,  $\Sigma_m^\delta(K, K_T)(S)$  is the set of equivalent classes of octuples  $(A_0, i_0, \lambda_0, \eta_0; A, i, \lambda, \eta)$  in which

- $(A_0, i_0, \lambda_0, \eta_0)$  is an element of  $\Sigma_0^\delta(K_T)(S)$ ,
- $A$  is an abelian scheme over  $S$  of relative dimension  $2m[F : \mathbb{Q}]$ ,
- $i: E \rightarrow \text{End}_S(A) \otimes \mathbb{Q}$  is an  $E$ -action such that for every  $x \in E$ ,  $\text{tr}(i(x) \mid \text{Lie}_S(A)) = m \text{Tr}_{E/\mathbb{Q}}(x)$ ,
- $\lambda: A \rightarrow A^\vee$  is a quasi-polarization under which the Rosati involution coincides with the complex conjugation on  $E$  under  $i$ ,
- $\eta: W_m^\delta \otimes_E \mathbb{A}_E^\infty \rightarrow \text{Hom}_{\mathbb{A}_E^\infty}(\text{H}_1^{\text{ét}}(A_0, \mathbb{A}^\infty), \text{H}_1^{\text{ét}}(A, \mathbb{A}^\infty))$  is a  $K$ -level structure, where  $W_m^\delta$  denotes the space  $E^{2m}$  equipped with the hermitian form  $\delta^{-1} \cdot \mathbf{w}_m$ .

It is known that  $\Sigma_m^\delta(K, K_T)$  is a scheme finite type over  $\Sigma_0^\delta(K_T)$ , which admits a natural lift of the action of  $T(\mathbb{A}^\infty)/T(\mathbb{Q})K_T$ . We denote by  $\Sigma_m^\delta(K, K_T)^b$  the quotient of  $\Sigma_m^\delta(K, K_T)$  by  $T(\mathbb{A}^\infty)/T(\mathbb{Q})K_T$ , as a presheaf on  $\text{Sch}'_{\mathbb{Q}_p}$ .

Now we discuss the relation between  $\Sigma_m^\delta(K, K_T)^b$  and usual Shimura varieties. For every CM type  $\Phi$ , we have the Deligne homomorphism

$$\begin{aligned} \mathbf{h}_m^\Phi: \text{Res}_{\mathbb{C}/\mathbb{R}} \mathbf{G} &\rightarrow (\text{Res}_{F/\mathbb{Q}} G_m) \otimes_{\mathbb{Q}} \mathbb{R} \\ z &\rightarrow ([1_m, (\bar{z}/z)1_m], \dots, [1_m, (\bar{z}/z)1_m]) \in K_{m, \infty}, \end{aligned}$$

in which for every archimedean place  $v$  of  $F$ , the notation  $[1_m, (\bar{z}/z)1_m]$  is understood via the unique complex embedding of  $E$  in  $\Phi$  inducing  $v$ . Then we obtain a projective system of Shimura varieties  $\{\Sigma_m^\Phi(K)\}_K$  associated with the Shimura data  $(\text{Res}_{F/\mathbb{Q}} G_m, \mathbf{h}_m^\Phi)$  indexed by neat open compact subgroups  $K \subseteq G_m(\mathbb{A}_F^\infty)$ , which are smooth quasi-projective complex schemes of dimension  $m^2[F : \mathbb{Q}]$ , with the complex analytification

$$\Sigma_m^\Phi(K)^{\text{an}} = G_m(F) \backslash G_m(\mathbb{A}_F) / K_{m, \infty} K.$$

For every embedding  $\iota: \mathbb{Q}_p \rightarrow \mathbb{C}$ , we denote by  $\Phi_\iota$  the set of complex embeddings  $i: E \rightarrow \mathbb{C}$  such that the  $p$ -adic place induced by the embedding  $i: E \hookrightarrow i(E) \cdot \iota(\mathbb{Q}_p)$  belongs to  $\text{P}_{\text{CM}}$  (§2.1(F2)). Then  $\Phi_\iota$  is a CM type of  $E$ .

**Lemma 2.1.** *The presheaf  $\Sigma_m^\delta(K, K_T)^b$  is a scheme over  $\mathbb{Q}_p$  independent of the choices of  $K_T$ ,  $\delta$ , and the orbit  $\Sigma_0^\delta(K_T)$ . Moreover, for every embedding  $\iota: \mathbb{Q}_p \rightarrow \mathbb{C}$ , we have a canonical isomorphism*

$$\Sigma_m^\delta(K, K_T)^b \otimes_{\mathbb{Q}_p, \iota} \mathbb{C} \xrightarrow{\sim} \Sigma_m^{\Phi_\iota}(K).$$

*Proof.* By definition, the reflex field  $E_{\Phi_t} \subseteq \mathbb{C}$  of  $\Phi_t$  is contained in  $\iota(\mathbb{Q}_p)$ . Then there is a canonical isomorphism

$$\left( X_K \otimes_{E_{\Phi_t}} Y_{K_T} \right) \otimes_{E_{\Phi_t, t^{-1}}} \mathbb{Q}_p \simeq \Sigma_m^\delta(K, K_T)$$

of schemes over  $\mathbb{Q}_p$ , where  $X_K$  and  $Y_{K_T}$  are the usual Shimura varieties for  $G_m$  and  $T$  of level  $K$  and  $K_T$ , respectively, over their common reflex field  $E_{\Phi_t}$ . Under such isomorphism,  $T(\mathbb{A}^\infty)/T(\mathbb{Q})K_T$  acts on the left side via the second factor  $Y_{K_T}$  whose quotient is nothing but  $\text{Spec } E_{\Phi_t}$ . Thus, we obtain a canonical isomorphism  $X_K \otimes_{E_{\Phi_t, t^{-1}}} \mathbb{Q}_p \simeq \Sigma_m^\delta(K, K_T)^b$ . The lemma follows.  $\square$

**Definition 2.2.** We define the Siegel hermitian variety (of genus  $m$  and level  $K$ ) over  $\mathbb{Q}_p$ , denoted as  $\Sigma_m(K)$ , to be  $\Sigma_m^\delta(K, K_T)^b$ , which makes sense by the lemma above.<sup>4</sup>

Now we define the *automorphy line bundle* on  $\Sigma_m(K)$ . Denote by  $A$  the ( $A$  part) of the universal object over  $\Sigma_m^\delta(K, K_T)$ . Then  $\text{Lie}(A)$  is a projective  $\mathcal{O}_{\mathbb{Q}} E$ -module of rank  $m$ , where  $\mathcal{O} = \mathcal{O}_{\Sigma_m^\delta(K, K_T)}$  is the structure sheaf. Put

$$\omega_m^\delta := \det_{\mathcal{O}} \left( \det_{\mathcal{O} \otimes_{\mathbb{Q}} E} \text{Lie}(A)^\vee \right),$$

which is a line bundle on  $\Sigma_m^\delta(K, K_T)$ . Since  $T(\mathbb{A}^\infty)/T(\mathbb{Q})K_T$  acts trivially on  $\omega_m^\delta$ ,  $\omega_m^\delta$  descends to a line bundle  $\omega_m$  on  $\Sigma_m(K)$ . It is easy too see that  $\omega_m$  does not depend on the choices of  $K_T$ ,  $\delta$ , and the orbit  $\Sigma_0^\delta(K_T)$ .

Now suppose that we are given a partition  $m = m_1 + \cdots + m_s$  of  $m$  into positive integers. We have a natural isometry

$$(2.1) \quad W_{m_1} \oplus \cdots \oplus W_{m_s} \simeq W_m$$

such that if we write  $\{e_1^j, \dots, e_{2m_j}^j\}$  as the standard bases for  $W_{m_j}$  for  $1 \leq j \leq s$ , then the standard basis of  $W_m$  is identified with  $\{e_1^1, \dots, e_{m_1}^1, \dots, e_1^s, \dots, e_{m_s}^s, e_{m_1+1}^1, \dots, e_{2m_1}^1, \dots, e_{m_s+1}^s, \dots, e_{2m_s}^s\}$ . In particular, we may regard  $G_{m_1, \dots, m_s} = G_{m_1} \times \cdots \times G_{m_s}$  as a subgroup of  $G_m$ . We obtain a map

$$(2.2) \quad \rho_{m_1, \dots, m_s} : \mathcal{A}_{m, \text{hol}}^{[w]} \rightarrow \mathcal{A}_{m_1, \dots, m_s, \text{hol}}^{[w]}$$

(see §2.1(G8)) given by the restriction to the subgroup  $G_{m_1, \dots, m_s}(\mathbb{A}_F)$ .

For neat open compact subgroups  $K_j \subseteq G_{m_j}(\mathbb{A}_F^\infty)$  for  $1 \leq j \leq s$ , we put

$$\begin{aligned} \Sigma_{m_1, \dots, m_s}^\delta(K_1 \times \cdots \times K_s, K_T) &:= \Sigma_{m_1}^\delta(K_1, K_T) \times_{\Sigma_0^\delta(K_T)} \cdots \times_{\Sigma_0^\delta(K_T)} \Sigma_{m_s}^\delta(K_s, K_T), \\ \omega_{m_1, \dots, m_s}^\delta &:= \omega_{m_1}^\delta \boxtimes \cdots \boxtimes \omega_{m_s}^\delta; \end{aligned}$$

and

$$\begin{aligned} \Sigma_{m_1, \dots, m_s}(K_1 \times \cdots \times K_s) &:= \Sigma_{m_1}(K_1) \times_{\mathbb{Q}_p} \cdots \times_{\mathbb{Q}_p} \Sigma_{m_s}(K_s), \\ \omega_{m_1, \dots, m_s} &:= \omega_{m_1} \boxtimes \cdots \boxtimes \omega_{m_s}. \end{aligned}$$

We have the natural quotient map

$$\xi_{m_1, \dots, m_s} : \Sigma_{m_1, \dots, m_s}^\delta(K_1 \times \cdots \times K_s, K_T) \rightarrow \Sigma_{m_1, \dots, m_s}(K_1 \times \cdots \times K_s)$$

under which  $\xi_{m_1, \dots, m_s}^* \omega_{m_1, \dots, m_s} \simeq \omega_{m_1, \dots, m_s}^\delta$ .

For a neat open compact subgroup  $K \subseteq G_m(\mathbb{A}_F^\infty)$  containing  $K_1 \times \cdots \times K_s$ , there is a natural morphism

$$\sigma_{m_1, \dots, m_s}^\delta : \Sigma_{m_1, \dots, m_s}^\delta(K_1 \times \cdots \times K_s, K_T) \rightarrow \Sigma_m^\delta(K, K_T)$$

sending  $((A_0, i_0, \lambda_0, \eta_0; A_j, i_j, \lambda_j, \eta_j))_{1 \leq j \leq s}$  to

$$(A_0, i_0, \lambda_0, \eta_0; A_1 \times \cdots \times A_s, (i_1, \dots, i_s), \lambda_1 \times \cdots \times \lambda_s, (\eta_1, \dots, \eta_s)).$$

It is clear that  $\sigma_{m_1, \dots, m_s}^\delta$  descends to morphism

$$\sigma_{m_1, \dots, m_s} : \Sigma_{m_1, \dots, m_s}(K_1 \times \cdots \times K_s) \rightarrow \Sigma_m(K)$$

<sup>4</sup>By construction,  $\Sigma_m(K)$  also depends on the choice of the subset  $P_{\text{CM}}$  of  $P$  (§2.1(F2)).

rendering the following diagram

$$(2.3) \quad \begin{array}{ccc} \Sigma_{m_1, \dots, m_s}^\delta(K_1 \times \cdots \times K_s, K_T) & \xrightarrow{\sigma_{m_1, \dots, m_s}^\delta} & \Sigma_m^\delta(K, K_T) \\ \xi_{m_1, \dots, m_s} \downarrow & & \downarrow \xi_m \\ \Sigma_{m_1, \dots, m_s}(K_1 \times \cdots \times K_s) & \xrightarrow{\sigma_{m_1, \dots, m_s}} & \Sigma_m(K) \end{array}$$

in  $\text{Sch}'_{/\mathbb{Q}_p}$  commutative. It is independent of the choices of  $K_T$ ,  $\delta$ , and the orbit  $\Sigma_0^\delta(K_T)$ . For the automorphy line bundles, we have  $(\sigma_{m_1, \dots, m_s}^\delta)^* \omega_m^\delta \simeq \omega_{m_1, \dots, m_s}^\delta$ , and hence  $\sigma_{m_1, \dots, m_s}^* \omega_m \simeq \omega_{m_1, \dots, m_s}$ .

For every integer  $w \geq 0$ , put

$$\begin{aligned} \mathcal{H}_{m_1, \dots, m_s}^w(K_1 \times \cdots \times K_s) &:= H^0(\Sigma_{m_1, \dots, m_s}(K_1 \times \cdots \times K_s), \omega_{m_1, \dots, m_s}^{\otimes w}), \\ \mathcal{H}_{m_1, \dots, m_s}^w &:= \varinjlim_{K_1, \dots, K_s} \mathcal{H}_{m_1, \dots, m_s}^w(K_1 \times \cdots \times K_s). \end{aligned}$$

For every embedding  $\iota: \mathbb{Q}_p \rightarrow \mathbb{C}$ , we have an injective map

$$(2.4) \quad h_{m_1, \dots, m_s}^\iota: \mathcal{A}_{m_1, \dots, m_s, \text{hol}}^{[w]} \rightarrow \mathcal{H}_{m_1, \dots, m_s}^w \otimes_{\mathbb{Q}_p, \iota} \mathbb{C},$$

which fits into the following commutative diagram

$$\begin{array}{ccc} \mathcal{A}_{m, \text{hol}}^{[w]} & \xrightarrow[\text{(2.2)}]{\rho_{m_1, \dots, m_s}} & \mathcal{A}_{m_1, \dots, m_s, \text{hol}}^{[w]} \\ h_m^\iota \downarrow & & \downarrow h_{m_1, \dots, m_s}^\iota \\ \mathcal{H}_m^w \otimes_{\mathbb{Q}_p, \iota} \mathbb{C} & \xrightarrow{\sigma_{m_1, \dots, m_s}^*} & \mathcal{H}_{m_1, \dots, m_s}^w \otimes_{\mathbb{Q}_p, \iota} \mathbb{C} \end{array}$$

of complex vector spaces.

**Definition 2.3.** Let the notation be as above.

- (1) We define  $\mathcal{H}_{m_1, \dots, m_s}^{[w]}$  to be the maximal subspace of  $\mathcal{H}_{m_1, \dots, m_s}^w$  such that for every embedding  $\iota: \mathbb{Q}_p \rightarrow \mathbb{C}$ ,  $\mathcal{H}_{m_1, \dots, m_s}^{[w]} \otimes_{\mathbb{Q}_p, \iota} \mathbb{C}$  is contained in the image of  $\mathcal{A}_{m_1, \dots, m_s, \text{hol}}^{[w]}$  under  $h_{m_1, \dots, m_s}^\iota$ .
- (2) For every  $\varphi \in \mathcal{H}_{m_1, \dots, m_s}^{[w]}$  and every embedding  $\iota: \mathbb{Q}_p \rightarrow \mathbb{C}$ , we denote by  $\varphi^\iota$  the unique element in  $\mathcal{A}_{m_1, \dots, m_s, \text{hol}}^{[w]}$  such that  $h_{m_1, \dots, m_s}^\iota(\varphi^\iota) = \iota\varphi$ .

*Remark 2.4.* We have the following remarks concerning  $\mathcal{H}_{m_1, \dots, m_s}^{[w]}$ .

- (1) The inclusion  $\mathcal{H}_{m_1, \dots, m_s}^{[w]} \subseteq \mathcal{H}_{m_1, \dots, m_s}^w$  is proper in general since in the definition of  $\mathcal{H}_{m_1, \dots, m_s}^w$ , we do not impose any growth condition along the boundary.
- (2) It is clear that the subspace  $\mathcal{H}_{m_1, \dots, m_s}^{[w]}$  is closed under the action of  $G_{m_1, \dots, m_s}(\mathbb{A}_F^\infty)$ . Moreover, in its definition, it suffices to check for *some* embedding  $\iota$ .
- (3) The natural map  $\mathcal{H}_{m_1}^{[w]} \otimes_{\mathbb{Q}_p} \cdots \otimes_{\mathbb{Q}_p} \mathcal{H}_{m_s}^{[w]} \rightarrow \mathcal{H}_{m_1, \dots, m_s}^{[w]}$  given by exterior product is an isomorphism. Indeed, it suffices to check it at every finite level, which is then an isomorphism of *finite-dimensional*  $\mathbb{Q}_p$ -vector spaces.

To end this subsection, we review the notion of analytic  $q$ -expansion (or Siegel–Fourier expansion).

**Definition 2.5.** For every ring  $R$ , we denote by  $\text{SF}_{m_1, \dots, m_s}(R)$  the  $R$ -module of formal power series

$$\sum_{(T_1, \dots, T_s) \in \text{Herm}_{m_1}(F)^+ \times \cdots \times \text{Herm}_{m_s}(F)^+} a_{T_1, \dots, T_s} q^{T_1, \dots, T_s}, \quad a_{T_1, \dots, T_s} \in R$$

in which  $a_{T_1, \dots, T_s}$  vanishes unless the entries of  $T_1, \dots, T_s$  are in a certain fractional ideal of  $E$ . We have a restriction map

$$\varrho_{m_1, \dots, m_s}: \text{SF}_m(R) \rightarrow \text{SF}_{m_1, \dots, m_s}(R)$$

sending

$$\sum_{T \in \text{Herm}_m(F)^+} a_T q^T$$

to

$$\sum_{(T_1, \dots, T_s) \in \text{Herm}_{m_1}(F)^+ \times \dots \times \text{Herm}_{m_s}(F)^+} \left( \sum_{\substack{T \in \text{Herm}_m(F)^+ \\ \partial_{m_1, \dots, m_s} T = (T_1, \dots, T_s)}} a_T \right) q^{T_1, \dots, T_s}.$$

It is an easy exercise to show that the interior summation is always a finite sum.

For every integer  $w \geq 0$ , we have a map

$$(2.5) \quad \mathbf{q}_{m_1, \dots, m_s}^{\text{an}} : \mathcal{A}_{m_1, \dots, m_s, \text{hol}}^{[w]} \rightarrow \text{SF}_{m_1, \dots, m_s}(\mathbb{C})$$

$$\varphi \mapsto \sum_{(T_1, \dots, T_s) \in \text{Herm}_{m_1}(F)^+ \times \dots \times \text{Herm}_{m_s}(F)^+} a_{T_1, \dots, T_s}(\varphi) q^{T_1, \dots, T_s}$$

in which  $a_{T_1, \dots, T_s}(\varphi)$  equals

$$\int_{\text{Herm}_{m_1}(F) \backslash \text{Herm}_{m_1}(\mathbb{A}_F)} \dots \int_{\text{Herm}_{m_s}(F) \backslash \text{Herm}_{m_s}(\mathbb{A}_F)} \varphi(n(b_1), \dots, n(b_s)) \psi_F(\text{tr } T_1 b_1)^{-1} \dots \psi_F(\text{tr } T_s b_s)^{-1} db_1 \dots db_s$$

with  $db_1, \dots, db_s$  being Tamagawa measures.

We have the following commutative diagram

$$\begin{array}{ccc} \mathcal{A}_{m, \text{hol}}^{[w]} & \xrightarrow[\text{(2.2)}]{\rho_{m_1, \dots, m_s}} & \mathcal{A}_{m_1, \dots, m_s, \text{hol}}^{[w]} \\ \mathbf{q}_m^{\text{an}} \downarrow & & \downarrow \mathbf{q}_{m_1, \dots, m_s}^{\text{an}} \\ \text{SF}_m(\mathbb{C}) & \xrightarrow{\varrho_{m_1, \dots, m_s}} & \text{SF}_{m_1, \dots, m_s}(\mathbb{C}) \end{array}$$

under restrictions.

**2.3. Relation with PEL type moduli spaces.** In order to show the rationality of some Eisenstein series later, we need the theory of algebraic  $q$ -expansions. However, since such theory was only developed for PEL type shimura variety (in the sense of Kottwitz), we need to study its relation with our Siegel hermitian varieties.

Let  $\widetilde{W}_m$  be the space  $E^{2m}$  equipped with the pairing  $\text{Tr}_{E/\mathbb{Q}} \circ \langle \cdot, \cdot \rangle_{W_{2m}} : E^{2m} \times E^{2m} \rightarrow \mathbb{Q}$ . Let  $\widetilde{G}_m$  be the similitude group of  $\widetilde{W}_m$ , which is a reductive group over  $\mathbb{Q}$ . Let  $\widetilde{P}_m \subseteq \widetilde{G}_m$  be the parabolic subgroup stabilizing the subspace generated by  $\{e_{m+1}, \dots, e_{2m}\}$ ,

Consider a partition  $m = m_1 + \dots + m_s$  of  $m$  into positive integers. We denote by  $\widetilde{G}_{m_1, \dots, m_s}$  the subgroup of  $\widetilde{G}_{m_1} \times \dots \times \widetilde{G}_{m_s}$  of common similitudes; in other words, it fits into a Cartesian diagram

$$\begin{array}{ccc} \widetilde{G}_{m_1, \dots, m_s} & \longrightarrow & \widetilde{G}_{m_1} \times \dots \times \widetilde{G}_{m_s} \\ \downarrow & & \downarrow \\ \mathbf{G}_{\mathbb{Q}} & \xrightarrow{\text{diagonal}} & \mathbf{G}_{\mathbb{Q}}^s \end{array}$$

in which the vertical arrows are similitude maps. In particular, we may regard  $\widetilde{G}_{m_1, \dots, m_s}$  as a subgroup of  $\widetilde{G}_m$ . Put  $\widetilde{P}_{m_1, \dots, m_s} := \widetilde{G}_{m_1, \dots, m_s} \cap \widetilde{P}_m$ .

For every neat open compact subgroup  $\widetilde{K}_{m_1, \dots, m_s} \subseteq \widetilde{G}_{m_1, \dots, m_s}(\mathbb{A}^{\infty})$ , we consider the PEL type moduli problem  $\widetilde{\Sigma}_{m_1, \dots, m_s}(\widetilde{K}_{m_1, \dots, m_s})$  on  $\text{Sch}'_{\mathbb{Q}}$  as follows: for every  $S \in \text{Sch}'_{\mathbb{Q}}$ ,  $\widetilde{\Sigma}_{m_1, \dots, m_s}(\widetilde{K}_{m_1, \dots, m_s})(S)$  is the set of equivalence classes of  $s$ -tuples of quadruples  $((A_1, i_1, \lambda_1, \widetilde{\eta}_1), \dots, (A_s, i_s, \lambda_s, \widetilde{\eta}_s))$  in which

- for  $1 \leq j \leq s$ ,  $A_j$  is an abelian scheme over  $S$  of relative dimension  $2m_j[F : \mathbb{Q}]$ ,
- for  $1 \leq j \leq s$ ,  $i_j : E \rightarrow \text{End}_S(A_j) \otimes \mathbb{Q}$  is an  $E$ -action such that for every  $x \in E$ ,  $\text{tr}(i_j(x) | \text{Lie}_S(A_j)) = m_j \text{Tr}_{E/\mathbb{Q}}(x)$ ,
- for  $1 \leq j \leq s$ ,  $\lambda_j : A_j \rightarrow A_j^{\vee}$  is a quasi-polarization under which the Rosati involution coincides with the complex conjugation on  $E$  under  $i_j$ ,
- $\{\widetilde{\eta}_j : \widetilde{W}_m \otimes_{\mathbb{Q}} \mathbb{A}^{\infty} \rightarrow \text{H}_1^{\text{ét}}(A_j, \mathbb{A}^{\infty})\}_{1 \leq j \leq s}$  is a  $\widetilde{K}_{m_1, \dots, m_s}$ -orbit of skew-hermitian similitudes with similitude factors independent of  $j$ .

Then  $\widetilde{\Sigma}_{m_1, \dots, m_s}(\widetilde{K}_{m_1, \dots, m_s})$  is a scheme of finite type over  $\mathbb{Q}$ . Now for a neat open compact subgroup  $\widetilde{K} \subseteq \widetilde{G}_m(\mathbb{A}^\infty)$  containing  $\widetilde{K}_{m_1, \dots, m_s}$ , we have an obvious morphism

$$\widetilde{\sigma}_{m_1, \dots, m_s} : \widetilde{\Sigma}_{m_1, \dots, m_s}(\widetilde{K}_{m_1, \dots, m_s}) \rightarrow \widetilde{\Sigma}_m(\widetilde{K})$$

over  $\mathbb{Q}$  by “taking the product of all factors”. For neat open compact subgroups  $\widetilde{K}_j \subseteq \widetilde{G}_{m_j}(\mathbb{A}^\infty)$  containing the image of  $\widetilde{K}_{m_1, \dots, m_s}$  under the natural projection map  $\widetilde{G}_{m_1, \dots, m_s} \rightarrow \widetilde{G}_{m_j}$ , we have another obvious map

$$\tau_{m_1, \dots, m_s} : \widetilde{\Sigma}_{m_1, \dots, m_s}(\widetilde{K}_{m_1, \dots, m_s}) \rightarrow \widetilde{\Sigma}_{m_1}(\widetilde{K}_1) \times_{\mathbb{Q}} \cdots \times_{\mathbb{Q}} \widetilde{\Sigma}_{m_s}(\widetilde{K}_s)$$

over  $\mathbb{Q}$ . On  $\widetilde{\Sigma}_m(\widetilde{K})$ , we have the automorphy line bundle  $\widetilde{\omega}_m$  similar to  $\omega_m^\delta$ , which satisfies

$$\widetilde{\sigma}_{m_1, \dots, m_s}^* \widetilde{\omega}_m \simeq \tau_{m_1, \dots, m_s}^* (\widetilde{\omega}_{m_1} \boxtimes \cdots \boxtimes \widetilde{\omega}_{m_s}).$$

Put  $\widetilde{\omega}_{m_1, \dots, m_s} := \widetilde{\sigma}_{m_1, \dots, m_s}^* \widetilde{\omega}_m$  for future use.

**Notation 2.6.** For every  $1 \leq j \leq s$ , we have an isometry  $W_{m_j}^\delta \otimes_E W^\delta \xrightarrow{\sim} \widetilde{W}_{m_j}$ . These isometries induce a homomorphism

$$\zeta_{m_1, \dots, m_s} : \text{Res}_{F/\mathbb{Q}} G_{m_1, \dots, m_s} \times T \rightarrow \widetilde{G}_{m_1, \dots, m_s}$$

sending  $(g_1, \dots, g_s, t)$  to  $(g_1 t, \dots, g_s t)$ , which is independent of the choice of  $\delta$ . Using this map, we regard  $\text{Res}_{F/\mathbb{Q}} G_{m_1, \dots, m_s}$  as a subgroup of  $\widetilde{G}_{m_1, \dots, m_s}$  in what follows.

For neat open compact subgroups  $K_j \subseteq G_{m_j}(\mathbb{A}_F^\infty)$  for  $1 \leq j \leq s$  and  $K_T \subseteq T(\mathbb{A}^\infty)$  such that  $K_1 \times \cdots \times K_s \times K_T$  is contained in  $\widetilde{K}_{m_1, \dots, m_s}$ , we have a natural morphism

$$\zeta_{m_1, \dots, m_s} : \Sigma_{m_1, \dots, m_s}^\delta(K_1 \times \cdots \times K_s, K_T) \rightarrow \widetilde{\Sigma}_{m_1, \dots, m_s}(\widetilde{K}_{m_1, \dots, m_s}) \otimes_{\mathbb{Q}} \mathbb{Q}_p$$

sending  $((A_0, i_0, \lambda_0, \eta_0; A_j, i_j, \lambda_j, \eta_j))_{1 \leq j \leq s}$  to  $((A_j, i_j, \lambda_j, \widetilde{\eta}_j))_{1 \leq j \leq s}$ , where  $\widetilde{\eta}_j$  sends  $w \otimes v$  to  $\eta_j(w)(\eta_0(v))$ . The morphism  $\zeta_{m_1, \dots, m_s}$  is finite étale.

In summary, for every neat open compact subgroup  $K \subseteq G_m(\mathbb{A}_F^\infty)$  containing  $K_1 \times \cdots \times K_s$  and such that  $\zeta_m(K \times K_T)$  is contained in  $\widetilde{K}$ , we have a diagram

(2.6)

$$\begin{array}{ccccc} \widetilde{\Sigma}_{m_1}(\widetilde{K}_1)_{\mathbb{Q}_p} \times_{\mathbb{Q}_p} \cdots \times_{\mathbb{Q}_p} \widetilde{\Sigma}_{m_s}(\widetilde{K}_s)_{\mathbb{Q}_p} & \xleftarrow{\tau_{m_1, \dots, m_s}} & \widetilde{\Sigma}_{m_1, \dots, m_s}(\widetilde{K}_{m_1, \dots, m_s})_{\mathbb{Q}_p} & \xrightarrow{\widetilde{\sigma}_{m_1, \dots, m_s}} & \widetilde{\Sigma}_m(\widetilde{K})_{\mathbb{Q}_p} \\ \zeta_{m_1} \times \cdots \times \zeta_{m_s} \uparrow & & \zeta_{m_1, \dots, m_s} \uparrow & & \zeta_m \uparrow \\ \Sigma_{m_1}^\delta(K_1, K_T) \times_{\Sigma_0^\delta(K_T)} \cdots \times_{\Sigma_0^\delta(K_T)} \Sigma_{m_s}^\delta(K_s, K_T) & \xlongequal{\text{def}} & \Sigma_{m_1, \dots, m_s}^\delta(K_1 \times \cdots \times K_s, K_T) & \xrightarrow{\sigma_{m_1, \dots, m_s}^\delta} & \Sigma_m^\delta(K, K_T) \\ \xi_{m_1} \times \cdots \times \xi_{m_s} \downarrow & & \xi_{m_1, \dots, m_s} \downarrow & & \xi_m \downarrow \\ \Sigma_{m_1}(K_1) \times_{\mathbb{Q}_p} \cdots \times_{\mathbb{Q}_p} \Sigma_{m_s}(K_s) & \xlongequal{\text{def}} & \Sigma_{m_1, \dots, m_s}(K_1 \times \cdots \times K_s) & \xrightarrow{\sigma_{m_1, \dots, m_s}} & \Sigma_m(K) \end{array}$$

in  $\text{Sch}'_{/\mathbb{Q}_p}$  expanding (2.3) as the lower-right square, in which various automorphy line bundles are compatible under pullbacks.

Similar to  $\mathcal{A}_{m_1, \dots, m_s, \text{hol}}^{[w]}$  (§2.1(G8)), we define a space  $\widetilde{\mathcal{A}}_{m_1, \dots, m_s, \text{hol}}^{[w]}$  of certain automorphic forms on  $\widetilde{G}_{m_1, \dots, m_s}(\mathbb{A})$  with the additional requirement that  $(t1_{m_1}, \dots, t1_{m_s})$  acts trivially for every  $t \in T(\mathbb{R})$ . We have a map

$$(2.7) \quad \widetilde{\rho}_{m_1, \dots, m_s} : \widetilde{\mathcal{A}}_{m, \text{hol}}^{[w]} \rightarrow \widetilde{\mathcal{A}}_{m_1, \dots, m_s, \text{hol}}^{[w]}$$

given by the restriction to the subgroup  $\widetilde{G}_{m_1, \dots, m_s}(\mathbb{A})$ .

For every integer  $w \geq 0$ , put

$$\begin{aligned} \widetilde{\mathcal{H}}_{m_1, \dots, m_s}^w(\widetilde{K}_{m_1, \dots, m_s}) &:= H^0(\widetilde{\Sigma}_{m_1, \dots, m_s}(\widetilde{K}_{m_1, \dots, m_s}), \widetilde{\omega}_{m_1, \dots, m_s}^{\otimes w}), \\ \widetilde{\mathcal{H}}_{m_1, \dots, m_s}^w &:= \varinjlim_{\widetilde{K}_{m_1, \dots, m_s}} \widetilde{\mathcal{H}}_{m_1, \dots, m_s}^w(\widetilde{K}_{m_1, \dots, m_s}). \end{aligned}$$

**Definition 2.7.** Similar to (2.4), we have an injective map

$$\tilde{h}_{m_1, \dots, m_s} : \tilde{\mathcal{A}}_{m_1, \dots, m_s, \text{hol}}^{[w]} \rightarrow \tilde{\mathcal{H}}_{m_1, \dots, m_s}^w \otimes_{\mathbb{Q}} \mathbb{C}$$

for  $w \geq 0$ . We define  $\tilde{\mathcal{H}}_{m_1, \dots, m_s}^{[w]}$  to be the subspace of  $\tilde{\mathcal{H}}_{m_1, \dots, m_s}^w$  such that the image of  $\tilde{h}_{m_1, \dots, m_s}$  coincides with  $\tilde{\mathcal{H}}_{m_1, \dots, m_s}^{[w]} \otimes_{\mathbb{Q}} \mathbb{C}$ . Thus, we obtain an isomorphism

$$(2.8) \quad \tilde{h}_{m_1, \dots, m_s} : \tilde{\mathcal{A}}_{m_1, \dots, m_s, \text{hol}}^{[w]} \xrightarrow{\sim} \tilde{\mathcal{H}}_{m_1, \dots, m_s}^{[w]} \otimes_{\mathbb{Q}} \mathbb{C}.$$

Now we review the algebraic theory of  $q$ -expansion for  $\tilde{\Sigma}_{m_1, \dots, m_s}$  from [Lan12]. Take an open compact subgroup  $\tilde{K}_{m_1, \dots, m_s} \subseteq \tilde{G}_{m_1, \dots, m_s}(\mathbb{A}^\infty)$ . We choose a smooth projective toroidal compactification  $\tilde{\Sigma}_{m_1, \dots, m_s}(\tilde{K}_{m_1, \dots, m_s})^{\text{tor}}$  of  $\tilde{\Sigma}_{m_1, \dots, m_s}(\tilde{K}_{m_1, \dots, m_s})$  over  $\mathbb{Q}$ , and let  $\tilde{\omega}_{m_1, \dots, m_s}^{\text{tor}}$  be the canonical extension of  $\tilde{\omega}_{m_1, \dots, m_s}$  to  $\tilde{\Sigma}_{m_1, \dots, m_s}(\tilde{K}_{m_1, \dots, m_s})^{\text{tor}}$ . Then by [Lan12, Definition 5.3.4], for every  $w \geq 0$ , we have the algebraic  $q$ -expansion map

$$H^0(\tilde{\Sigma}_{m_1, \dots, m_s}(\tilde{K}_{m_1, \dots, m_s})^{\text{tor}}, (\tilde{\omega}_{m_1, \dots, m_s}^{\text{tor}})^{\otimes w}) \otimes_{\mathbb{Q}} \mathbb{C} \rightarrow \text{SF}_{m_1, \dots, m_s}(\mathbb{C})$$

(Definition 2.5) at the cusp “at infinity”. We remark that the map  $\mathbf{q}_{m_1, \dots, m_s}$  is not necessarily injective, since we only expand the section on the connected component of  $\tilde{\Sigma}_{m_1, \dots, m_s}(\tilde{K}_{m_1, \dots, m_s})^{\text{tor}} \otimes_{\mathbb{Q}} \mathbb{C}$  that contains the cusp “at infinity”. By [Lan12, Remark 5.2.14], the map  $\tilde{h}_{m_1, \dots, m_s}$  (2.8) lifts uniquely to a map

$$\tilde{\mathcal{H}}_{m_1, \dots, m_s}^{[w]}(\tilde{K}_{m_1, \dots, m_s}) \rightarrow H^0(\tilde{\Sigma}_{m_1, \dots, m_s}(\tilde{K}_{m_1, \dots, m_s})^{\text{tor}}, (\tilde{\omega}_{m_1, \dots, m_s}^{\text{tor}})^{\otimes w}),$$

hence we obtain a map

$$(2.9) \quad \mathbf{q}_{m_1, \dots, m_s} : \tilde{\mathcal{H}}_{m_1, \dots, m_s}^{[w]}(\tilde{K}_{m_1, \dots, m_s}) \otimes_{\mathbb{Q}} \mathbb{C} \rightarrow \text{SF}_{m_1, \dots, m_s}(\mathbb{C}),$$

which is independent of the choice of the toroidal compactification. Thus, by passing to the colimit, we obtain a map

$$(2.10) \quad \mathbf{q}_{m_1, \dots, m_s} : \tilde{\mathcal{H}}_{m_1, \dots, m_s}^{[w]} \otimes_{\mathbb{Q}} \mathbb{C} \rightarrow \text{SF}_{m_1, \dots, m_s}(\mathbb{C}),$$

which fits into the following commutative diagram

$$(2.11) \quad \begin{array}{ccc} \tilde{\mathcal{A}}_{m, \text{hol}}^{[w]} & \xrightarrow[\text{(2.7)}]{\tilde{\rho}_{m_1, \dots, m_s}} & \tilde{\mathcal{A}}_{m_1, \dots, m_s, \text{hol}}^{[w]} \\ \tilde{h}_m \downarrow & & \downarrow \tilde{h}_{m_1, \dots, m_s} \\ \tilde{\mathcal{H}}_m^{[w]} \otimes_{\mathbb{Q}} \mathbb{C} & \xrightarrow{\tilde{\sigma}_{m_1, \dots, m_s}^*} & \tilde{\mathcal{H}}_{m_1, \dots, m_s}^{[w]} \otimes_{\mathbb{Q}} \mathbb{C} \\ \mathbf{q}_m \downarrow & & \downarrow \mathbf{q}_{m_1, \dots, m_s} \\ \text{SF}_m(\mathbb{C}) & \xrightarrow[\text{Def. 2.5}]{\mathcal{Q}_{m_1, \dots, m_s}} & \text{SF}_{m_1, \dots, m_s}(\mathbb{C}) \end{array}$$

of complex vector spaces.

**Definition 2.8.** Denote by  $\mathfrak{D}_E \subseteq O_E$  the different ideal of  $E/\mathbb{Q}$ . The (projective)  $O_E$ -lattice  $\mathcal{W}_m := (O_E)^m \oplus (\mathfrak{D}_E^{-1})^m$  of  $W_m$  defines an integral model  $\mathcal{G}_m$  (resp.  $\tilde{\mathcal{G}}_m$ ) of  $G_m$  (resp.  $\tilde{G}_m$ ) over  $O_F$  (resp.  $\mathbb{Z}$ ).<sup>5</sup> Similarly, we have  $\mathcal{G}_{m_1, \dots, m_s}$  and  $\tilde{\mathcal{G}}_{m_1, \dots, m_s}$  and their parabolic subgroups  $\mathcal{P}_{m_1, \dots, m_s}$  and  $\tilde{\mathcal{P}}_{m_1, \dots, m_s}$ , respectively.

For future use, we introduce some standard open compact subgroups. Take two positive integers  $\Delta$  and  $\Delta'$  that are coprime to each other. We put

$$\tilde{K}_{m_1, \dots, m_s}(\Delta, \Delta') := \tilde{\mathcal{G}}_{m_1, \dots, m_s}(\widehat{\mathbb{Z}}) \times_{\tilde{\mathcal{G}}_{m_1, \dots, m_s}(\mathbb{Z}/\Delta\Delta')} \tilde{\mathcal{P}}_{m_1, \dots, m_s}(\mathbb{Z}/\Delta).$$

**Lemma 2.9.** When  $\tilde{K}_{m_1, \dots, m_s} = \tilde{K}_{m_1, \dots, m_s}(\Delta, \Delta')$ , the map (2.9) is equivariant under  $\text{Aut}(\mathbb{C}/\mathbb{Q}(\Delta'))$ , where we recall that  $\mathbb{Q}(\Delta') \subseteq \mathbb{C}$  is the subfield generated by  $\Delta'^l$ -th roots of unity for all  $l \geq 1$ .

*Proof.* This follows from the fact that the cusp “at infinity” is defined over the subfield  $\mathbb{Q}(\Delta')$  at this level structure. See [Lan12] for more details.  $\square$

<sup>5</sup>For  $v \in \mathbb{V}_F^{\text{in}}$ ,  $\mathcal{G}_m(O_{F,v}) = K_{m,v}$  if and only if  $d_v = 0$  and  $v \notin \mathbb{V}_F^{\text{ram}}$ .

*Remark 2.10.* Denote by  $\widetilde{G}_{m_1, \dots, m_s}^{\text{der}}$  the derived subgroup of  $\widetilde{G}_{m_1, \dots, m_s}$  and consider the maximal abelian quotient  $\widetilde{G}_{m_1, \dots, m_s}^{\text{ab}} := \widetilde{G}_{m_1, \dots, m_s} / \widetilde{G}_{m_1, \dots, m_s}^{\text{der}}$ . Since  $\widetilde{G}_{m_1, \dots, m_s}^{\text{der}}$  is simply connected, for every open compact subgroup  $\widetilde{K}_{m_1, \dots, m_s} \subseteq \widetilde{G}_{m_1, \dots, m_s}(\mathbb{A}^\infty)$ , the natural map

$$\Sigma_{m_1, \dots, m_s}(\widetilde{K}_{m_1, \dots, m_s})(\mathbb{C}) \rightarrow \widetilde{G}_{m_1, \dots, m_s}^{\text{ab}}(\mathbb{Q}) \backslash \widetilde{G}_{m_1, \dots, m_s}^{\text{ab}}(\mathbb{A}^\infty) / \widetilde{K}_{m_1, \dots, m_s}^{\text{ab}}$$

has connected fibers, where  $\widetilde{K}_{m_1, \dots, m_s}^{\text{ab}}$  denotes the image of  $\widetilde{K}_{m_1, \dots, m_s}$  in  $\widetilde{G}_{m_1, \dots, m_s}^{\text{ab}}(\mathbb{A}^\infty)$ . It is clear that  $\widetilde{K}_{m_1, \dots, m_s}(\Delta, \Delta')^{\text{ab}}$  depends only on  $\Delta'$ , which we denote by  $\widetilde{K}_{m_1, \dots, m_s}^{\text{ab}}(\Delta')$ .

### 3. CYCLOTOMIC $p$ -ADIC $L$ -FUNCTION

In this section, we construct the  $p$ -adic  $L$ -function. We fix an even positive integer  $n = 2r$ .

**3.1. Doubling space and degenerate principal series.** We have the doubling skew-hermitian space  $W_r^\square := W_r \oplus \bar{W}_r$ . Let  $G_r^\square$  be the unitary group of  $W_r^\square$ , which admits a canonical embedding  $\iota: G_r \times G_r \hookrightarrow G_r^\square$ . We now take a basis  $\{e_1^\square, \dots, e_{4r}^\square\}$  of  $W_r^\square$  by the formula

$$e_i^\square = e_i, \quad e_{r+i}^\square = -\bar{e}_i, \quad e_{2r+i}^\square = e_{r+i}, \quad e_{3r+i}^\square = \bar{e}_{r+i}$$

for  $1 \leq i \leq r$ , under which we may identify  $W_r^\square$  with  $W_{2r}$  and  $G_r^\square$  with  $G_{2r}$ . Put  $w_r^\square := w_{2r}$ ,  $P_r^\square := P_{2r}$  and  $N_r^\square := N_{2r}$ . We denote by

$$\delta_r^\square: P_r^\square \rightarrow \mathbf{G}_F$$

the composition of the Levi quotient map  $P_r^\square = P_{2r} \rightarrow M_{2r}$ , the isomorphism  $m^{-1}: M_{2r} \rightarrow \text{Res}_{E/F} \text{GL}_{2r}$ , the determinant  $\text{Res}_{E/F} \text{GL}_{2r} \rightarrow \text{Res}_{E/F} \mathbf{G}$  and the norm  $\text{Nm}_{E/F}: \text{Res}_{E/F} \mathbf{G} \rightarrow \mathbf{G}_F$ . Put

$$\mathbf{w}_r := \begin{pmatrix} & & & 1_r \\ & & 1_r & \\ -1_r & & 1_r & \\ & & & 1_r \quad 1_r \end{pmatrix} \in G_r^\square(F).$$

Then  $P_r^\square \cdot \mathbf{w}_r \cdot \iota(G_r \times G_r)$  is Zariski open in  $G_r^\square$ .

*Remark 3.1.* The embedding  $\iota: G_r \times G_r \hookrightarrow G_r^\square = G_{2r}$  differs from the one induced by the isometry (2.1) by the involution  $\text{id} \times \dagger$  on  $G_r \times G_r$ .

Let  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times$  be a finite character, regarded as an automorphic character of  $\mathbb{A}_F^\times$ . For every place  $v$  of  $F$ , we have the degenerate principal series of  $G_r^\square(F_v)$ , which is defined as the normalized induced representation

$$\mathbf{I}_{r,v}^\square(\chi_v) := \text{Ind}_{P_r^\square(F_v)}^{G_r^\square(F_v)}(\chi_v \circ \delta_{r,v}^\square)$$

of  $G_r^\square(F_v)$  with complex coefficients. For every  $f \in \mathbf{I}_{r,v}^\square(\chi_v)$  and every  $T^\square \in \text{Herm}_{2r}^\circ(F_v)$ , we can regularize the following integral

$$(3.1) \quad W_{T^\square}(f) := \int_{\text{Herm}_{2r}(F_v)} f(\mathbf{w}_r^\square n(b)) \psi_{F,v}(\text{tr } T^\square b)^{-1} db,$$

where  $db$  is the self-dual measure on  $\text{Herm}_{2r}(F_v)$  with respect to  $\psi_{F,v}$ , which is not necessarily absolutely convergent. It is defined by (the value at  $s = 0$  of) the analytic continuation of a family of integrals  $W_{T^\square}(s, f)$  indexed by  $s \in \mathbb{C}$ , which is absolutely convergent when  $\text{Re } s$  is large enough (see [Wal88, Theorem 8.1] and [Kar79, Corollary 3.6.1]).

In order to show the rationality of our  $p$ -adic  $L$ -function, we need to extend the degenerate principal series to  $\widetilde{G}_{2r}$ . Recall that we have a natural inclusion  $\text{Res}_{F/\mathbb{Q}} G_r^\square = \text{Res}_{F/\mathbb{Q}} G_{2r} \hookrightarrow \widetilde{G}_{2r}$ . We have a map

$$s: \mathbf{G}_{\mathbb{Q}} \rightarrow \widetilde{G}_{2r}$$

sending  $c$  to  $\begin{pmatrix} c^{1_{2r}} & \\ & 1_{2r} \end{pmatrix}$ . Then the natural map  $\text{Res}_{F/\mathbb{Q}} P_{2r} \times s(\mathbf{G}_{\mathbb{Q}}) \rightarrow \widetilde{P}_{2r}$  is an isomorphism.

Take a place  $w$  of  $\mathbb{Q}$ . Put

$$\psi_{F,w} := \prod_{v \in \mathbb{V}_F^{(w)}} \psi_{F,v}, \quad \chi_w := \prod_{v \in \mathbb{V}_F^{(w)}} \chi_v, \quad \mathbf{I}_{r,w}^\square(\chi_w) := \bigotimes_{v \in \mathbb{V}_F^{(w)}} \mathbf{I}_{r,v}^\square(\chi_v),$$



and

$$\delta_{r,w}^\square := \prod_{v \in \mathbf{V}_F^{(w)}} \delta_{r,v}^\square : \prod_{v \in \mathbf{V}_F^{(w)}} P_r^\square(F_v) = (\text{Res}_{F/\mathbb{Q}} P_r^\square)(\mathbb{Q}_w) \rightarrow (F \otimes \mathbb{Z}_w)^\times.$$

The map  $\delta_{r,w}^\square$  extends uniquely to a map  $\widetilde{\delta}_{r,w}^\square$  along the inclusion  $(\text{Res}_{F/\mathbb{Q}} P_r^\square)(\mathbb{Q}_w) = (\text{Res}_{F/\mathbb{Q}} P_{2r})(\mathbb{Q}_w) \subseteq \widetilde{P}_{2r}(\mathbb{Q}_w)$  that sends  $s(c)$  to  $c^{2r}$  for  $c \in \mathbb{Q}_w^\times$ . Then we have a canonical isomorphism

$$\mathbf{I}_{r,w}^\square(\chi_w) \simeq \text{Ind}_{\widetilde{P}_{2r}(\mathbb{Q}_w)}^{\widetilde{G}_{2r}(\mathbb{Q}_w)}(\chi_w \circ \widetilde{\delta}_{r,w}^\square)$$

so that  $\mathbf{I}_{r,w}^\square(\chi_w)$  becomes a representation of  $\widetilde{G}_{2r}(\mathbb{Q}_w)$ . For every  $T^\square \in \text{Herm}_{2r}^\circ(F \otimes \mathbb{Z}_w)$ , we define the functional  $W_{T^\square}(-)$  on  $\mathbf{I}_{r,w}^\square(\chi_w)$  to be the product of the corresponding ones over  $v \in \mathbf{V}_F^{(w)}$ .

**Lemma 3.2.** *For every  $v \in \mathbf{V}_F^{(\infty)}$ , denote by  $f_v^{[r]} \in \mathbf{I}_{r,v}^\square(\chi_v) = \mathbf{I}_{r,v}^\square(\mathbf{1})$  the unique section whose restriction to  $K_{2r,v}$  is the character  $\kappa_{2r,v}^r$ . Put  $f_\infty^{[r]} := \otimes_{v \in \mathbf{V}_F^{(\infty)}} f_v^{[r]}$ . Then there exists  $W_{2r} \in \mathbb{Q}_{>0}$  such that*

$$W_{T^\square}(f_\infty^{[r]}) = W_{2r} \cdot b_{2r}^\infty(\mathbf{1}) \cdot \exp(-2\pi \text{Tr}_{F/\mathbb{Q}} \text{tr } T^\square)$$

for every  $T^\square \in \text{Herm}_{2r}^\circ(F)^+$ .

*Proof.* For two elements  $x, y \in \mathbb{C}^\times$ , we write  $x \sim y$  if their quotient is rational.

By [Liu11a, Proposition 4.5(2)], we have

$$W_{T^\square}(f_\infty^{[r]}) = \left( \frac{(2\pi)^{r(2r+1)}}{\Gamma(1)\Gamma(2)\cdots\Gamma(2r)} \right)^{[F:\mathbb{Q}]} \exp(-2\pi \text{Tr}_{F/\mathbb{Q}} \text{tr } T^\square)$$

for every  $T^\square \in \text{Herm}_{2r}^\circ(F)^+$ . The positivity of  $W_{2r}$  then follows. Thus, it remains to show that  $b_{2r}^\infty(\mathbf{1}) \sim \pi^{r(2r+1)[F:\mathbb{Q}]}$ .

Write  $L(s, \eta_{E/F}^i)$  for the complete  $L$ -function for the self-dual character  $\eta_{E/F}^i$ . Then by the functional equation, we have

$$\prod_{i=1}^{2r} L(i, \eta_{E/F}^i) \sim \prod_{i=1}^{2r} L(1-i, \eta_{E/F}^i).$$

By a well-known result of Siegel,  $\prod_{i=1}^{2r} L^\infty(1-i, \eta_{E/F}^i)$  is rational. It follows that

$$b_{2r}^\infty(\mathbf{1}) \sim \frac{\prod_{i=1}^{2r} L_\infty(1-i, \eta_{E/F}^i)}{\prod_{i=1}^{2r} L_\infty(i, \eta_{E/F}^i)} = \left( \frac{\prod_{i=1}^{2r} L_{\mathbb{R}}(1-i, \text{sgn}^i)}{\prod_{i=1}^{2r} L_{\mathbb{R}}(i, \text{sgn}^i)} \right)^{[F:\mathbb{Q}]} \sim \left( \frac{\pi^{r^2}}{\pi^{-r(r+1)}} \right)^{[F:\mathbb{Q}]} = \pi^{r(2r+1)[F:\mathbb{Q}]}.$$

The lemma follows.  $\square$

From now to the end of this subsection, we assume  $w \neq \infty$ .

**Lemma 3.3.** *We have*

(1) *For  $v \in \mathbf{V}_F^{(w)}$  and  $b \in \text{Herm}_{2r}(F_v)$ , the relation*

$$W_{T^\square}(n(b)f) = \psi_{F,v}(\text{tr } T^\square b) \cdot W_{T^\square}(f)$$

*holds for every  $f \in \mathbf{I}_{r,v}^\square(\chi_v)$  and every  $T^\square \in \text{Herm}_{2r}^\circ(F_v)$ .*

(2) *For  $v \in \mathbf{V}_F^{(w)}$  and  $a \in \text{GL}_{2r}(E_v)$ , the relation*

$$W_{T^\square}(m(a)f) = \chi_v(\text{Nm}_{E/F} \det a)^{-1} |\det a|_E^r \cdot W_{T^\square}(f)$$

*holds for every  $f \in \mathbf{I}_{r,v}^\square(\chi_v)$  and every  $T^\square \in \text{Herm}_{2r}^\circ(F_v)$ .*

(3) *For  $c \in \mathbb{Q}_w^\times$ , the relation*

$$W_{T^\square}(s(c)f) = \chi_w(c)^{-2r} |c|_F^{2r^2} \cdot W_{T^\square}(f)$$

*holds for every  $f \in \mathbf{I}_{r,w}^\square(\chi_w)$  and every  $T^\square \in \text{Herm}_{2r}^\circ(F \otimes \mathbb{Z}_w)$ .*

*Proof.* This is well-known. For readers' convenience, we give a (formal) proof.

For (1), we have

$$\begin{aligned} W_{T^\square}(n(b)f) &= \int_{\text{Herm}_{2r}(F_v)} f(\mathbf{w}_r^\square n(b')n(b))\psi_{F,v}(\text{tr } T^\square b')^{-1} db' \\ &= \psi_{F,v}(\text{tr } T^\square b) \int_{\text{Herm}_{2r}(F_v)} f(\mathbf{w}_r^\square n(b'+b))\psi_{F,v}(\text{tr } T^\square (b'+b))^{-1} db' = \psi_{F,v}(\text{tr } T^\square b) \cdot W_{T^\square}(f). \end{aligned}$$

For (2), we have

$$\begin{aligned} W_{T^\square}(m(a)f) &= \int_{\text{Herm}_{2r}(F_v)} f(\mathbf{w}_r^\square n(b)m(a))\psi_{F,v}(\text{tr } T^\square b)^{-1} db \\ &= \int_{\text{Herm}_{2r}(F_v)} f(\mathbf{w}_r^\square m(a)n(a^{-1}b^t a^{c,-1}))\psi_{F,v}(\text{tr } T^\square b)^{-1} db \\ &= \int_{\text{Herm}_{2r}(F_v)} f(m({}^t a^{c,-1})\mathbf{w}_r^\square n(a^{-1}b^t a^{c,-1}))\psi_{F,v}(\text{tr}({}^t a^c T^\square a)(a^{-1}b^t a^{c,-1}))^{-1} db \\ &= \chi_v(\text{Nm}_{E/F} \det a)^{-1} |\det a|_E^r \int_{\text{Herm}_{2r}(F_v)} f(\mathbf{w}_r^\square n(b))\psi_{F,v}(\text{tr}({}^t a^c T^\square a)b)^{-1} db \\ &= \chi_v(\text{Nm}_{E/F} \det a)^{-1} |\det a|_E^r \cdot W_{{}^t a^c T^\square a}(f). \end{aligned}$$

The proof for (3) is similar to (2) and we omit it. The lemma is proved.  $\square$

**Notation 3.4.** Let  $v \in \mathbf{V}_F^{\text{fin}}$  be a finite place.

- (1) We denote by  $\mathbf{I}_{r,v}^\square(\chi_v)^\circ$  the subspace of  $\mathbf{I}_{r,v}^\square(\chi_v)$  consisting of sections that are supported on the big Bruhat cell  $P_r^\square(F_v) \cdot \mathbf{w}_r^\square \cdot N_r^\square(F_v)$ .
- (2) When  $v \in \mathbf{V}_F^{\text{fin}} \setminus \mathbf{V}_F^{(p)}$ , we denote by  $f_{\chi_v}^{\text{sph}} \in \mathbf{I}_{r,v}^\square(\chi_v)$  the unique section that takes value 1 on  $K_{2r,v}$ .

It is clear that  $\mathbf{I}_{r,v}^\square(\chi_v)^\circ$  is stable under the action of  $P_r^\square(F_v)$ . For  $f \in \mathbf{I}_{r,v}^\square(\chi_v)^\circ$  and  $T^\square \in \text{Herm}_{2r}(F_v)$ , we put

$$W_{T^\square}(f) := \int_{\text{Herm}_{2r}(F_v)} f(\mathbf{w}_r^\square n(b))\psi_{F,v}(\text{tr } T^\square b)^{-1} db,$$

which is in fact a finite sum and coincides with (3.1) for  $T^\square \in \text{Herm}_{2r}^\circ(F_v)$ . It is clear that the assignment  $T^\square \mapsto W_{T^\square}(f)$  is a Schwartz function on  $\text{Herm}_{2r}(F_v)$ . Conversely, using the Fourier inversion formula, we know that for every  $\mathbf{f} \in \mathcal{S}(\text{Herm}_{2r}(F_v))$ , there exists a unique section  $\mathbf{f}^{\chi_v} \in \mathbf{I}_{r,v}^\square(\chi_v)^\circ$  such that  $W_{T^\square}(\mathbf{f}^{\chi_v}) = \mathbf{f}(T^\square)$  holds for every  $T^\square \in \text{Herm}_{2r}(F_v)$ . In other words, we obtain a bijection

$$(3.2) \quad -\chi_v : \mathcal{S}(\text{Herm}_{2r}(F_v)) \xrightarrow{\sim} \mathbf{I}_{r,v}^\square(\chi_v)^\circ.$$

Put  $\mathbf{I}_{r,w}^\square(\chi_w)^\circ := \bigotimes_{v \in \mathbf{V}_F^{(w)}} \mathbf{I}_{r,v}^\square(\chi_v)^\circ$  and we obtain an isomorphism

$$-\chi_w : \mathcal{S}(\text{Herm}_{2r}(F \otimes \mathbb{Z}_w)) \xrightarrow{\sim} \mathbf{I}_{r,w}^\square(\chi_w)^\circ$$

by taking product over  $v \in \mathbf{V}_F^{(w)}$

**Lemma 3.5.** *Suppose that  $w \neq p$ .*

- (1) *For every  $v \in \mathbf{V}_F^{(w)} \setminus \mathbf{V}_F^{\text{ram}}$ , every  $T^\square \in \text{Herm}_{2r}^\circ(F_v)$  and every  $g \in G_{2r}(F_v)$ , there exists a unique element  ${}^s\mathbf{W}_{T^\square,v}^{\text{sph}} \in \mathbb{Z}_{(p)}\langle w \rangle[X, X^{-1}]$  such that*

$${}^s\mathbf{W}_{T^\square,v}^{\text{sph}}(\chi_v(\varpi_v)) = b_{2r,v}(\chi) \cdot W_{T^\square}(g \cdot f_{\chi_v}^{\text{sph}})$$

*holds for every finite character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times$ , where  $\varpi_v$  is an arbitrary uniformizer of  $F_v$ . Moreover,*

$$\mathbf{W}_{T^\square,v}^{\text{sph}} := {}_{14r}\mathbf{W}_{T^\square,v}^{\text{sph}} \in \mathbb{Z}[X].$$

- (2) *For every  $f \in \mathbf{I}_{r,w}^\square(\chi_w)$  and every  $T^\square \in \text{Herm}_{2r}^\circ(F \otimes \mathbb{Z}_w)$ , we have*

$$W_{T^\square}(\sigma f) = \sigma W_{T^\square}(f)$$

*for  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}\langle w \rangle)$ .*

(3) For every  $f \in \mathbb{I}_{r,w}^\square(\chi_w)^\circ$  that is fixed by  $\widetilde{\mathcal{P}}_{2r}(\mathbb{Z}_w)$  and every  $T^\square \in \text{Herm}_{2r}^\circ(F \otimes \mathbb{Z}_w)$ , we have

$$W_{T^\square}(\sigma f) = \sigma W_{T^\square}(f)$$

for  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q})$ .

*Proof.* For (1), by Lemma 3.3(1,2) and the Iwasawa decomposition  $G_{2r}(F_v) = P_{2r}(F_v)K_{2r,v}$ , it suffices to consider the case where  $g = 1_{4r}$ . Then the statement follows from [LZ, Theorem 3.5.1], together with the discussion in [LZ, Sections 3.2 & 3.3].<sup>6</sup>

Part (2) follows from the proof of [Kar79, Corollary 3.6.1] and the fact that  $\psi_{F,w}$  takes values in  $\mathbb{Q}\langle w \rangle$ .

For (3), put  $\mathfrak{D}_b := \{c^t a^c b a \mid a \in \text{GL}_{2r}(O_E \otimes \mathbb{Z}_w), c \in \mathbb{Z}_w^\times\}$  for every  $b \in \text{Herm}_{2r}(F \otimes \mathbb{Z}_w)$ , which is an open compact subset of  $\text{Herm}_{2r}(F \otimes \mathbb{Z}_w)$ . It follows easily that

$$\int_{\mathfrak{D}_b} \psi_{F,v}(\text{tr } T^\square b')^{-1} db' \in \mathbb{Q}.$$

Since  $\chi_w$  is unramified, the assignment  $b' \mapsto f(w_r^\square n(b'))$  is constant on each subset  $\mathfrak{D}_b$ . Thus, (3) follows.  $\square$

**Lemma 3.6.** *The representation  $\mathbb{I}_{r,w}^\square(\chi_w)$  is semisimple as a representation of  $\widetilde{G}_{2r}(\mathbb{Q}_w)$ . When  $w \neq p$ , every irreducible summand of  $\mathbb{I}_{r,w}^\square(\chi_w)$  contains a nonzero element  $f$  in  $\mathbb{I}_{r,w}^\square(\chi_w)^\circ$  that is fixed by  $\widetilde{\mathcal{P}}_{2r}(\mathbb{Z}_w)$ .*

*Proof.* For the first statement, it suffices to show that  $\mathbb{I}_{r,w}^\square(\chi_w)$  is semisimple as a representation of  $(\text{Res}_{F/\mathbb{Q}} G_{2r})(\mathbb{Q}_w)$ , which is due to [KS97].

Now we show the second statement. For every  $v \in \mathbb{V}_F^{(w)}$ , by [KS97, Theorem 1.2 & Theorem 1.3],  $\mathbb{I}_{r,v}^\square(\chi_v)$  is an irreducible representation of  $G_{2r}(F_v)$  unless  $\chi_v^2 = \mathbf{1}$ . Moreover, when  $\chi_v^2 = \mathbf{1}$ , each direct summand of  $\mathbb{I}_{r,v}^\square(\chi_v)$  is of the form  $\mathbb{I}(V_v)$  for some (nondegenerate) hermitian space  $V_v$  over  $E_v$  of rank  $2r$ . Here,  $\mathbb{I}(V_v)$  is the image of the Siegel–Weil section map  $\mathcal{S}(V_v^{2r}) \rightarrow \mathbb{I}_{r,v}^\square(\chi_v)$  under the Weil representation with respect to (the standard additive character  $\psi_{F,v}$  and) the splitting character  $\chi_v \circ \text{Nm}_{E/F}$  (again see [KS97]). Put  $\mathbb{V} := \{v \in \mathbb{V}_F^{(w)} \mid \chi_v^2 = \mathbf{1}\}$ .

Now let  $\mathbb{I}$  be an irreducible summand of  $\mathbb{I}_{r,w}^\square(\chi_w)$  as a representation of  $\widetilde{G}_{2r}(\mathbb{Q}_w)$ . One can find a collection of hermitian spaces  $V_v$  over  $E_v$  of rank  $2r$  for  $v \in \mathbb{V}$  such that  $\mathbb{I}$  contains

$$\left( \bigotimes_{v \in \mathbb{V}} \mathbb{I}(V_v) \right) \otimes \left( \bigotimes_{v \in \mathbb{V}_F^{(w)} \setminus \mathbb{V}} \mathbb{I}_{r,v}^\square(\chi_v) \right).$$

For every  $v \in \mathbb{V}_F^{(w)}$ , we define a subset  $\mathfrak{I}_v$  of  $\text{Herm}_{2r}^\circ(F_v)$  as follows. If  $v \in \mathbb{V}$ , then we define  $\mathfrak{I}_v$  to be the intersection of  $\text{Herm}_{2r}^\circ(F_v)$  and the image of the moment map  $V_v^{2r} \rightarrow \text{Herm}_{2r}(F_v)$  (see §4.1(H1) if one needs recall). If  $v \notin \mathbb{V}$ , then we define  $\mathfrak{I}_v$  to be  $\text{Herm}_{2r}^\circ(F_v)$ . Take any open compact subset  $\mathfrak{I}$  of  $\text{Herm}_{2r}(F \otimes \mathbb{Z}_w) = \prod_{v \in \mathbb{V}_F^{(w)}} \text{Herm}_{2r}(F_v)$  that is contained in  $\prod_{v \in \mathbb{V}_F^{(w)}} \mathfrak{I}_v \cap \text{Herm}_{2r}(O_{F_v})$  satisfying that  $c^t a^c \mathfrak{I} a = \mathfrak{I}$  for every  $a \in \text{GL}_{2r}(O_E \otimes \mathbb{Z}_w)$  and every  $c \in \mathbb{Z}_w^\times$ . Then  $(\mathbf{1}_{\mathfrak{I}})^{\chi_w} \in \mathbb{I}_{r,w}^\square(\chi_w)^\circ$  is a nonzero element of  $\mathbb{I}$ . Moreover, by Lemma 3.3, it is fixed by  $\widetilde{\mathcal{P}}_{2r}(\mathbb{Z}_w)$ .

The lemma is proved.  $\square$

In the rest of this subsection, we construct some explicit sections in  $\mathbb{I}_{r,p}^\square(\chi_p)^\circ$ .

**Notation 3.7.** For every place  $v \in \mathbb{V}_F^{(p)}$ , we

- fix a uniformizer  $\varpi_v$  of  $F_v$ ,
- for every element  $e = (e_u)_u \in \mathbb{Z}^{\mathbb{P}_v}$ , put  $|e| := \sum_{u \in \mathbb{P}_v} e_u$  and denote by  $\varpi_v^e$  the element in  $E_v = \prod_{u \in \mathbb{P}_v} E_u$  whose component in  $E_u$  is  $\varpi_v^{e_u}$ ,
- for  $u \in \mathbb{P}_v$ , denote by  $1_u \in \mathbb{Z}^{\mathbb{P}_v}$  the element that takes values 1 at  $u$  and 0 at  $u^c$ ,
- for every  $u \in \mathbb{P}_v$ , introduce an element

$$U_u := \sum_{b \in \text{Herm}_r(O_{F_v}/\varpi_v)} \left[ \begin{pmatrix} 1_r & \varpi_v^{-d_v} b^\sharp \\ & 1_r \end{pmatrix} \begin{pmatrix} \varpi_v^{1_u} \cdot 1_r & \\ & \varpi_v^{-1_{u^c}} \cdot 1_r \end{pmatrix} \right] \in \mathbb{Z}[G_r(F_v)],$$

where  $b^\sharp \in \text{Herm}_r(O_{F_v})$  denotes the Teichmüller lift of  $b$ ,

<sup>6</sup>Though [LZ] only treats the case where  $v$  is inert in  $E$ , the same argument works in the case where  $v$  splits in  $E$  as well.

- for every  $e = (e_u)_u \in \mathbb{N}^{\mathbb{P}_v}$ , define

$$U_v^e := \prod_{u \in \mathbb{P}_v} U_u^{e_u} \in \mathbb{Z}[G_r(F_v)],$$

where we note that the subalgebra of  $\mathbb{Z}[G_r(F_v)]$  generated by  $U_u$  for  $u \in \mathbb{P}_v$  is commutative.

**Construction 3.8.** For  $v \in \mathbb{V}_F^{(p)}$  and every element  $e \in \mathbb{Z}^{\mathbb{P}_v}$ , let  $\mathfrak{T}_v^{[e]}$  be the subset of  $\text{Herm}_{2r}(F_v)$  consisting of elements

$$T^\square = \begin{pmatrix} T_{11}^\square & T_{12}^\square \\ T_{21}^\square & T_{22}^\square \end{pmatrix}$$

satisfying  $T_{11}^\square, T_{22}^\square \in \text{Herm}_r(O_{F_v})$  and  $T_{12}^\square \in \varpi_v^{-e} \cdot \text{GL}_r(O_{E_v})$ . Define a function  $\mathbf{f}_{\chi_v}^{[e]} \in \mathcal{S}(\text{Herm}_{2r}(F_v))$  by the formula

$$\mathbf{f}_{\chi_v}^{[e]}(T^\square) := \chi_v(\text{Nm}_{E/F} \det T_{12}^\square) \cdot \mathbf{1}_{\mathfrak{T}_v^{[e]}}(T^\square).$$

In particular, we obtain a section  $(\mathbf{f}_{\chi_v}^{[e]})_{\chi_v} \in \mathbb{I}_{r,v}^\square(\chi_v)^\circ$  by (3.2).

In what follows, we will identify  $\mathbb{Z}^{\mathbb{P}}$  and  $\mathbb{N}^{\mathbb{P}}$  with  $\prod_{v \in \mathbb{V}_F^{(p)}} \mathbb{Z}^{\mathbb{P}_v}$  and  $\prod_{v \in \mathbb{V}_F^{(p)}} \mathbb{N}^{\mathbb{P}_v}$ , respectively. For  $e \in \mathbb{Z}^{\mathbb{P}}$ , we put

$$\|e\| := \max_{v \in \mathbb{V}_F^{(p)}} |e_v|, \quad \mathfrak{T}_p^{[e]} := \prod_{v \in \mathbb{V}_F^{(p)}} \mathfrak{T}_v^{[e_v]}, \quad \mathbf{f}_{\chi_p}^{[e]} := \bigotimes_{v \in \mathbb{V}_F^{(p)}} \mathbf{f}_{\chi_v}^{[e_v]}.$$

For  $e \in \mathbb{N}^{\mathbb{P}}$ , we put

$$U_p^e := \bigotimes_{v \in \mathbb{V}_F^{(p)}} U_v^{e_v} \in \bigotimes_{v \in \mathbb{V}_F^{(p)}} \mathbb{Z}[G_r(F_v)] = \mathbb{Z}[G_r(F \otimes \mathbb{Z}_p)].$$

**Lemma 3.9.** For every element  $e \in \mathbb{Z}^{\mathbb{P}}$ , the section  $(\mathbf{f}_{\chi_p}^{[e]})_{\chi_p} \in \mathbb{I}_{r,p}^\square(\chi_p)^\circ$  is invariant under the subgroup  $\widetilde{\mathcal{P}}_{r,p}(\mathbb{Z}_p)$  of  $\widetilde{G}_{2r}(\mathbb{Q}_p)$ .

*Proof.* This follows immediately from Lemma 3.3. □

**Lemma 3.10.** For every element  $e \in \mathbb{Z}^{\mathbb{P}}$  and every  $e_1, e_2 \in \mathbb{N}^{\mathbb{P}}$ , we have

$$(U_p^{e_1} \times U_p^{e_2})(\mathbf{f}_{\chi_p}^{[e]})_{\chi_p} = (\mathbf{f}_{\chi_p}^{[e+e_1^c+e_2]})_{\chi_p},$$

where  $e_1^c := e_1 \circ c$ .

*Proof.* By induction, we may assume either  $e_1 = 0$  or  $e_2 = 0$ . We consider the case where  $e_2 = 0$  and leave the other similar case to the reader. Again by induction, we may assume  $e_1 = 1_u$  for some  $u \in \mathbb{P}$ , with  $v \in \mathbb{V}_F^{(p)}$  its underlying place.

For two square matrices  $a$  and  $b$ , we write  $[a, b]$  for the block diagonal matrix. As an element in  $\mathbb{Z}[\widetilde{G}_{2r}(\mathbb{Q}_p)]$ , we have

$$U_p^{e_1} \times U_p^{e_2} = \sum_{b \in \text{Herm}_r(O_{F_v}/\varpi_v)} \left[ n([\varpi_v^{-d_v} \cdot b^\sharp, 1_r]) \cdot m([\varpi_v^{1_u} \cdot 1_r, 1_r]) \right]$$

in which all components away from  $v$  are  $1_{4r}$ . By Lemma 3.3, we have

$$\begin{aligned} & W_{T^\square}((U_p^{e_1} \times U_p^{e_2})(\mathbf{f}_{\chi_p}^{[e]})_{\chi_p}) \\ &= \sum_{b \in \text{Herm}_r(O_{F_v}/\varpi_v)} W_{T^\square}(n([\varpi_v^{-d_v} \cdot b^\sharp, 1_r]) \cdot m([\varpi_v^{1_u} \cdot 1_r, 1_r]) \cdot (\mathbf{f}_{\chi_p}^{[e]})_{\chi_p}) \\ &= \left( \sum_{b \in \text{Herm}_r(O_{F_v}/\varpi_v)} \psi_{F,v}(\varpi_v^{-d_v} \text{tr } T_{11,v}^\square b^\sharp) \right) W_{T^\square}(m([\varpi_v^{1_u} \cdot 1_r, 1_r]) \cdot (\mathbf{f}_{\chi_p}^{[e]})_{\chi_p}) \\ &= \left( \sum_{b \in \text{Herm}_r(O_{F_v}/\varpi_v)} \psi_{F,v}(\varpi_v^{-d_v} \text{tr } T_{11,v}^\square b^\sharp) \right) \chi_v(\varpi_v^r)^{-1} q_v^{-r^2} \cdot \mathbf{f}_{\chi_p}^{[e]}([\varpi_v^{1_{u^c}} \cdot 1_r, 1_r] T^\square [\varpi_v^{1_u} \cdot 1_r, 1_r]) \\ (3.3) \quad &= \left( \sum_{b \in \text{Herm}_r(O_{F_v}/\varpi_v)} \psi_{F,v}(\varpi_v^{-d_v} \text{tr } T_{11,v}^\square b^\sharp) \right) \chi_v(\varpi_v^r)^{-1} q_v^{-r^2} \cdot \mathbf{f}_{\chi_p}^{[e]} \left( \begin{pmatrix} \varpi_v \cdot T_{11}^\square & \varpi_v^{1_{u^c}} \cdot T_{12}^\square \\ \varpi_v^{1_u} \cdot T_{21}^\square & T_{22}^\square \end{pmatrix} \right). \end{aligned}$$

Since

$$\sum_{b \in \text{Herm}_r(O_{F_v}/\varpi_v)} \psi_{F,v}(\varpi_v^{-d_v} \text{tr } T_{11,v}^\square b^\sharp) = \begin{cases} q_v^{r^2} & \text{if } T_{11,v}^\square \in \text{Herm}_{2r}(O_{F_v}), \\ 0 & \text{if } T_{11,v}^\square \in \varpi_v^{-1} \text{Herm}_{2r}(O_{F_v}) \setminus \text{Herm}_{2r}(O_{F_v}), \end{cases}$$

we have

$$(3.3) = \chi_v(\varpi_v^r)^{-1} \chi_p(\text{Nm}_{E/F} \det \varpi_v^{1,rc} T_{12}^\square) \cdot \mathbf{1}_{\mathfrak{F}_p^{[e+e_1]}(T^\square)} = \chi_p(\text{Nm}_{E/F} \det T_{12}^\square) \cdot \mathbf{1}_{\mathfrak{F}_p^{[e+e_1]}(T^\square)} = \mathbf{f}_{\chi_p}^{[e+e_1]}(T^\square).$$

The lemma follows.  $\square$

**3.2. Siegel hermitian Eisenstein series.** Let  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times$  be a finite character, regarded as an automorphic character of  $\mathbb{A}_F^\times$ . We define  $I_r^\square(\chi)$  to be the restricted tensor product of  $I_{r,v}^\square(\chi_v)$  over all places  $v$  of  $F$ , which is a smooth representation of  $\widetilde{G}_{2r}(\mathbb{A})$ . For  $f_\chi \in I_r^\square(\chi)$ , we have the Siegel hermitian Eisenstein series<sup>7</sup>

$$\begin{aligned} E(g, f_\chi) &:= \sum_{\gamma \in P_{2r}(F) \backslash G_{2r}(F)} f_\chi(\gamma g), \quad g \in G_{2r}(\mathbb{A}_F), \\ \widetilde{E}(g, f_\chi) &:= \sum_{\gamma \in \widetilde{P}_{2r}(\mathbb{Q}) \backslash \widetilde{G}_{2r}(\mathbb{Q})} f_\chi(\gamma g), \quad g \in \widetilde{G}_{2r}(\mathbb{A}). \end{aligned}$$

For a finite set  $\diamond$  of places of  $\mathbb{Q}$  containing  $\{\infty, p\}$ , an element  $e \in \mathbb{Z}^p$  and a section  $f \in I_r^\square(\chi)^{\otimes p}$ , we put

$$(3.4) \quad \widetilde{E}_{[e]}^\diamond(-, \chi, f) := b_{2r}^\diamond(\mathbf{1})^{-1} \cdot b_{2r}^\diamond(\chi) \cdot \widetilde{E}(-, f_\infty^{[r]} \otimes (\mathbf{f}_{\chi_p}^{[e]})^{\chi_p} \otimes f),$$

where  $b_{2r}^\diamond$  is defined in §2.1(F3);  $f_\infty^{[r]}$  is introduced in Lemma 3.2; and  $(\mathbf{f}_{\chi_p}^{[e]})^{\chi_p}$  is introduced in Construction 3.8. It is clear that  $\widetilde{E}_{[e]}^\diamond(-, \chi, f)$  belongs to  $\widetilde{\mathcal{A}}_{2r, \text{hol}}^{[r]}$ . Put

$$(3.5) \quad W_{2r}^\diamond := W_{2r} \cdot b_{2r, \diamond \setminus \{\infty\}}(\mathbf{1}) \in \mathbb{Q}^\times,$$

where  $W_{2r}$  is the constant in Lemma 3.2.

**Lemma 3.11.** *Suppose that  $\|e\| > 0$ . Then for  $f = \otimes_{w \nmid \infty p} f_w$  that is a pure tensor,*

$$\mathbf{q}_{2r} \widetilde{\mathbf{h}}_{2r} \left( \widetilde{E}_{[e]}^\diamond(-, \chi, f) \right) = W_{2r}^\diamond \sum_{T^\square \in \text{Herm}_{2r}^\circ(F)^+} \left( \chi_p(\text{Nm}_{E/F} \det T_{12}^\square) \mathbf{1}_{\mathfrak{F}_p^{[e]}(T^\square)} \cdot \prod_{w \nmid \infty p} W_{T^\square}^\diamond(f_w) \right) q^T$$

in which the product is finite. Here,  $\widetilde{\mathbf{h}}_{2r}$  is the map (2.8);  $\mathbf{q}_{2r}$  is the map (2.10); and

$$W_{T^\square}^\diamond(f_w) := \begin{cases} W_{T^\square}(f_w) & \text{if } w \in \diamond, \\ b_{2r,w}(\chi) \cdot W_{T^\square}(f_w) & \text{if } w \notin \diamond. \end{cases}$$

*Proof.* First, note that when  $\|e\| > 0$ , we have  $\mathbf{f}_{\chi_p}^{[e]}(T^\square) = 0$  for  $T^\square \in \text{Herm}_{2r}(F) \setminus \text{Herm}_{2r}^\circ(F)$ . By the discussion in [Liu11b, Section 2B] and Lemma 3.2, the analytic  $q$ -expansion (2.5) of  $\widetilde{E}(-, f_\infty^{[r]} \otimes (\mathbf{f}_{\chi_p}^{[e]})^{\chi_p} \otimes f)$  equals

$$W_{2r} \cdot b_{2r}^\infty(\mathbf{1}) \sum_{T^\square \in \text{Herm}_{2r}^\circ(F)^+} \left( \chi_p(\text{Nm}_{E/F} \det T_{12}^\square) \mathbf{1}_{\mathfrak{F}_p^{[e]}(T^\square)} \cdot \prod_{w \nmid \infty p} W_{T^\square}(f_w) \right) q^T.$$

It follows that the analytic  $q$ -expansion of  $\widetilde{E}_{[e]}^\diamond(-, \chi, f)$  equals

$$W_{2r}^\diamond \sum_{T^\square \in \text{Herm}_{2r}^\circ(F)^+} \left( \chi_p(\text{Nm}_{E/F} \det T_{12}^\square) \mathbf{1}_{\mathfrak{F}_p^{[e]}(T^\square)} \cdot \prod_{w \nmid \infty p} W_{T^\square}^\diamond(f_w) \right) q^T$$

in which the product is actually finite by [Tan99, Proposition 3.2]. The lemma follows by the coincidence of the analytic and the algebraic  $q$ -expansions [Lan12, Theorem 5.3.5].  $\square$

<sup>7</sup>We remind the reader that the sums in the following expressions are not absolutely convergent in general; they are rather defined by analytic continuation.

Put

$$(3.6) \quad \widetilde{D}_{[e]}^\diamond(-, \chi, f) := \widetilde{\rho}_{r,r} \left( \widetilde{E}_{[e]}^\diamond(-, \chi, f) \right) \in \widetilde{\mathcal{A}}_{r,r,\text{hol}}^{[r]}$$

(see (2.7) for the map  $\widetilde{\rho}_{r,r}$ ).<sup>8</sup> The following proposition concerns the rationality of  $\widetilde{D}_{[e]}^\diamond(-, \chi, f)$ , which is the main result of this subsection.

**Proposition 3.12.** *Suppose that  $\|e\| > 0$  and let  $f \in \mathbb{I}_r^\square(\chi)^{\infty p}$  be a section. For every  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q})$ , we have*

$$\widetilde{h}_{r,r} \left( \widetilde{D}_{[e]}^\diamond(-, \sigma\chi, \sigma f) \right) = \sigma \widetilde{h}_{r,r} \left( \widetilde{D}_{[e]}^\diamond(-, \chi, f) \right),$$

where  $\widetilde{h}_{r,r}$  is the map (2.8).

Note that for  $f \in \mathbb{I}_r^\square(\chi)^{\infty p}$ ,  $\sigma f \in \mathbb{I}_r^\square(\sigma\chi)^{\infty p}$ . Thus, the statement of the proposition makes sense.

*Proof.* Take an integer  $d \geq 1$  such that  $(\mathbf{f}_{\chi_p}^{[e]})^{\chi_p}$  is fixed by the kernel of the map  $\widetilde{\mathcal{G}}_{2r}(\mathbb{Z}_p) \rightarrow \widetilde{\mathcal{G}}_{2r}(\mathbb{Z}/p^d)$ .

We first show that for every  $f \in \mathbb{I}_r^\square(\chi)^{\infty p}$  and every  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}\langle p \rangle)$ , we have

$$(3.7) \quad \widetilde{h}_{2r} \left( \widetilde{E}_{[e]}^\diamond(-, \sigma\chi, \sigma f) \right) = \sigma \widetilde{h}_{2r} \left( \widetilde{E}_{[e]}^\diamond(-, \chi, f) \right).$$

Take an irreducible summand  $\mathbb{I}$  of  $\mathbb{I}_r^\square(\chi)^{\infty p}$  (as a representation of  $\widetilde{\mathcal{G}}_{2r}(\mathbb{A}^{\infty p})$ ). Choose a positive integer  $\Delta = \Delta_1 > 1$  that is coprime to  $p$  such that

- (1) for every rational prime  $w$  not dividing  $p\Delta$ ,  $\mathbb{I}_w$  has nonzero invariants under  $\widetilde{\mathcal{G}}_{2r}(\mathbb{Z}_w)$ ;
- (2) one can write  $\Delta = \Delta_1 \cdot \Delta_2$  with  $(\Delta_1, \Delta_2) = 1$  such that for  $i = 1, 2$ ,  $\prod_{w|\Delta_i} \widetilde{\mathcal{P}}_{2r}(\mathbb{Q}_w)$  maps surjectively to  $\widetilde{\mathcal{G}}_{2r}^{\text{ab}}(\mathbb{Q}) \backslash \widetilde{\mathcal{G}}_{2r}^{\text{ab}}(\mathbb{A}^\infty) / \widetilde{\mathcal{K}}_{2r}^{\text{ab}}(p^d)$  (Remark 2.10).

For every  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}\langle p \rangle)$ , since the map

$$f \mapsto \widetilde{h}_{2r} \left( \widetilde{E}_{[e]}^\diamond(-, \sigma\chi, \sigma f) \right) - \sigma \widetilde{h}_{2r} \left( \widetilde{E}_{[e]}^\diamond(-, \chi, f) \right)$$

is  $\widetilde{\mathcal{G}}_{2r}(\mathbb{A}^{\infty p})$ -equivariant, it suffices to show that there exists a nonzero element  $f = f_\sigma \in \mathbb{I}$  such that

$$(3.8) \quad \widetilde{h}_{2r} \left( \widetilde{E}_{[e]}^\diamond(-, \sigma\chi, \sigma f) \right) - \sigma \widetilde{h}_{2r} \left( \widetilde{E}_{[e]}^\diamond(-, \chi, f) \right) = 0.$$

We choose a nonzero element  $f = \otimes_{w \nmid \infty p} f_w \in \mathbb{I}$  such that  $f_w$  satisfies the condition in Lemma 3.6 for  $w \mid \Delta$  and that  $f_w$  is the unique section that is fixed by  $\widetilde{\mathcal{G}}_{2r}(\mathbb{Z}_w)$  and satisfies  $f_w(1_{4r}) = 1$  for  $w \nmid \Delta$ . Replacing  $\Delta$  by a power of  $\Delta$ , we may assume that  $f_w$  is invariant under  $\widetilde{\mathcal{G}}_{2r}(\mathbb{Z}_w) \times_{\widetilde{\mathcal{G}}_{2r}(\mathbb{Z}_w/\Delta)} \widetilde{\mathcal{P}}_{2r}(\mathbb{Z}_w/\Delta)$  for every  $w \mid \infty p$ . In particular, we have

$$\widetilde{E}_{[e]}^\diamond(-, \chi, f) \in \widetilde{\mathcal{A}}_{2r,\text{hol}}^{[r]}(\widetilde{\mathcal{K}}_{2r}(\Delta, p^d)).$$

Now we claim that (3.8) holds for  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}\langle p\Delta_1 \rangle)$ . By property (2) for  $\Delta$ , we may choose finitely many elements  $g_1, \dots, g_s \in \prod_{w|\Delta_1} \widetilde{\mathcal{P}}_{2r}(\mathbb{Q}_w)$  that map surjectively to  $\widetilde{\mathcal{G}}_{2r}^{\text{ab}}(\mathbb{Q}) \backslash \widetilde{\mathcal{G}}_{2r}^{\text{ab}}(\mathbb{A}^\infty) / \widetilde{\mathcal{K}}_{2r}^{\text{ab}}(p^d)$ . For  $1 \leq j \leq s$ , there exists an integer  $d_j \geq 1$  such that  $\widetilde{E}_{[e]}^\diamond(-, \chi, g_j \cdot f)$  belongs to  $\widetilde{\mathcal{A}}_{2r,\text{hol}}^{[r]}(\widetilde{\mathcal{K}}_{2r}(\Delta_2, p^d \Delta_1^{d_j}))$ . By Lemma 2.9 and Remark 2.10, it suffices to check that

$$\mathbf{q}_{2r} \widetilde{h}_{2r} \left( \widetilde{E}_{[e]}^\diamond(-, \sigma\chi, g_j \cdot \sigma f) \right) - \sigma \mathbf{q}_{2r} \widetilde{h}_{2r} \left( \widetilde{E}_{[e]}^\diamond(-, \chi, g_j \cdot f) \right) = 0$$

for every  $1 \leq j \leq s$  and every  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}\langle p\Delta_1 \rangle)$ . But this follows from Lemma 3.11 and Lemma 3.5. Since the roles of  $\Delta_1$  and  $\Delta_2$  are symmetric, (3.8) also holds for  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}\langle p\Delta_2 \rangle)$ . Together, (3.8) holds for  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}\langle p \rangle)$ . Thus, (3.7) holds.

We continue the proof of the proposition. By Lemma 3.9 and Lemma 2.9, it suffices to show that for every  $f \in \mathbb{I}_r^\square(\chi)^{\infty p}$ , there exists a positive integer  $\Delta$  that is coprime to  $p$  such that

$$(3.9) \quad \mathbf{q}_{r,r} \widetilde{h}_{r,r} \left( \widetilde{D}_{[e]}^\diamond(-, \sigma\chi, \sigma f) \right) - \sigma \mathbf{q}_{r,r} \widetilde{h}_{r,r} \left( \widetilde{D}_{[e]}^\diamond(-, \chi, f) \right) = 0$$

holds for every  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}\langle \Delta \rangle)$ . By linearity, we may assume that  $f = \otimes_{w \nmid \infty p} f_w$  is a pure tensor. Let  $\Delta$  be a positive integer that is coprime to  $p$  such that

- (3) for every prime  $w$  not dividing  $p\Delta$ ,  $f_w$  is the unique section that is fixed by  $\widetilde{\mathcal{G}}_{2r}(\mathbb{Z}_w)$  and satisfies  $f_w(1_{4r}) = 1$ ;
- (4) for every prime  $w$  dividing  $\Delta$ ,  $f_w$  is fixed by the kernel of the map  $\widetilde{\mathcal{G}}_{2r}(\mathbb{Z}_w) \rightarrow \widetilde{\mathcal{G}}_{2r}(\mathbb{Z}_w/\Delta)$ .

<sup>8</sup>The letter  $D$  stands for pullback along the diagonal block.

By Lemma 3.9,  $\widetilde{D}_{[e]}^\diamond(-, \chi, f)$  belongs to  $\widetilde{\mathcal{A}}_{r,r,\text{hol}}^{[r]}(\widetilde{K}_{r,r}(p^d, \Delta))$ . By (2.11), Lemma 3.11 and Lemma 3.5(1,2), (3.9) holds for  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}(\Delta))$ .

The proposition is proved.  $\square$

### 3.3. Relevant representations.

**Lemma 3.13.** *Let  $\mathbb{L}/\mathbb{Q}_p$  be a finite extension and let  $\pi$  be a relevant  $\mathbb{L}$ -representation of  $G_r(\mathbb{A}_F^\infty)$  (Definition 1.1).*

- (1) *The representation  $\hat{\pi} := (\pi^\vee)^\dagger$  is also a relevant  $\mathbb{L}$ -representation of  $G_r(\mathbb{A}_F^\infty)$ .*
- (2) *The  $\mathbb{L}$ -vector space  $\text{Hom}_{G_r(\mathbb{A}_F^\infty)}(\pi, \mathcal{H}_r^{[r]} \otimes_{\mathbb{Q}_p} \mathbb{L})$  has dimension 1.*

See Definition 2.3 for the notation  $\mathcal{H}_r^{[r]}$ .

*Proof.* Part (1) follows from the fact that for every  $v \in \mathbb{V}_F^{(\infty)}$ ,  $((\pi_v^{[r]})^\vee)^\dagger$  is isomorphic to  $\pi_v^{[r]}$ .

For (2), it suffices to show that for every embedding  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ , the complex vector space  $\text{Hom}_{G_r(\mathbb{A}_F^\infty)}(\iota\pi, \mathcal{A}_{r,\text{hol}}^{[r]})$  has dimension 1. However, this follows from Arthur's multiplicity one property proved in [Mok15].  $\square$

Now we fix a relevant  $\mathbb{L}$ -representation  $\pi$  of  $G_r(\mathbb{A}_F^\infty)$  for some finite extension  $\mathbb{L}/\mathbb{Q}_p$  contained in  $\overline{\mathbb{Q}_p}$  such that  $\pi_v$  is *unramified* for every  $v \in \mathbb{V}_F^{(p)}$ . After Lemma 3.13, we let  $\mathcal{V}_\pi$  and  $\mathcal{V}_{\hat{\pi}}$  be the unique subspaces of  $\mathcal{H}_r^{[r]} \otimes_{\mathbb{Q}_p} \mathbb{L}$  that are isomorphic to  $\pi$  and  $\hat{\pi}$ , respectively.

**Notation 3.14.** We fix a  $G_r(\mathbb{A}_F^\infty)$ -invariant pairing  $\langle \cdot, \cdot \rangle_\pi: \mathcal{V}_{\hat{\pi}}^\dagger \times \mathcal{V}_\pi \rightarrow \mathbb{L}$ . For every embedding  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ , since  $\pi$  is absolutely irreducible, there is a unique element  $\mathbf{P}_\pi^\iota \in \mathbb{C}^\times$  such that

$$\int_{G_r(F) \backslash G_r(\mathbb{A}_F)} \varphi_1^\iota(g^\dagger) \varphi_2^\iota(g) dg = \mathbf{P}_\pi^\iota \cdot \langle \varphi_1^\dagger, \varphi_2 \rangle_\pi$$

for every  $\varphi_1 \in \mathcal{V}_{\hat{\pi}}$  and  $\varphi_2 \in \mathcal{V}_\pi$ . See Definition 2.3 for the notation  $\varphi_i^\iota$ .

*Remark 3.15.* Since  $\hat{\hat{\pi}} = \pi$ , the pairing  $\langle \cdot, \cdot \rangle_\pi$  is equivalent to a similar pairing  $\langle \cdot, \cdot \rangle_{\hat{\pi}}: \mathcal{V}_{\hat{\pi}}^\dagger \times \mathcal{V}_{\hat{\pi}} \rightarrow \mathbb{L}$  for  $\hat{\pi}$ , for which we have  $\mathbf{P}_{\hat{\pi}}^\iota = \mathbf{P}_\pi^\iota$  for every embedding  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ .

**Lemma 3.16.** *There is a unique  $\mathbb{L}$ -linear map*

$$\text{pr}_\pi: \mathcal{H}_r^{[r]} \otimes_{\mathbb{Q}_p} \mathbb{L} \rightarrow \mathcal{V}_\pi$$

*satisfying that for every  $\mathcal{Z} \in \mathcal{H}_r^{[r]} \otimes_{\mathbb{Q}_p} \mathbb{L}$ , every  $\varphi \in \mathcal{V}_{\hat{\pi}}$  and every embedding  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ ,*

$$\int_{G_r(F) \backslash G_r(\mathbb{A}_F)} \varphi^\iota(g^\dagger) \mathcal{Z}^\iota(g) dg = \mathbf{P}_\pi^\iota \cdot \iota \langle \varphi^\dagger, \text{pr}_\pi(\mathcal{Z}) \rangle_\pi$$

*holds.*

*Proof.* Take an open compact subgroup  $K$  of  $G_r(\mathbb{A}_F^\infty)$ . The  $\mathbb{L}$ -vector space  $\mathcal{H}_r^{[r]}(K) \otimes_{\mathbb{Q}_p} \mathbb{L}$  is a semisimple module over  $\mathbb{L}[K \backslash G_r(\mathbb{A}_F^\infty)/K]$ , in which  $\mathcal{V}_\pi(K)$  is the unique summand that is isomorphic to  $\pi^K$ . We denote by  $\mathcal{H}_r^{[r]}(K)^\pi \subseteq \mathcal{H}_r^{[r]}(K) \otimes_{\mathbb{Q}_p} \mathbb{L}$  the direct sum of the remaining summands. Then we have a direct sum decomposition  $\mathcal{H}_r^{[r]}(K) \otimes_{\mathbb{Q}_p} \mathbb{L} = \mathcal{V}_\pi(K) \oplus \mathcal{H}_r^{[r]}(K)^\pi$  of  $\mathbb{L}[K \backslash G_r(\mathbb{A}_F^\infty)/K]$ -modules. Denote by  $\text{pr}_\pi^K: \mathcal{H}_r^{[r]}(K) \otimes_{\mathbb{Q}_p} \mathbb{L} \rightarrow \mathcal{V}_\pi(K) \subseteq \mathcal{V}_\pi$  the corresponding projection map. It is clear that the maps  $\text{pr}_\pi^K$  are compatible with each other for different  $K$ , hence defining a map  $\text{pr}_\pi: \mathcal{H}_r^{[r]} \otimes_{\mathbb{Q}_p} \mathbb{L} \rightarrow \mathcal{V}_\pi$  which satisfies the property of the lemma by construction. The lemma is proved as the uniqueness is obvious.  $\square$

Take an element  $v \in \mathbb{V}_F^{(p)}$ . For every  $u \in \mathbb{P}_v$ , we have the representation  $\pi_u$  of  $\text{GL}_n(F_v)$  as a local component of  $\pi$  via the isomorphism  $G_r(F_v) \simeq \text{GL}_n(E_u) = \text{GL}_n(F_v)$ . In particular,  $\pi_u^\vee \simeq \pi_{u^c} \simeq (\pi^\vee)_u$ . Note that we will also speak of  $\pi_v$ , a representation of  $G_r(F_v)$  without any identification with  $\text{GL}_n(F_v)$ .

**Definition 3.17.** We let  $\{\alpha_{u,1}, \dots, \alpha_{u,n}\} \subseteq \overline{\mathbb{Q}_p}^\times$  (as a multi-subset) be the Satake parameter of  $\pi_u$ .

- (1) Define the *Satake polynomial* of  $\pi_u$  to be

$$\mathbf{P}_{\pi_u}(T) := \prod_{j=1}^n (T - \alpha_{u,j} \sqrt{q_v^{n-1}}).$$

(2) For every integer  $1 \leq m \leq n$ , put

$$\mathbf{A}(\pi_u, m) := \left\{ \left( \prod_{j \in J} \alpha_{u,j} \right) \sqrt{q_v}^{m(n-m)} \mid J \subseteq \{1, \dots, n\}, |J| = m \right\}$$

as a multi-subset of  $\overline{\mathbb{Q}}_p$ .

Note that to define the Satake parameter, one needs to choose a square root of  $q_v$  in  $\overline{\mathbb{Q}}_p$ . However, both  $\mathbf{P}_{\pi_u}(T)$  and  $\mathbf{A}(\pi_u, m)$  are independent of such choice.

**Lemma 3.18.** *We have*

(1) *There exist  $\beta_{u,1}, \dots, \beta_{u,n} \in O_{\mathbb{L}}$  such that*

$$\mathbf{P}_{\pi_u}(T) = T^n + \beta_{u,1} \cdot T^{n-1} + \beta_{u,2} \cdot q_v \cdot T^{n-2} + \dots + \beta_{u,r} \cdot q_v^{\frac{r(r-1)}{2}} \cdot T^r + \dots + \beta_{u,n} \cdot q_v^{\frac{n(n-1)}{2}}.$$

(2) *For every integer  $1 \leq m \leq n$ ,  $\mathbf{A}(\pi_u, m)$  is contained in  $\overline{\mathbb{Z}}_p$  and contains at most one element (with multiplicity one) in  $\overline{\mathbb{Z}}_p^{\times}$ . Moreover,  $\mathbf{A}(\pi_u, m) \cap \overline{\mathbb{Z}}_p^{\times} \neq \emptyset$  if and only if  $\beta_{u,m} \in O_{\mathbb{L}}^{\times}$ .*

(3) *We have that  $\mathbf{A}(\pi_u, m) \cap \overline{\mathbb{Z}}_p^{\times} \neq \emptyset$  is equivalent to that  $\mathbf{A}(\pi_{u^c}, n-m) \cap \overline{\mathbb{Z}}_p^{\times} \neq \emptyset$ .*

*Proof.* Part (1) follows from Definition 1.1 and [Hid98, Theorem 8.1(3)], that is, the Newton polygon is above the Hodge polygon. Part (2) is a direct consequence of (1). Part (3) follows from the fact that  $\prod_{j=1}^n \alpha_{u,j}$  is a root of unity and the fact that  $\{\alpha_{u^c,1}, \dots, \alpha_{u^c,n}\} = \{\alpha_{u,1}^{-1}, \dots, \alpha_{u,n}^{-1}\}$ .  $\square$

Put

$$I_v := \mathcal{G}_r(O_{F_v}) \times_{\mathcal{G}_r(O_{F_v}/\varpi_v)} \mathcal{P}_r(O_{F_v}/\varpi_v)$$

which is an open compact subgroup of  $G_r(F_v)$ . For every  $u \in \mathbf{P}_v$ , define two Hecke operators

$$\mathbf{T}_u^{\pm} := I_v \begin{pmatrix} \varpi_v^{\pm 1_u} \cdot 1_r & \\ & \varpi_v^{\mp 1_{u^c}} \cdot 1_r \end{pmatrix} I_v$$

(in which the volume of  $I_v$  is regarded as 1). In particular,  $\mathbf{T}_u^+ = \mathbf{U}_u \cdot I_v$  (Notation 3.7).

**Lemma 3.19.** *For every  $u \in \mathbf{P}_v$ , the multisets of generalized eigenvalues of the actions of  $\mathbf{T}_u^+$  and  $\mathbf{T}_u^-$  on  $\pi_v^{I_v}$  are  $\mathbf{A}(\pi_u, r)$  and  $\mathbf{A}(\pi_{u^c}, r)$ , respectively.*

The proof of this lemma will be given at the end of this subsection.

**Definition 3.20.** We say that the (unramified) representation  $\pi_v$  of  $G_r(F_v)$  is *Panchishkin* if  $\beta_{u,r} \in O_{\mathbb{L}}^{\times}$  for every  $u \in \mathbf{P}_v$  under the notation in Lemma 3.18.

**Lemma 3.21.** *The following statements are equivalent:*

- (1)  $\pi_v$  is Panchishkin unramified;
- (2)  $\hat{\pi}_v$  is Panchishkin unramified;
- (3)  $\mathbf{A}(\pi_u, r)$  contains a unique element in  $O_{\mathbb{L}}^{\times}$  for some  $u \in \mathbf{P}_v$ .

*Proof.* This is an immediate consequence of Lemma 3.18. The fact that the unique element in  $\mathbf{A}(\pi_u, r) \cap \overline{\mathbb{Z}}_p^{\times}$  belongs to  $\mathbb{L}$  follows from the Galois action and the uniqueness.  $\square$

**Lemma 3.22.** *Suppose that  $\pi_v$  is Panchishkin unramified.*

- (1) *The 1-dimensional subspace of  $\pi_v^{I_v}$  that is the eigenspace of the operator  $\mathbf{T}_u^+$  for the eigenvalue that is the unique element in  $\mathbf{A}(\pi_u, r) \cap O_{\mathbb{L}}^{\times}$  is independent of  $u \in \mathbf{P}_v$ .*
- (2) *The 1-dimensional subspace of  $\pi_v^{I_v}$  that is the eigenspace of the operator  $\mathbf{T}_u^-$  for the eigenvalue that is the unique element in  $\mathbf{A}(\pi_{u^c}, r) \cap O_{\mathbb{L}}^{\times}$  is independent of  $u \in \mathbf{P}_v$ .*

The proof of this lemma will be given at the end of this subsection.

**Notation 3.23.** Suppose that  $\pi_v$  is Panchishkin unramified.

- (1) For every  $u \in \mathbf{P}_v$ , we denote by  $\alpha(\pi_u) \in O_{\mathbb{L}}^{\times}$  the unique element in Lemma 3.21(2),



- (2) In view of Lemma 3.19 and Lemma 3.22, we denote by  $\pi_v^+$  and  $\pi_v^-$  the 1-dimensional subspaces of  $\pi_v^I$  that are the eigenspaces of the operators  $T_u^+$  and  $T_u^-$  for the eigenvalues  $\alpha(\pi_u)$  and  $\alpha(\pi_{u^c})$  for every  $u \in P_v$ , respectively.

**Proposition 3.24.** *Suppose that  $\pi_v$  is Panchishkin unramified.*

- (1) *For every  $u \in P_v$ , there is a unique polynomial  $Q_{\pi_u}(T) \in \mathbb{L}[T]$  that divides  $P_{\pi_u}(T)$  and has the form*

$$Q_{\pi_u}(T) = T^r + \gamma_{u,1} \cdot T^{r-1} + \gamma_{u,2} \cdot q_v \cdot T^{r-2} + \cdots + \gamma_{u,r} \cdot q_v^{\frac{r(r-1)}{2}}$$

with  $\gamma_{u,r} \in O_{\mathbb{L}}^\times$ .

- (2) *There is a unique  $\mathbb{L}$ -valued locally constant function  $\xi_{\pi_v}$  on  $G_r(F_v)$  such that*  
 (a) *there exist  $\varphi_v^\vee \in (\pi_v^\vee)^-$  and  $\varphi_v \in \pi_v^-$  such that  $\xi_{\pi_v} = \langle \pi_v^\vee(g)\varphi_v^\vee, \varphi_v \rangle_{\pi_v}$  for  $g \in G_r(F_v)$ ;*  
 (b)  $\xi_{\pi_v}(\mathfrak{w}_r) = 1$ .

*In particular,  $\xi_{\pi_v}$  is bi- $I_v$ -invariant.*

- (3) *For  $u \in P_v$ , denote by  $\underline{\pi}_u$  the unramified principal series of  $GL_r(F_v)$  with  $Q_{\pi_u}(T)$  as its Satake polynomial, which is defined over  $\mathbb{L}$ . Then there exist  $GL_r(O_{F_v})$ -invariant vectors  $\phi_u \in \underline{\pi}_u$  and  $\phi_u^\vee \in (\underline{\pi}_u)^\vee$  for every  $u \in P_v$  such that*

$$\xi_{\pi_v}(m(a)\mathfrak{w}_r) = \prod_{u \in P_v} \langle \underline{\pi}_u(a_{u^c})\phi_u, \phi_u^\vee \rangle_{(\underline{\pi}_u)^\vee}$$

*holds for every  $a = (a_u)_u \in GL_r(E_v) = \prod_{u \in P_v} GL_r(E_u)$ .*

In this rest of this subsection, we prove Lemma 3.19, Lemma 3.22 and Proposition 3.24. To ease the notation, we will suppress the subscript  $v$  hence  $P = P_v$  temporarily. We need some preparation on Jacquet modules.

For every subset  $J \subseteq \{1, \dots, n\}$ , put  $\bar{J} := \{1, \dots, n\} \setminus J$ . For every subset  $J \subseteq \{1, \dots, n\}$  of cardinality  $r$ , every  $u \in P$  and every sign  $\epsilon \in \{+, -\}$ , we denote by  $I(\alpha_{u,j} \sqrt{q}^{\epsilon r} \mid j \in J)$  the unramified principal series of  $GL_r(F)$  with the Satake parameter  $\{\alpha_{u,j} \sqrt{q}^{\epsilon r} \mid j \in J\}$ , which is defined over  $\mathbb{L}$ .

Put  $\bar{P}_r := \mathfrak{w}_r^{-1} P_r \mathfrak{w}_r$  and let  $\bar{N}_r$  be its unipotent radical. We identify both Levi quotients  $P_r/N_r$  and  $\bar{P}_r/\bar{N}_r$  with  $\text{Res}_{E/F} GL_r$  via the map  $m$  in §2.1(G4). We define the Jacquet modules

$$\begin{aligned} \pi_{N_r} &:= \pi / \{\varphi - \pi(n)\varphi \mid n \in N_r(F), \varphi \in \pi\}, \\ \pi_{\bar{N}_r} &:= \pi / \{\varphi - \pi(n)\varphi \mid n \in \bar{N}_r(F), \varphi \in \pi\}, \end{aligned}$$

which are admissible representations of  $GL_r(E)$  of finite length. Fix an order  $\{u_1, u_2\}$  of  $P$ . It is well-known that

$$\begin{aligned} \pi_{N_r}^{\text{ss}} &\simeq \bigoplus_{\substack{J \subseteq \{1, \dots, n\} \\ |J|=r}} I(\alpha_{u_1,j} \sqrt{q}^{-r} \mid j \in J) \boxtimes I(\alpha_{u_2,j} \sqrt{q}^{-r} \mid j \in \bar{J}), \\ \pi_{\bar{N}_r}^{\text{ss}} &\simeq \bigoplus_{\substack{J \subseteq \{1, \dots, n\} \\ |J|=r}} I(\alpha_{u_1,j} \sqrt{q}^r \mid j \in \bar{J}) \boxtimes I(\alpha_{u_2,j} \sqrt{q}^r \mid j \in J), \end{aligned}$$

as representations of  $GL_r(E) = GL_r(E_{u_1}) \times GL_r(E_{u_2})$ . Moreover, the isomorphism  $\mathfrak{w}_r: \pi \rightarrow \pi$  descends to an isomorphism  $\pi_{N_r}^{\text{ss}} \rightarrow \pi_{\bar{N}_r}^{\text{ss}}$  that sends  $I(\alpha_{u_1,j} \sqrt{q}^{-r} \mid j \in J) \boxtimes I(\alpha_{u_2,j} \sqrt{q}^{-r} \mid j \in \bar{J})$  to  $I(\alpha_{u_1,j} \sqrt{q}^r \mid j \in \bar{J}) \boxtimes I(\alpha_{u_2,j} \sqrt{q}^r \mid j \in J)$ .

*Proof of Lemma 3.19 and Lemma 3.22.* The element

$$\begin{pmatrix} & \varpi^{-1u^c} \cdot 1_r \\ -\varpi^{1u} \cdot 1_r & \end{pmatrix} \in G_r(F)$$

normalizes  $I$  and induces an operator on  $\pi^I$  that switches  $T_u^+$  and  $\varpi^{1u-1u^c} \cdot T_u^-$ . In particular, if the multiset of generalized eigenvalues of  $T_u^+$  on  $\pi^I$  is  $A(\pi_u, r)$ , then the multiset for  $T_u^-$  is

$$\left\{ \left( \prod_{j \in J} \alpha_{u,j}^{-1} \right) \sqrt{q}^{r^2} \mid J \subseteq \{1, \dots, n\}, |J| = r \right\},$$

which is nothing but  $A(\pi_{u^c}, r)$ . Thus, it suffices to study  $T_u^+$  in both lemmas.

The quotient map  $\pi \rightarrow \pi_{N_r}$  induces an isomorphism

$$\pi^I \xrightarrow{\sim} \pi_{N_r}^{\mathrm{GL}_r(O_E)}$$

under which the operator  $T_u^+$  (which is nothing but the operator  $U_u$  in Notation 3.7) corresponds to the operator  $q^{r^2} \cdot (\varpi^{1_u \cdot 1_r})_{1_r}$ .

Since the operator  $(\varpi^{1_u \cdot 1_r})_{1_r}$  acts on  $\mathrm{I}(\alpha_{u,j} \sqrt{q}^{-r} \mid j \in J) \boxtimes \mathrm{I}(\alpha_{u^c,j} \sqrt{q}^{-r} \mid j \in \bar{J})$  by the scalar  $\prod_{j \in J} \alpha_{u,j} \sqrt{q}^{-r^2}$ , the multiset of (generalized) eigenvalues of  $T_u^+$  on  $\pi^I$  is  $\mathbf{A}(\pi_u, r)$ . Lemma 3.19 is proved.

Now we consider Lemma 3.22, for which it suffices to show (1). For  $i = 1, 2$ , let  $J_i$  be the unique subset of  $\{1, \dots, n\}$  of cardinality  $r$  such that  $\prod_{j \in J_i} \alpha_{u_i,j} \sqrt{q}^{-r^2} \in \overline{\mathbb{Z}}_p^\times$ . Then  $J_1 \cup J_2 = \{1, \dots, n\}$ . Thus, for both  $i = 1, 2$ , the 1-dimensional subspace of  $\pi^I$  that is the eigenspace of the operator  $T_{u_i}^+$  for the eigenvalue that is the unique element in  $\mathbf{A}(\pi_{u_i}, r) \cap O_{\mathbb{L}}^\times$  is the  $\mathrm{GL}_r(O_E)$ -invariant subspace of  $\mathrm{I}(\alpha_{u_1,j} \sqrt{q}^{-r} \mid j \in J_1) \boxtimes \mathrm{I}(\alpha_{u_2,j} \sqrt{q}^{-r} \mid j \in J_2)$ . Lemma 3.22 is proved.  $\square$

*Proof of Proposition 3.24.* Without loss of generality, by Lemma 3.21, we may assume that the unique subset  $J$  of  $\{1, \dots, n\}$  with  $|J| = r$  such that  $\sqrt{q}^{r^2} \prod_{j \in J} \alpha_{u_1,j} \in O_{\mathbb{L}}^\times$  is  $\{1, \dots, r\}$ . It follows that the unique subset  $J$  of  $\{1, \dots, n\}$  with  $|J| = r$  such that  $\sqrt{q}^{r^2} \prod_{j \in J} \alpha_{u_2,j} \in O_{\mathbb{L}}^\times$  is  $\{r+1, \dots, n\}$ .

For (1), note that every factor of  $\mathcal{P}_{\pi_u}(T)$  in  $\mathbb{L}[T]$  that is monic of degree  $r$  has the form

$$\prod_{j \in J} (T - \alpha_{u,j} \sqrt{q}^{n-1})$$

for some  $J \subseteq \{1, \dots, n\}$  with  $|J| = r$ . In particular, the corresponding term  $\gamma_{u,r}$  equals  $\sqrt{q}^{r^2} \prod_{j \in J} \alpha_{u,j}$ . Thus, we must have

$$\mathcal{Q}_{\pi_{u_1}}(T) = \prod_{j=1}^r (T - \alpha_{u_1,j} \sqrt{q}^{n-1}), \quad \mathcal{Q}_{\pi_{u_2}}(T) = \prod_{j=r+1}^n (T - \alpha_{u_2,j} \sqrt{q}^{n-1}).$$

For (2) and (3), it suffices to show the following claim: For nonzero vectors  $\varphi^\vee \in (\pi^\vee)^-$  and  $\varphi \in \pi^-$ , there exist nonzero  $\mathrm{GL}_r(O_F)$ -invariant vectors  $\phi_1 \in \underline{\pi}_{u_1}$ ,  $\phi_1^\vee \in (\underline{\pi}_{u_1})^\vee$ ,  $\phi_2 \in \underline{\pi}_{u_2}$ ,  $\phi_2^\vee \in (\underline{\pi}_{u_2})^\vee$  such that

$$\langle \pi^\vee(m(a_1, a_2) \mathbf{w}_r) \varphi^\vee, \varphi \rangle_\pi = \prod_{i=1}^2 \langle \underline{\pi}_{u_i}(a_{3-i}) \phi_i, \phi_i^\vee \rangle_{(\underline{\pi}_{u_i})^\vee}$$

holds for every  $(a_1, a_2) \in \mathrm{GL}_r(E) = \mathrm{GL}_r(E_{u_1}) \times \mathrm{GL}_r(E_{u_2})$ .

Again by Lemma 3.21, the two factors

$$\begin{aligned} & \mathrm{I}(\alpha_{u_1,j} \sqrt{q}^{-r} \mid 1 \leq j \leq r) \boxtimes \mathrm{I}(\alpha_{u_2,j} \sqrt{q}^{-r} \mid r+1 \leq j \leq n), \\ & \mathrm{I}(\alpha_{u_1,j} \sqrt{q}^{-r} \mid r+1 \leq j \leq n) \boxtimes \mathrm{I}(\alpha_{u_2,j} \sqrt{q}^{-r} \mid 1 \leq j \leq r) \end{aligned}$$

are direct summands of  $\pi_{N_r}$ . We see from the proof of Lemma 3.22 that under the projection  $\pi \rightarrow \pi_{N_r}$ , the 1-dimensional subspaces  $\pi^+, \pi^- \subseteq \pi^I$  map to

$$\begin{aligned} & \mathrm{I}(\alpha_{u_1,j} \sqrt{q}^{-r} \mid 1 \leq j \leq r)^{\mathrm{GL}_r(O_F)} \boxtimes \mathrm{I}(\alpha_{u_2,j} \sqrt{q}^{-r} \mid r+1 \leq j \leq n)^{\mathrm{GL}_r(O_F)}, \\ & \mathrm{I}(\alpha_{u_1,j} \sqrt{q}^{-r} \mid r+1 \leq j \leq n)^{\mathrm{GL}_r(O_F)} \boxtimes \mathrm{I}(\alpha_{u_2,j} \sqrt{q}^{-r} \mid 1 \leq j \leq r)^{\mathrm{GL}_r(O_F)}, \end{aligned}$$

respectively. However, we observe that

$$\mathrm{I}(\alpha_{u_1,j} \sqrt{q}^{-r} \mid r+1 \leq j \leq n) \simeq (\underline{\pi}_{u_2})^\vee, \quad \mathrm{I}(\alpha_{u_2,j} \sqrt{q}^{-r} \mid 1 \leq j \leq r) \simeq (\underline{\pi}_{u_1})^\vee.$$

The claim follows.

The proposition is proved.  $\square$

**3.4. Local doubling zeta integral.** Let  $\pi$  be as in §3.3. Let  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times$  be a finite character, regarded as an automorphic character of  $\mathbb{A}_F^\times$ .

Take a finite place  $v \in \mathbb{V}_F^{\text{fin}}$ . For every  $\varphi_v^\vee \in \pi_v^\vee$ ,  $\varphi_v \in \pi_v$  and  $f \in \mathbb{I}_{r,v}^\square(\chi_v)$ , we have the local doubling zeta integral

$$(3.10) \quad Z^t(\varphi_v^\vee \otimes \varphi_v, f) := \int_{G_r(F_v)} \iota \langle \pi_v^\vee(g) \varphi_v^\vee, \varphi_v \rangle_{\pi_v} \cdot f(\mathbf{w}_r(g, 1_{2r})) \, dg$$

for every embedding  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ . Since  $\iota\pi_v$  is tempered, the above integral is absolute convergent by [Yam14, Lemma 7.2].

**Lemma 3.25.** *Define a map  $\varsigma: \text{Res}_{E/F} \text{GL}_r \times \text{Herm}_F \times \text{Herm}_F \rightarrow G_r$  by the formula*

$$\varsigma(a, u_1, u_2) := \begin{pmatrix} 1_r & u_2 \\ 0 & 1_r \end{pmatrix} \begin{pmatrix} -a & 0 \\ 0 & -{}^t a^{c,-1} \end{pmatrix} \mathbf{w}_r \begin{pmatrix} 1_r & u_1 \\ 0 & 1_r \end{pmatrix}$$

whose image is contained in the big Bruhat cell  $P_r \mathbf{w}_r N_r$ . Then we have

$$Z^t(\varphi_v^\vee \otimes \varphi_v, f) = \int_{G_r(F_v)} \iota \langle \pi_v^\vee(\varsigma(a, u_1, u_2)) \varphi_v^\vee, \varphi_v \rangle_{\pi_v} \cdot \chi(\text{Nm}_{E/F} \det a) \cdot |\det a|_E^r \cdot f \left( \mathbf{w}_r \begin{pmatrix} u_1 & {}^t a^c \\ a & u_2 \end{pmatrix} \right) \, d\varsigma(a, u_1, u_2)$$

in which the integral is absolutely convergent.

*Proof.* This formula is deduced in the proof of [LL21, Proposition 3.13].  $\square$

**Definition 3.26.** For a pair  $\varphi_v^\vee \in \pi_v^\vee$  and  $\varphi_v \in \pi_v$ , we say that an element  $\mathbf{f} \in \mathcal{S}(\text{Herm}_{2r}(F_v))$  is  $(\varphi_v^\vee, \varphi_v)$ -typical if its Fourier transform  $\widehat{\mathbf{f}} \in \mathcal{S}(\text{Herm}_{2r}(F_v))$  with respect to  $\psi_{F,v}$  satisfies

- (1)  $\widehat{\mathbf{f}}$  takes values in  $\mathbb{Q}$ ;
- (2)  $\widehat{\mathbf{f}}$  is supported on the subset

$$\left\{ \begin{pmatrix} u_1 & {}^t a^c \\ a & u_2 \end{pmatrix} \middle| a \in \text{GL}_r(O_{E,v}), u_1, u_2 \in \text{Herm}_r(O_{F,v}) \right\} \subseteq \text{Herm}_{2r}(F_v);$$

- (3)  $\widehat{\mathbf{f}}$  satisfies

$$\int_{G_r(F_v)} \langle \pi_v^\vee(\varsigma(a, u_1, u_2)) \varphi_v^\vee, \varphi_v \rangle_{\pi_v} \cdot \widehat{\mathbf{f}} \left( \begin{pmatrix} u_1 & {}^t a^c \\ a & u_2 \end{pmatrix} \right) \cdot d\varsigma(a, u_1, u_2) = 1,$$

where the integration is in fact a finite sum by (2) and  $\varsigma$  is the map in Lemma 3.25.

*Remark 3.27.* It is easy to see that  $(\varphi_v^\vee, \varphi_v)$ -typical element exists if  $\langle \pi_v^\vee(\mathbf{w}_r) \varphi_v^\vee, \varphi_v \rangle_{\pi_v} \in \mathbb{Q}^\times$ .

**Lemma 3.28.** *Consider  $\varphi_v^\vee \in \pi_v^\vee$ ,  $\varphi_v \in \pi_v$  and a  $(\varphi_v^\vee, \varphi_v)$ -typical element  $\mathbf{f} \in \mathcal{S}(\text{Herm}_{2r}(F_v))$ . If  $\chi_v$  is unramified, then  $\mathbf{f}^{X_v}$  is valued in  $\mathbb{Q}$  and*

$$Z^t(\varphi_v^\vee \otimes \varphi_v, \mathbf{f}^{X_v}) = 1$$

holds for every  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ .

*Proof.* This is immediate from Lemma 3.25 and Definition 3.26.  $\square$

This following lemma will not be used until Section 4.

**Lemma 3.29.** *For every  $\varphi_v^\vee \in \pi_v^\vee$ ,  $\varphi_v \in \pi_v$  and a  $\mathbb{Q}$ -valued section  $f \in \mathbb{I}_{r,v}^\square(\mathbf{1})$ , there exists a unique element*

$$Z(\varphi_v^\vee \otimes \varphi_v, f) \in \mathbb{L}$$

such that for every embedding  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ ,  $\iota Z(\varphi_v^\vee \otimes \varphi_v, f)$  coincides with  $Z^t(\varphi_v^\vee \otimes \varphi_v, f)$ .

*Proof.* We may regard  $\mathbb{I}_{r,v}^\square(\mathbf{1})$  as a representation with coefficients in  $\mathbb{Q}$ . Let  $\Omega$  be the set of all embeddings  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ . The assignment

$$(\varphi_v^\vee \otimes \varphi_v, f) \mapsto \{Z^t(\varphi_v^\vee \otimes \varphi_v, f)\}_{t \in \Omega}$$

defines an element

$$\mathfrak{Z} \in \text{Hom}_{G_r(F_v) \times G_r(F_v)} \left( (\pi_v^\vee \boxtimes \pi_v) \otimes \mathbb{I}_{r,v}^\square(\mathbf{1}), \mathbb{C}^\Omega \right).$$

We need to show that  $\mathfrak{Z}$  takes values in  $\mathbb{L}$ , which is tautologically a subring of  $\mathbb{C}^\Omega$ . By [LL21, Proposition 3.6(1)], it suffices to find one pair of elements  $(\varphi_v^\vee \otimes \varphi_v, f)$  such that  $\mathfrak{Z}(\varphi_v^\vee \otimes \varphi_v, f) \in \mathbb{L}^\times$ . Indeed, choose  $\varphi_v^\vee \in \pi_v^\vee$ ,  $\varphi_v \in \pi_v$

such that  $\langle \pi_v^\vee(\mathbf{w}_r)\varphi_v^\vee, \varphi_v \rangle_{\pi_v} = 1$ , and a  $(\varphi_v^\vee, \varphi_v)$ -typical element  $\mathbf{f} \in \mathcal{S}(\text{Herm}_{2r}(F_v))$  (which exists by Remark 3.27). Then by Lemma 3.28,  $\mathbf{f}^1$  is  $\mathbb{Q}$ -valued and  $\mathfrak{Z}(\varphi_v^\vee \otimes \varphi_v, \mathbf{f}^1) = 1 \in \mathbb{L}^\times$ .

The lemma is proved.  $\square$

**Lemma 3.30.** *Suppose that  $v \notin \mathbb{V}_F^{(p)}$ . If  $\pi_v$  is unramified with respect to  $K_{r,v}$  and  $\varphi_v^\vee, \varphi_v$  are both  $K_{r,v}$ -invariant such that  $\langle \varphi_v^\vee, \varphi \rangle_{\pi_v} = 1$ , then for every  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ ,*

$$Z^\iota(\varphi_v^\vee \otimes \varphi_v, f_{\chi_v}^{\text{sph}}) = \frac{L(\frac{1}{2}, \text{BC}(\iota\pi_v) \otimes (\chi_v \circ \text{Nm}_{E/F}))}{b_{2r,v}(\chi)},$$

where  $f_{\chi_v}^{\text{sph}}$  is defined in Notation 3.4(2).

*Proof.* This is a well-known calculation of Piatetski-Shapiro and Rallis. See [Li92, Theorem 3.1] for a full account including our case.  $\square$

**Proposition 3.31.** *Suppose that  $v \in \mathbb{V}_F^{(p)}$  and that  $\pi_v$  is Panchishkin unramified. For every embedding  $\iota: \mathbb{L} \rightarrow \mathbb{C}$ , we have*

$$\int_{G_r(F_v)} \iota \xi_{\pi_v}(g) \cdot (\mathbf{f}_{\chi_v}^{[0]})^{\chi_v}(\mathbf{w}_r(g, 1_{2r})) \, dg = q_v^{d_v r^2} \prod_{u \in \mathbb{P}_v} \gamma(\frac{1+\iota r}{2}, \iota \pi_u \otimes \chi_v, \psi_{F,v})^{-1}$$

where  $\xi_{\pi_v}$  and  $\pi_u$  are introduced in Proposition 3.24.

Note that the left-hand side is a local doubling zeta integral.

*Proof.* To ease notation, we omit  $v$  and  $\iota$  in the proof. In particular,  $\varpi^d$  generates the different ideal of  $F/\mathbb{Q}_p$ , and  $\xi_\pi$  is  $\mathbb{C}$ -valued.

By Lemma 3.25, we have

$$(3.11) \quad \begin{aligned} & \int_{G_r(F_v)} \iota \xi_\pi(g) \cdot (\mathbf{f}_\chi^{[0]})^\chi(\mathbf{w}_r(g, 1_{2r})) \, dg \\ &= \int_{G_r(F)} \xi_\pi(\mathcal{S}(a, u_1, u_2)) \cdot \chi(\text{Nm}_{E/F} \det a) \cdot |\text{Nm}_{E/F} \det a|_F^r \cdot \widehat{\mathbf{f}_\chi^{[0]}}(a, u_1, u_2) \cdot d\mathcal{S}(a, u_1, u_2), \end{aligned}$$

where

$$\widehat{\mathbf{f}_\chi^{[0]}}(a, u_1, u_2) := \int_{\text{Herm}_{2r}(F)} \mathbf{f}_\chi^{[0]}(T^\square) \psi_F \left( \text{tr} \begin{pmatrix} u_1 & {}^t a^c \\ a & u_2 \end{pmatrix} \begin{pmatrix} T_{11}^\square & T_{12}^\square \\ T_{21}^\square & T_{22}^\square \end{pmatrix} \right) dT^\square$$

in which  $dT^\square$  is the self-dual measure with respect to  $\psi_F$ .

It follows easily that

$$(3.12) \quad \widehat{\mathbf{f}_\chi^{[0]}}(a, u_1, u_2) = \begin{cases} q^{-dr^2} \int_{\text{GL}_r(O_E)} \chi(\text{Nm}_{E/F} \det T) \psi_F(\text{Tr}_{E/F} \text{tr } aT) \, dT & \text{if } u_1, u_2 \in \varpi^{-d} \text{Herm}_r(O_F), \\ 0 & \text{otherwise,} \end{cases}$$

in which  $dT$  is the self-dual measure on  $\text{Mat}_{r,r}(E)$  with respect to  $\psi_F$ .

Since  $\xi_\pi$  is bi- $I$ -invariant, (3.12) implies that

$$(3.11) = q^{dr^2} \int_{\text{GL}_r(E)} \xi_\pi(m(a)\mathbf{w}_r) \cdot \chi(\text{Nm}_{E/F} \det a) \cdot |\text{Nm}_{E/F} \det a|_F^r \\ \times \left( \int_{\text{GL}_r(O_E)} \chi(\text{Nm}_{E/F} \det T) \psi_F(\text{Tr}_{E/F} \text{tr } aT) \, dT \right) da,$$

which, by Proposition 3.24(3), equals

$$\begin{aligned} &= q^{dr^2} \prod_{u \in \mathbb{P}} \left( \int_{\text{GL}_r(F)} \langle \pi_u(a)\phi_u, \phi_u^\vee \rangle_{\pi_u^\vee} \cdot \chi(\det a) \cdot |\det a|_F^r \left( \int_{\text{GL}_r(O_F)} \chi(\det T) \psi_F(\text{tr } aT) \, dT \right) da \right) \\ &= q^{dr^2} \prod_{u \in \mathbb{P}} \left( \int_{\text{GL}_r(F)} \langle (\pi_u \otimes \chi)(a)\phi_u, \phi_u^\vee \rangle_{(\pi_u \otimes \chi)^\vee} \cdot |\det a|_F^r \left( \int_{\text{GL}_r(O_F)} \chi(\det T) \psi_F(\text{tr } aT) \, dT \right) da \right). \end{aligned}$$

Applying [Jac79, Proposition 1.2(3)] with  $\Phi = (\chi \circ \det) \cdot \mathbf{1}_{\mathrm{GL}_r(O_F)}$ , we have

$$\begin{aligned}
& \int_{\mathrm{GL}_r(F)} \langle (\underline{\pi}_u \otimes \chi)(a) \phi_u, \phi_u^\vee \rangle_{(\underline{\pi}_u \otimes \chi)^\vee} \cdot |\det a|_F^r \left( \int_{\mathrm{GL}_r(O_F)} \chi(\det T) \psi_F(\mathrm{tr} aT) \, dT \right) da \\
&= \gamma\left(\frac{1-r}{2}, (\underline{\pi}_u \otimes \chi)^\vee, \psi_F\right) \int_{\mathrm{GL}_r(F)} \langle \phi_u, (\underline{\pi}_u \otimes \chi)^\vee(a) \phi_u^\vee \rangle_{(\underline{\pi}_u \otimes \chi)^\vee} \cdot |\det a|_F^{\frac{1-r}{2}} \cdot \chi(\det a) \cdot \mathbf{1}_{\mathrm{GL}_r(O_F)}(a) \, da \\
&= \gamma\left(\frac{1-r}{2}, (\underline{\pi}_u \otimes \chi)^\vee, \psi_F\right) \cdot \langle \phi_u^\vee, \phi_u \rangle_{\underline{\pi}_u} \\
&= \gamma\left(\frac{1+r}{2}, \underline{\pi}_u \otimes \chi, \psi_F\right)^{-1} \cdot \langle \phi_u^\vee, \phi_u \rangle_{\underline{\pi}_u}.
\end{aligned}$$

Together, we have

$$\begin{aligned}
(3.11) &= q^{dr^2} \prod_{u \in \mathcal{P}} \gamma\left(\frac{1+r}{2}, \underline{\pi}_u \otimes \chi, \psi_F\right)^{-1} \cdot \langle \phi_u^\vee, \phi_u \rangle_{\underline{\pi}_u} \\
&= q^{dr^2} \xi_\pi(\mathbf{w}_r) \prod_{u \in \mathcal{P}} \gamma\left(\frac{1+r}{2}, \underline{\pi}_u \otimes \chi, \psi_F\right)^{-1} = q^{dr^2} \prod_{u \in \mathcal{P}} \gamma\left(\frac{1+r}{2}, \underline{\pi}_u \otimes \chi, \psi_F\right)^{-1}.
\end{aligned}$$

The proposition is proved.  $\square$

**3.5. Construction of  $p$ -adic  $L$ -function.** Let  $\pi$  be a relevant  $\mathbb{L}$ -representation of  $G_r(\mathbb{A}_F^\infty)$  for some finite extension  $\mathbb{L}/\mathbb{Q}_p$  contained in  $\overline{\mathbb{Q}}_p$  such that  $\pi_v$  is *Panchishkin unramified* for every  $v \in \mathbb{V}_F^{(p)}$ .

Choose a finite set  $\diamond$  of places of  $\mathbb{Q}$  containing  $\{\infty, p\}$  such that  $\pi_v$  is unramified for every  $v \in \mathbb{V}_F \setminus \mathbb{V}_F^{(\diamond)}$ .

We choose decomposable elements  $\varphi_1 = \otimes_v \varphi_{1,v} \in \mathcal{V}_{\hat{\pi}}$  and  $\varphi_2 = \otimes_v \varphi_{2,v} \in \mathcal{V}_\pi$  satisfying

- (T1)  $\langle \pi_v^\vee(\mathbf{w}_r) \varphi_{1,v}^\dagger, \varphi_{2,v} \rangle_{\pi_v} = 1$  for  $v \in \mathbb{V}_F^{(\diamond \setminus \{\infty, p\})}$ ,
- (T2)  $\varphi_{1,v}^\dagger \in (\pi_v^\vee)^-, \varphi_{2,v} \in \pi_v^-$  and  $\langle \pi_v^\vee(\mathbf{w}_r) \varphi_{1,v}^\dagger, \varphi_{2,v} \rangle_{\pi_v} = 1$  for  $v \in \mathbb{V}_F^{(p)}$ ,
- (T3)  $\varphi_{1,v}^\dagger \in (\pi_v^\vee)^{K_{r,v}}, \varphi_{2,v} \in \pi_v^{K_{r,v}}$  and  $\langle \varphi_{1,v}^\dagger, \varphi_{2,v} \rangle_{\pi_v} = 1$  for  $v \in \mathbb{V}_F \setminus \mathbb{V}_F^{(\diamond)}$ .

We also choose a  $(\varphi_{1,v}^\dagger, \varphi_{2,v})$ -typical element  $\mathbf{f}_v \in \mathcal{S}(\mathrm{Herm}_{2r}(F_v))$  (which exists by (T1) and Remark 3.27) for  $v \in \mathbb{V}_F^{(\diamond \setminus \{\infty, p\})}$ .

For every finite character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times$ , put

$$f_{\chi^{\infty p}} := \bigotimes_{v \in \mathbb{V}_F^{\mathrm{fin}} \setminus \mathbb{V}_F^{(p)}} f_{\chi_v} \in \mathbb{I}_r^\square(\chi)^{\infty p},$$

where  $f_{\chi_v} \in \mathbb{I}_{r,v}^\square(\chi_v)$  is the section  $\mathbf{f}_v^{\chi_v}$  (resp.  $f_{\chi_v}^{\mathrm{sph}}$ ) for  $v \in \mathbb{V}_F^{(\diamond \setminus \{\infty, p\})}$  (resp.  $v \in \mathbb{V}_F \setminus \mathbb{V}_F^{(\diamond)}$ ).

*Remark 3.32.* We have the following two remarks.

- (1) For every  $w \notin \diamond$ , we may choose a nonnegative power  $\Delta_w$  of  $w$ , such that  $\prod_{v \in \mathbb{V}_F^{(w)}} K_{m,v}$  contains

$$G_m(F \otimes \mathbb{Z}_w) \cap \widetilde{\mathcal{G}}_m(\mathbb{Z}_w) \times_{\widetilde{\mathcal{G}}_m(\mathbb{Z}_w/\Delta_w)} \widetilde{\mathcal{P}}_m(\mathbb{Z}_w/\Delta_w)$$

for every integer  $m \geq 1$ . Moreover, we may and will take  $\Delta_w = 1$  when  $w$  is unramified in  $E$ .

- (2) For every  $w \in \diamond \setminus \{\infty, p\}$ , we may choose a nonnegative power  $\Delta_w$  of  $w$  such that  $\otimes_{v \in \mathbb{V}_F^{(w)}} f_{\chi_v}$  is fixed by the kernel of the map  $\widetilde{\mathcal{G}}_{2r}(\mathbb{Z}_w) \rightarrow \widetilde{\mathcal{G}}_{2r}(\mathbb{Z}_w/\Delta_w)$  for every finite character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times$ . Indeed, by Definition 3.26(2), the restriction of  $f_{\chi_v}$  to  $K_{2r,v}$  is independent of  $\chi_v$ , which implies the existence of  $\Delta_w$ .

Consider an open compact subset  $\Omega \subseteq \Gamma_{F,p}$ . By the linear independence of characters, one can write

$$\mathbf{1}_\Omega = \sum_i c_i \cdot \chi_i$$

as a finite sum in a unique way with  $c_i \in \mathbb{C}$  and finite characters  $\chi_i: \Gamma_{F,p} \rightarrow \mathbb{C}^\times$ . For an element  $e \in \mathbb{Z}^{\mathcal{P}}$ , we put

$$\widetilde{D}_{[e]}^\diamond(-, \Omega) := \sum_i c_i \widetilde{D}_{[e]}^\diamond(-, \chi_i, f_{\chi_i^{\infty p}}),$$

where  $\widetilde{D}_{[e]}^\diamond(-, \chi_i, f_{\chi_i^{\infty p}})$  is defined in (3.6). Put

$$\Delta := \prod_{w \notin \diamond} \Delta_w, \quad \Delta' := \prod_{w \in \diamond \setminus \{\infty, p\}} \Delta_w$$

as the product of those in Remark 3.32 (1) and (2), respectively.

**Lemma 3.33.** *For every open compact subset  $\Omega \subseteq \Gamma_{F,p}$ , if  $\|e\| > 0$ , then*

$$\widetilde{h}_{r,r} \left( \widetilde{D}_{[e]}^\diamond(-, \Omega) \right) \in \varinjlim_{d \in \mathbb{N}} \widetilde{\mathcal{H}}_{r,r}^{[r]}(\widetilde{K}_{r,r}(p^d \Delta, \Delta')).$$

*Proof.* By construction and Lemma 3.9, it is clear that

$$\widetilde{h}_{r,r} \left( \widetilde{D}_{[e]}^\diamond(-, \Omega) \right) \in \varinjlim_{d \in \mathbb{N}} \widetilde{\mathcal{H}}_{r,r}^{[r]}(\widetilde{K}_{r,r}(p^d \Delta, \Delta')) \otimes_{\mathbb{Q}} \mathbb{C}.$$

It remains to show the rationality when  $\|e\| > 0$ .

Take an arbitrary element  $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q})$ . We have  $\sigma f_{\chi_v} = f_{\sigma \chi_v}$  for every  $v \in \mathbf{V}_F^{\text{fin}} \setminus \mathbf{V}_F^{(p)}$  and every finite character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times$  by construction. By Proposition 3.12, we have

$$\sigma \widetilde{h}_{r,r} \left( \widetilde{D}_{[e]}^\diamond(-, \Omega) \right) = \widetilde{h}_{r,r} \left( \sum_i \sigma(c_i) \widetilde{D}_{[e]}^\diamond(-, \sigma \chi_i, f_{\sigma \chi_i^{\infty p}}) \right).$$

On the other hand, we have  $\mathbf{1}_\Omega = \sigma \mathbf{1}_\Omega = \sum_i \sigma(c_i) \cdot \sigma \chi_i$ , which implies that

$$\widetilde{h}_{r,r} \left( \sum_i \sigma(c_i) \widetilde{D}_{[e]}^\diamond(-, \sigma \chi_i, f_{\sigma \chi_i^{\infty p}}) \right) = \widetilde{h}_{r,r} \left( \widetilde{D}_{[e]}^\diamond(-, \Omega) \right).$$

The lemma is proved.  $\square$

**Lemma 3.34.** *For every open compact subset  $\Omega \subseteq \Gamma_{F,p}$ , if  $\|e\| > 0$ , then there is a unique element*

$$\mathcal{D}_{[e]}^\diamond(-, \Omega) \in \varinjlim_{d \in \mathbb{N}} \mathcal{H}_{r,r}^{[r]}(K_{r,r}(p^d \Delta, \Delta'))$$

such that

$$\xi_{r,r}^* \mathcal{D}_{[e]}^\diamond(-, \Omega) = \zeta_{r,r}^* \widetilde{h}_{r,r} \left( \widetilde{D}_{[e]}^\diamond(-, \Omega) \right)$$

in terms of the diagram (2.6), where  $K_{r,r}(p^d \Delta, \Delta') := G_{r,r}(\mathbb{A}_F^\infty) \cap \widetilde{K}_{r,r}(p^d \Delta, \Delta')$ .

*Proof.* Since the center of  $\widetilde{G}_{2r}(\mathbb{A}^\infty)$  (as a subgroup of  $\widetilde{G}_{r,r}(\mathbb{A}^\infty)$ ) acts trivially on  $\widetilde{D}_{[e]}^\diamond(-, \Omega)$ , the element  $\zeta_{r,r}^* \widetilde{D}_{[e]}^\diamond(-, \Omega)$  descends to the desired element  $\mathcal{D}_{[e]}^\diamond(-, \Omega)$ .  $\square$

**Notation 3.35.** By Remark 2.4(2), we have a map

$$\text{pr}_{\pi, \hat{\pi}} := \text{pr}_\pi \otimes \text{pr}_{\hat{\pi}}: \mathcal{H}_{r,r}^{[r]} = \mathcal{H}_r^{[r]} \otimes_{\mathbb{Q}_p} \mathcal{H}_r^{[r]} \rightarrow \mathcal{V}_\pi \otimes_{\mathbb{L}} \mathcal{V}_{\hat{\pi}}$$

that is the tensor product of  $\text{pr}_\pi$  and  $\text{pr}_{\hat{\pi}}$  from Lemma 3.16. In what follows, for every  $\Psi \in \mathcal{H}_{r,r}^{[r]}$ ,  $\varphi_1 \in \mathcal{V}_{\hat{\pi}}$  and  $\varphi_2 \in \mathcal{V}_\pi$ , we put

$$\langle \varphi_1 \otimes \varphi_2, \Psi \rangle_{\pi, \hat{\pi}} := \left\langle \varphi_2^\dagger, \left\langle \varphi_1^\dagger, \text{pr}_{\pi, \hat{\pi}} \Psi \right\rangle_\pi \right\rangle_{\hat{\pi}}.$$

**Definition 3.36.** We define an  $\mathbb{L}$ -valued distribution  $d\mathcal{L}_p^\diamond(\pi)$  on  $\Gamma_{F,p}$  to be the following assignment

$$\Omega \subseteq \Gamma_{F,p} \mapsto C_p^{-1} \left( \prod_{u \in \mathbb{P}} \alpha(\pi_u) \right)^{-1} \left\langle \varphi_1 \otimes \varphi_2, \mathcal{D}_{[1]}^\diamond(-, \Omega) \right\rangle_{\pi, \hat{\pi}},$$

where  $C_p := \prod_{v \in \mathbf{V}_F^{(p)}} q_v^{d_v r^2}$ , which is additive from the construction. Here, 1 is regarded as a constant tuple in  $\mathbb{N}^{\mathbb{P}}$ .

**Theorem 3.37.** *The distribution  $d\mathcal{L}_p^\diamond(\pi)$  on  $\Gamma_{F,p}$  in Definition 3.36 is a  $p$ -adic measure. Moreover, if we denote by  $\mathcal{L}_p^\diamond(\pi)$  the induced (bounded) analytic function on  $\mathcal{X}_{F,p} \otimes_{\mathbb{Q}_p} \mathbb{L}$ , then for every finite (continuous) character  $\chi: \Gamma_{F,p} \rightarrow \overline{\mathbb{Q}_p}^\times$  and every isomorphism  $\iota: \overline{\mathbb{Q}_p} \xrightarrow{\sim} \mathbb{C}$ , we have*

$$\iota \mathcal{L}_p^\diamond(\pi)(\chi) = \frac{1}{P_\pi} \cdot \frac{Z_r^{[F:\mathbb{Q}]}}{b_{2r}^\diamond(\mathbf{1})} \cdot \prod_{v \in \mathbb{V}_F^{(p)}} \prod_{u \in P_v} \gamma\left(\frac{1+r}{2}, \iota(\pi_u \otimes \chi_v), \psi_{F,v}\right)^{-1} \cdot L\left(\frac{1}{2}, \text{BC}(\iota\pi^\diamond) \otimes (\iota\chi^\diamond \circ \text{Nm}_{E/F})\right),$$

where

$$Z_r := (-1)^r 2^{-2r^2} \cdot 2^{r^2-r} \pi^{r^2} \frac{\Gamma(1) \cdots \Gamma(r)}{\Gamma(r+1) \cdots \Gamma(2r)}$$

and  $\pi_u$  is introduced in Proposition 3.24. In particular, in terms of the data chosen from this subsection,  $\mathcal{L}_p^\diamond(\pi)$  depends on  $\diamond$  only, justifying its notation.

*Proof.* For the first statement, it amounts to showing that the map

$$\Omega \mapsto \int_{\Omega} d\mathcal{L}_p^\diamond(\pi) := C_p^{-1} \left( \prod_{u \in \mathbb{P}} \alpha(\pi_u) \right)^{-1} \left\langle \varphi_1 \otimes \varphi_2, \mathcal{D}_{[1]}^\diamond(-, \Omega) \right\rangle_{\pi, \hat{\pi}} \in \mathbb{L}$$

is uniformly bounded.

We first claim that for every  $\Omega$ , there is an integer  $e = e(\Omega) \in \mathbb{N}$ , regarded as a constant tuple in  $\mathbb{N}^{\mathbb{P}}$ , such that

$$(U_p^e \times U_p^e) \widetilde{D}_{[1]}^\diamond(-, \Omega) \in \widetilde{\mathcal{H}}_{r,r}^{[r]}(K_{r,r}(p\Delta, \Delta')),$$

and hence

$$(U_p^e \times U_p^e) \mathcal{D}_{[1]}^\diamond(-, \Omega) \in \mathcal{H}_{r,r}^{[r]}(K_{r,r}(p\Delta, \Delta')).$$

Indeed, this follows from the following well-known property for the operator  $U_u$  for every  $u \in P_v$  for some  $v \in \mathbb{V}_F^{(p)}$ : For every integer  $d \geq 2$ , we have  $I_v^d U_u I_v^d = I_v^{d-1} U_u I_v^d$ , where  $I_v^d := \mathcal{G}_r(O_{F_v}) \times_{\mathcal{G}_r(O_{F_v}/\varpi_v)} \mathcal{P}_r(O_{F_v}/\varpi_v^d)$ .

We fix finitely many elements  $g_1, \dots, g_s \in \widetilde{G}_{r,r}(\mathbb{A}^\diamond)$  that map surjectively to  $\widetilde{G}_{r,r}^{\text{ab}}(\mathbb{Q}) \backslash \widetilde{G}_{r,r}^{\text{ab}}(\mathbb{A}^\infty) / \widetilde{K}_{r,r}^{\text{ab}}(\Delta)$  (Remark 2.10). Then the map

$$\begin{aligned} \widetilde{\mathcal{H}}_{r,r}^{[r]}(K_{r,r}(p\Delta, \Delta')) &\rightarrow \text{SF}_{r,r}(\mathbb{C})^{\oplus s} \\ \widetilde{D} &\mapsto (\mathbf{q}_{r,r}(g_1 \cdot \widetilde{D}), \dots, \mathbf{q}_{r,r}(g_s \cdot \widetilde{D})) \end{aligned}$$

is injective.

Now we are ready to show the uniform boundedness. By (T2), we have

$$\begin{aligned} \int_{\Omega} d\mathcal{L}_p^\diamond(\pi) &= C_p^{-1} \left( \prod_{u \in \mathbb{P}} \alpha(\pi_u) \right)^{-1-2e} \left\langle (T_p^-)^e \varphi_1 \otimes (T_p^-)^e \varphi_2, \mathcal{D}_{[1]}^\diamond(-, \Omega) \right\rangle_{\pi, \hat{\pi}} \\ &= C_p^{-1} \left( \prod_{u \in \mathbb{P}} \alpha(\pi_u) \right)^{-1-2e} \left\langle \varphi_1 \otimes \varphi_2, (U_p^e \times U_p^e) \mathcal{D}_{[1]}^\diamond(-, \Omega) \right\rangle_{\pi, \hat{\pi}}, \end{aligned}$$

where  $T_p^- := \prod_{u \in \mathbb{P}} T_u^-$ . Since  $\alpha(\pi_u) \in O_{\mathbb{L}}^\times$  for every  $u \in \mathbb{P}$  and  $\mathcal{H}_{r,r}^{[r]}(K(p\Delta, \Delta'))$  is a finite-dimensional  $\mathbb{Q}_p$ -vector space, by Lemma 3.10, it suffices to show that there exists an integer  $M \geq 0$  such that

$$p^M \mathbf{q}_{r,r}(g_j \cdot \widetilde{D}_{[2e+1]}^\diamond(-, \Omega)) \in \text{SF}_{r,r}(\overline{\mathbb{Z}(p)})$$

holds for every  $1 \leq j \leq s$ , every  $e \geq 1$  and every  $\Omega$ . By (2.11), it suffices to study  $\mathbf{q}_{2r}(g_j \cdot \widetilde{E}_{[2e+1]}^\diamond(-, \Omega))$ . We may choose  $M$  such that  $p^M W_{2r}^\diamond \cdot \prod_{v \in \mathbb{V}_F^{(\diamond \setminus \{\infty, p\})}} \mathbf{f}_v(T) \in \overline{\mathbb{Z}(p)}$  for every  $T \in \text{Herm}_{2r}^\circ(F)$ . Then by Lemma 3.11 and Lemma 3.5(1),  $p^M \mathbf{q}_{2r}(g_j \cdot \widetilde{E}_{[2e+1]}^\diamond(-, \Omega)) \in \text{SF}_{2r}(\overline{\mathbb{Z}(p)})$  holds for every  $1 \leq j \leq s$ , every  $e \geq 1$  and every  $\Omega$ . Thus, we have shown that  $d\mathcal{L}_p^\diamond(\pi)$  is a  $p$ -adic measure.

Next, we show the second statement, that is, the interpolation property. By construction, Remark 3.15 and Lemma 3.10, for every finite character  $\chi: \Gamma_{F,p} \rightarrow \overline{\mathbb{Q}}_p^\times$  and every  $\iota: \overline{\mathbb{Q}}_p \xrightarrow{\sim} \mathbb{C}$ ,

$$\begin{aligned}
\iota \mathcal{L}_p^\diamond(\pi)(\chi) &= C_p^{-1} \left( \iota \prod_{u \in \mathbb{P}} \alpha(\pi_u) \right)^{-1} \frac{1}{(\mathbf{P}_\pi^\iota)^2} \iint \varphi_1^\iota(g_1^\dagger) \varphi_2^\iota(g_2^\dagger) \widetilde{E}_{[1]}^\diamond((g_1, g_2), \iota\chi, f_{\iota\chi^{\infty p}}) dg_1 dg_2 \\
&= C_p^{-1} \frac{1}{(\mathbf{P}_\pi^\iota)^2} \iint \varphi_1^\iota(g_1^\dagger) \varphi_2^\iota(g_2^\dagger) \widetilde{E}_{[0]}^\diamond((g_1, g_2), \iota\chi, f_{\iota\chi^{\infty p}}) dg_1 dg_2 \\
(3.13) \quad &= C_p^{-1} \frac{1}{(\mathbf{P}_\pi^\iota)^2} \iint (\varphi_1^\dagger)^\iota(g_1) \varphi_2^\iota(g_2) \widetilde{E}_{[0]}^\diamond((g_1, g_2^\dagger), \iota\chi, f_{\iota\chi^{\infty p}}) dg_1 dg_2,
\end{aligned}$$

where the omitted limits of double integrals are all  $(G_r(F) \backslash G_r(\mathbb{A}_F))^2$ . By (3.4),

$$\begin{aligned}
(3.13) &= C_p^{-1} \frac{1}{(\mathbf{P}_\pi^\iota)^2} \cdot b_{2r}^\diamond(\mathbf{1})^{-1} \cdot b_{2r}^\diamond(\iota\chi) \iint (\varphi_1^\dagger)^\iota(g_1) \varphi_2^\iota(g_2) \widetilde{E}((g_1, g_2^\dagger), f_\infty^{[r]} \otimes (\mathbf{f}_{\iota\chi_p}^{[0]})^{\iota\chi_p} \otimes f_{\iota\chi^{\infty p}}) dg_1 dg_2 \\
(3.14) \quad &= C_p^{-1} \frac{1}{(\mathbf{P}_\pi^\iota)^2} \cdot b_{2r}^\diamond(\mathbf{1})^{-1} \cdot b_{2r}^\diamond(\iota\chi) \iint (\varphi_1^\dagger)^\iota(g_1) \varphi_2^\iota(g_2) \widetilde{E}(\iota(g_1, g_2), f_\infty^{[r]} \otimes (\mathbf{f}_{\iota\chi_p}^{[0]})^{\iota\chi_p} \otimes f_{\iota\chi^{\infty p}}) dg_1 dg_2,
\end{aligned}$$

where we have used  $(g_1, g_2^\dagger) = \iota(g_1, g_2)$  as in Remark 3.1. By the well-known doubling integral expansion (see [Ral82] or [Liu11a, Section 2B] in the case of unitary groups) and Lemma 3.30, we have

$$\begin{aligned}
\iint (\varphi_1^\dagger)^\iota(g_1) \varphi_2^\iota(g_2) \widetilde{E}(\iota(g_1, g_2), f_\infty^{[r]} \otimes (\mathbf{f}_{\iota\chi_p}^{[0]})^{\iota\chi_p} \otimes f_{\iota\chi^{\infty p}}) dg_1 dg_2 &= \frac{L(\frac{1}{2}, \mathbf{BC}(\iota\pi^\diamond) \otimes (\iota\chi^\diamond \circ \text{Nm}_{E/F}))}{b_{2r}^\diamond(\iota\chi)} \\
&\times Z((\varphi_1^\dagger)_\infty^\iota \otimes (\varphi_2)_\infty^\iota, f_\infty^{[r]}) \cdot \prod_{v \in \mathbb{V}_F^{(p)}} Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, (\mathbf{f}_{\iota\chi_v}^{[0]})^{\iota\chi_v}) \cdot \prod_{v \in \mathbb{V}_F^{\diamond \setminus \{\infty, p\}}} Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\iota\chi_v}).
\end{aligned}$$

There are three cases:

- By [EL, Theorem 1.3 & Proposition 3.3.2] (with  $n = k = 2r$ ,  $a = b = r$ ,  $\tau_1 = \dots = \tau_r = r$ ,  $\nu_1 = \dots = \nu_r = -r$ , and  $\chi_{\text{ac}}^r = 1$ ), we have (see the proof of [LL21, Proposition 3.7] for more details)

$$Z((\varphi_1^\dagger)_\infty^\iota \otimes (\varphi_2)_\infty^\iota, f_\infty^{[r]}) = \mathbf{P}_\pi^\iota \cdot Z_r^{[F:\mathbb{Q}]}.$$

- By (T2) and Proposition 3.31, for  $v \in \mathbb{V}_F^{(p)}$ , we have

$$Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, (\mathbf{f}_{\iota\chi_v}^{[0]})^{\iota\chi_v}) = q_v^{d_v r^2} \prod_{u \in \mathbb{P}_v} \gamma(\frac{1+r}{2}, \iota(\underline{\pi}_u \otimes \chi_v), \psi_{F,v})^{-1}.$$

- By Lemma 3.28, for  $v \in \mathbb{V}_F^{\diamond \setminus \{\infty, p\}}$ , we have  $Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\iota\chi_v}) = 1$ .

Putting together, we have

$$\begin{aligned}
(3.14) &= C_p^{-1} \cdot \frac{1}{\mathbf{P}_\pi^\iota} \cdot \frac{Z_r^{[F:\mathbb{Q}]}}{b_{2r}^\diamond(\mathbf{1})} \cdot \prod_{v \in \mathbb{V}_F^{(p)}} q_v^{d_v r^2} \prod_{u \in \mathbb{P}_v} \gamma(\frac{1+r}{2}, \iota(\underline{\pi}_u \otimes \chi_v), \psi_{F,v})^{-1} \cdot L(\frac{1}{2}, \mathbf{BC}(\iota\pi^\diamond) \otimes (\iota\chi^\diamond \circ \text{Nm}_{E/F})) \\
&= \frac{1}{\mathbf{P}_\pi^\iota} \cdot \frac{Z_r^{[F:\mathbb{Q}]}}{b_{2r}^\diamond(\mathbf{1})} \cdot \prod_{v \in \mathbb{V}_F^{(p)}} \prod_{u \in \mathbb{P}_v} \gamma(\frac{1+r}{2}, \iota(\underline{\pi}_u \otimes \chi_v), \psi_{F,v})^{-1} \cdot L(\frac{1}{2}, \mathbf{BC}(\iota\pi^\diamond) \otimes (\iota\chi^\diamond \circ \text{Nm}_{E/F})).
\end{aligned}$$

The theorem is proved.  $\square$

*Remark 3.38.* It is expected that the  $p$ -adic  $L$ -function  $\mathcal{L}_p^\diamond(\pi)$  in Theorem 3.37 can be completed to the one that interpolates away-from- $p$  complex  $L$ -values, hence is independent of  $\diamond$ . In fact, for every  $v \in \mathbb{V}_F^{\text{fin}} \setminus \mathbb{V}_F^{(p)}$ , there is a natural rigid analytic function  $\mathcal{L}_p(\pi_v)$  on  $\mathcal{X}_{F,p} \otimes_{\mathbb{Q}_p} \mathbb{L}$  with possible poles such that for every finite character  $\chi: \Gamma_{F,p} \rightarrow \overline{\mathbb{Q}}_p^\times$  and every isomorphism  $\iota: \overline{\mathbb{Q}}_p \xrightarrow{\sim} \mathbb{C}$ , we have  $\iota \mathcal{L}_p(\pi_v)(\chi) = L(\frac{1}{2}, \mathbf{BC}(\iota\pi_v) \otimes (\iota\chi_v \circ \text{Nm}_{E/F}))$ . Now put

$$\mathcal{L}_p(\pi) := \mathcal{L}_p^\diamond(\pi) \prod_{v \in \mathbb{V}_F^{\diamond \setminus \{\infty, p\}}} b_{2r,v}(\mathbf{1})^{-1} \cdot \mathcal{L}_p(\pi_v).$$



We conjecture that  $\mathcal{L}_p(\pi)$  is still a bounded analytic function on  $\mathcal{X}_{F,p} \otimes_{\mathbb{Q}_p} \mathbb{L}$ . By construction, for every finite character  $\chi: \Gamma_{F,p} \rightarrow \overline{\mathbb{Q}}_p^\times$  and every isomorphism  $\iota: \overline{\mathbb{Q}}_p \xrightarrow{\sim} \mathbb{C}$ , we have

$$\iota \mathcal{L}_p(\pi)(\chi) = \frac{1}{\mathfrak{P}_\pi} \cdot \frac{Z_r^{[F:\mathbb{Q}]}}{b_{2r}^{\infty p}(\mathbf{1})} \cdot \prod_{v \in \mathbb{V}_F^{(p)}} \prod_{u \in \mathbb{P}_v} \gamma\left(\frac{1+r}{2}, \iota(\pi_u \otimes \chi_v), \psi_{F,v}\right)^{-1} \cdot L\left(\frac{1}{2}, \mathrm{BC}(\iota\pi^p) \otimes (\iota\chi^p \circ \mathrm{Nm}_{E/F})\right).$$

#### 4. $p$ -ADIC HEIGHTS OF SELMER THETA LIFTS

In this section, we introduce Selmer theta lifts and study their  $p$ -adic heights. We fix an embedding  $E \hookrightarrow \mathbb{C}$  and regard  $E$  as a subfield of  $\mathbb{C}$ . Fix an even positive integer  $n = 2r$ .

**4.1. The hermitian space.** Let  $\pi$  be a relevant  $\mathbb{L}$ -representation of  $G_r(\mathbb{A}_F^\infty)$  for some finite extension  $\mathbb{L}/\mathbb{Q}_p$  contained in  $\overline{\mathbb{Q}}_p$ .

Choose a finite set  $\diamond$  of places of  $\mathbb{Q}$  containing  $\{\infty, p\}$  such that  $\pi_v$  is unramified for every  $v \in \mathbb{V}_F \setminus \mathbb{V}_F^{(\diamond)}$ . We assume that  $\diamond$  satisfies the following extra condition:

(4.1) The set of primes of  $E$  above  $\diamond \setminus \{\infty\}$  generate the relative class group of  $E/F$ .

For later use, we put  $\underline{\diamond} := \prod_{w \in \diamond \setminus \{\infty\}} w$  which is a positive integer divisible by  $p$ .

Let  $V, (\cdot, \cdot)_V$  be a hermitian space (that is nondegenerate and  $E$ -linear in the second variable) over  $E$  of rank  $n$  that is split at every  $v \in \mathbb{V}_F \setminus \mathbb{V}_F^{(\diamond)}$ , has signature  $(n-1, 1)$  along the induced inclusion  $F \subseteq \mathbb{R}$  and signature  $(n, 0)$  at other archimedean places of  $F$ . We introduce the following sets of notation.

(H1) For every  $F$ -ring  $R$  and every integer  $m \geq 0$ , we denote by

$$T(x) := \left( (x_i, x_j)_V \right)_{i,j} \in \mathrm{Herm}_m(R)$$

the moment matrix of an element  $x = (x_1, \dots, x_m) \in V^m \otimes_F R$ .

(H2) For every  $v \in \mathbb{V}_F$ , we put  $\epsilon_v := \eta_{E/F}((-1)^r \det V_v) \in \{\pm 1\}$ . In particular,  $\epsilon_v = 1$  for  $v \notin \mathbb{V}_F^{(\diamond)}$ .

(H3) Let  $v \in \mathbb{V}_F^{\mathrm{fin}}$  be an element and  $m \geq 0$  an integer.

- For  $T \in \mathrm{Herm}_m(F_v)$ , we put  $(V_v^m)_T := \{x \in V_v^m \mid T(x) = T\}$ , and

$$(V_v^m)_{\mathrm{reg}} := \bigcup_{T \in \mathrm{Herm}_m^\circ(F_v)} (V_v^m)_T.$$

- For every  $\mathbb{Q}\langle p_v \rangle$ -ring  $\mathbb{M}$ , we have a Fourier transform map  $\widehat{\cdot}: \mathcal{S}(V_v^m, \mathbb{M}) \rightarrow \mathcal{S}(V_v^m, \mathbb{M})$  sending  $\phi$  to  $\widehat{\phi}$  defined by the formula

$$\widehat{\phi}(x) := \int_{V_v^m} \phi(y) \psi_{F,v} \left( \mathrm{Tr}_{E/F} \sum_{i=1}^m (x_i, y_i)_V \right) dy,$$

which is in fact a finite sum, where  $dy$  is the self-dual Haar measure on  $V_v^m$  with respect to  $\psi_{F,v}$ . In what follows, we will always use this self-dual Haar measure on  $V_v^m$ .

(H4) We fix an  $O_E \otimes \widehat{\mathbb{Z}}$ -lattice  $\Lambda$  in  $V \otimes_F \mathbb{A}_F^\infty$  such that for every  $v \in \mathbb{V}_F^{\mathrm{fin}} \setminus \{v \in \mathbb{V}_F^{\mathrm{ram}} \mid \text{either } \epsilon_v = -1 \text{ or } v \mid 2\}$ ,  $\Lambda_v$  is a subgroup of  $\Lambda_v^\vee$  of index  $q_v^{1-\epsilon_v}$ , where

$$\Lambda_v^\vee := \{x \in V_v \mid \mathrm{Tr}_{E/F}(x, y)_V \in \mathfrak{p}_v^{-d_v} \text{ for every } y \in \Lambda_v\}.$$

(H5) Put  $H := \mathrm{U}(V)$ , which is a reductive group over  $F$ .

(H6) Denote by  $L_0 \subseteq H(\mathbb{A}_F^\infty)$  the stabilizer of  $\Lambda$ . For every finite set  $\blacklozenge$  of places of  $\mathbb{Q}$  containing  $\diamond$ , we have the (abstract) Hecke algebra away from  $\blacklozenge$ :

$$\mathbb{T}^{\blacklozenge} := \mathbb{Z}[L_0^{\blacklozenge} \backslash H(\mathbb{A}_F^{\blacklozenge}) / L_0^{\blacklozenge}],$$

which is a (commutative) ring with the unit  $\mathbf{1}_{L_0^{\blacklozenge}}$ ; and we denote by  $\mathbb{S}^{\blacklozenge} \subseteq \mathbb{T}^{\blacklozenge}$  the subring

$$\varinjlim_{\substack{\mathbb{T} \subseteq \mathbb{V}_F^{\mathrm{spl}} \setminus \mathbb{V}_F^{(\blacklozenge)} \\ |\mathbb{T}| < \infty}} \mathbb{Z}[L_{0,\mathbb{T}} \backslash H(F_{\mathbb{T}}) / L_{0,\mathbb{T}}] \otimes \mathbf{1}_{L_0^{\blacklozenge}}.$$

(H7) For every integer  $m \geq 1$ , every  $v \in \mathbf{V}_F^{\text{fin}}$  and every  $\mathbb{Q}\langle p_v \rangle$ -ring  $\mathbb{M}$ , we have the Weil representation  $\omega_{m,v}$  of  $G_m(F_v) \times H(F_v)$  on  $\mathcal{S}(V_v^m, \mathbb{M})$  given by the following formulae:

- for  $a \in \text{GL}_m(E_v)$  and  $\phi \in \mathcal{S}(V_v^m, \mathbb{M})$ , we have

$$\omega_{m,v}(m(a))\phi(x) = |\det a|_E^r \cdot \phi(xa);$$

- for  $b \in \text{Herm}_m(F_v)$  and  $\phi \in \mathcal{S}(V_v^m, \mathbb{M})$ , we have

$$\omega_{m,v}(n(b))\phi(x) = \psi_{F,v}(\text{tr } bT(x))\phi(x);$$

- for  $\phi \in \mathcal{S}(V_v^m, \mathbb{M})$ , we have

$$\omega_{m,v}(\mathbf{w}_m)\phi(x) = \gamma_{V_v, \psi_{F,v}}^m \cdot \widehat{\phi}(x),$$

where  $\gamma_{V_v, \psi_{F,v}} \in \{\pm 1\}$  is the Weil constant of  $V_v$  with respect to  $\psi_{F,v}$ ;

- for  $h \in H(F_v)$  and  $\phi \in \mathcal{S}(V_v^m, \mathbb{M})$ , we have

$$\omega_{m,v}(h)\phi(x) = \phi(h^{-1}x).$$

(H8) When  $m = 2r$ , we have the *Siegel–Weil section map*

$$f_{-}^{\text{SW}} : \mathcal{S}(V_v^{2r}) \rightarrow \mathbf{I}_{r,v}^{\square}(\mathbf{1})$$

for  $v \in \mathbf{V}_F^{\text{fin}}$  sending  $\Phi$  to  $f_{\Phi}^{\text{SW}}$  defined by the formula

$$f_{\Phi}^{\text{SW}}(g) = (\omega_{2r,v}(g)\Phi)(0), \quad g \in G_{2r}(F_v) = G_r^{\square}(F_v).$$

(H9) For every  $v \in \mathbf{V}_F^{\text{fin}}$ , there is a unique  $\mathbb{Q}$ -valued Haar measure  $dh_v$  on  $H(F_v)$ , called the *Siegel–Weil measure*, satisfying that for every  $T^{\square} \in \text{Herm}_{2r}^{\circ}(F_v)$  and every  $\Phi \in \mathcal{S}(V_v^{2r})$ ,

$$I_{T^{\square}}(\Phi) := \int_{H(F_v)} \Phi(h_v^{-1}x) dh_v = b_{2r,v}(\mathbf{1}) \cdot W_{T^{\square}}(f_{\Phi}^{\text{SW}})$$

where  $x$  is an arbitrary element in  $(V_v^{2r})_{T^{\square}}$ . When  $v$  is unramified over  $\mathbb{Q}$  and  $H \otimes_F F_v$  is unramified, the measure  $dh_v$  gives volume 1 to every hyperspecial maximal subgroup of  $H(F_v)$ .

**Definition 4.1.** By the local theta dichotomy [GG11, Theorem 1.8], for every  $v \in \mathbf{V}_F^{\text{fin}}$ , we have a unique up to isomorphism hermitian space  $V_{\pi_v}$  over  $E_v$  of rank  $n$  such that the local theta lift of  $\pi_v$  to  $V_{\pi_v}$  does not vanishes. We say that  $V$  is  $\pi$ -coherent if  $V_v \simeq V_{\pi_v}$  for every  $v \in \mathbf{V}_F^{\text{fin}}$ .

*Remark 4.2.* It is clear that a  $\pi$ -coherent hermitian space  $V$  exists if and only if

$$(4.2) \quad \prod_{v \in \mathbf{V}_F^{\diamond}(\infty)} \eta_{E/F}((-1)^r \det V_{\pi_v}) = -(-1)^{r[F:\mathbb{Q}]}.$$

Moreover, when (4.2) holds,  $\mathcal{L}_{\rho}^{\diamond}(\pi)$  vanishes at  $\mathbf{1}$ .

At last, we recall the following definition from [LL22] and refer to [LL22, Remark 1.2] for its technical nature.

**Definition 4.3.** We define the subset  $\mathbf{V}_F^{\heartsuit}$  of  $\mathbf{V}_F^{\text{spl}} \cup \mathbf{V}_F^{\text{int}}$  consisting of  $v$  satisfying that for every  $v' \in \mathbf{V}_F^{(p_v)} \cap \mathbf{V}_F^{\text{ram}}$ , the subfield of  $\overline{F_v}$  generated by  $F_v$  and the Galois closure of  $E_{v'}$  is unramified over  $F_v$ .

In particular,  $\mathbf{V}_F^{\heartsuit}$  contains  $\mathbf{V}_F^{(p)}$ .

**4.2. Selmer theta lifts.** From this subsection, we will assume  $F \neq \mathbb{Q}$ .

Back to the setup in §4.1, take a neat open compact subgroup  $L \subseteq H(\mathbb{A}_F^{\infty})$  of the form  $L_{\diamond} L_0^{\diamond}$ . We have the Shimura variety  $X_L$  associated with  $\text{Res}_{F/\mathbb{Q}} H$  of level  $L$ , which is a smooth *projective* scheme over  $E$  of dimension  $n - 1$ . Recall that for every element  $x \in V^m \otimes_F \mathbb{A}_F^{\infty}$ , we have *Kudla's special cycle*  $Z(x)_L \in Z^m(X_L)$  if  $T(x) \in \text{Herm}_m^{\circ}(F)^+$  and  $Z(x)_L \in \text{CH}^m(X_L)_{\mathbb{Q}}$  in general. See [LL21, Section 4] for more details in our setting.

For every  $\mathbb{Q}\langle \diamond \rangle$ -ring  $\mathbb{M}$ , every  $\phi \in \mathcal{S}(V^m \otimes_F \mathbb{A}_F^{\infty}, \mathbb{M})^{K_m^{\diamond} \times L}$  and every  $T \in \text{Herm}_m(F)$ , we put

$$Z_T(\phi)_L := \sum_{\substack{x \in L \setminus V^m \otimes_F \mathbb{A}_F^{\infty} \\ T(x)=T}} \phi(x) Z(x)_L$$

as an element in  $\mathbb{M} \otimes Z^m(X_L)$  if  $T \in \text{Herm}_m^{\diamond}(F)^+$  and in  $\mathbb{M} \otimes \text{CH}^m(X_L)$  in general. Denote by

$$Z_T^{\star}(\phi)_L \in \mathbb{M} \otimes_{\mathbb{Q}} \text{H}_{\text{ét}}^{2m}(X_L, \mathbb{Q}_p(m))$$

the image of  $Z_T(\phi)_L$  under the (absolute) cycle class map  $Z^m(X_L) \rightarrow \text{CH}^m(X_L) \rightarrow \text{H}_{\text{ét}}^{2m}(X_L, \mathbb{Q}_p(m))$ .

Finally, for every  $g \in \prod_{v \in \mathbb{V}_F^{\text{fin}} \cap \mathbb{V}_F^{(\diamond)}} G_m(F_v)$ , we define the *cohomological generating function* to be

$$Z_{\phi}^{\star}(g)_L := \sum_{T \in \text{Herm}_m(F)^+} Z_T^{\star}(\omega_m(g)\phi)_L \cdot q^T \in \text{SF}_m(\mathbb{M} \otimes_{\mathbb{Q}} \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} \text{H}_{\text{ét}}^{2m}(X_L, \mathbb{Q}_p(m)),$$

where  $\omega_m$  is the restricted tensor product of  $\omega_{m,v}$  (§4.1(H7)).

**Hypothesis 4.4** (Modularity of cohomological generating functions). *For  $\phi \in \mathcal{S}(V^m \otimes_F \mathbb{A}_F^{\infty}, \mathbb{Q}\langle \diamond \rangle)^{K_m^{\diamond} \times L}$ , there exists a (unique)<sup>9</sup> element*

$$Z_{\phi,L} \in \mathcal{A}_{r,\text{hol}}^{[r]} \otimes_{\mathbb{Q}} \text{H}_{\text{ét}}^{2m}(X_L, \mathbb{Q}_p(m))$$

that is  $K_m^{\diamond}$ -invariant such that  $\mathbf{q}_r^{\text{an}}(g \cdot Z_{\phi,L}) = Z_{\phi}^{\star}(g)_L$  holds for every  $g \in \prod_{v \in \mathbb{V}_F^{\text{fin}} \cap \mathbb{V}_F^{(\diamond)}} G_m(F_v)$ .

*Remark 4.5.* Hypothesis 4.4 is implied by [LL21, Hypothesis 4.5].

**Proposition 4.6.** *For every  $\phi \in \mathcal{S}(V^m \otimes_F \mathbb{A}_F^{\infty}, \mathbb{Q}\langle \diamond \rangle)^{K_m^{\diamond} \times L}$  for which Hypothesis 4.4 holds, there exists a unique element*

$$\mathcal{Z}_{\phi,L} \in \left( \mathcal{H}_m^{[r]} \otimes_{\mathbb{Q}} \mathbb{Q}\langle \diamond \rangle \right) \otimes_{\mathbb{Q}_p} \text{H}_{\text{ét}}^{2m}(X_L, \mathbb{Q}_p(m))$$

(see Definition 2.3) such that for every embedding  $\iota: \mathbb{Q}_p \rightarrow \mathbb{C}$ ,  $\mathcal{Z}_{\phi,L}$ , regarded as an element in  $\mathcal{A}_{m,\text{hol}}^{[r]} \otimes_{\mathbb{Q}} \text{H}_{\text{ét}}^{2m}(X_L, \mathbb{Q}_p(m))$  via the inclusion  $\mathbb{Q}\langle \diamond \rangle \subseteq \mathbb{C}$ , coincides with  $Z_{\phi,L}$ .

*Proof.* Put  $G'_m := \text{Res}_{F/\mathbb{Q}} G_m$ , which has been regarded as a subgroup of  $\widetilde{G}_m$  in Notation 2.6. Put

$$\widetilde{K}_m^{\diamond} := \prod_{w \notin \diamond} \mathcal{G}_m(\mathbb{Z}_w) \times_{\widetilde{\mathcal{G}}_m(\mathbb{Z}_w/\Delta_w)} \widetilde{\mathcal{P}}_m(\mathbb{Z}_w/\Delta_w)$$

where  $\Delta_w$  is introduced in Remark 3.32(1), and  $K'_m{}^{\diamond} := G'_m(\mathbb{A}^{\diamond}) \cap \widetilde{K}_m^{\diamond}$ , which is contained in  $K_m^{\diamond}$ .

We claim that for every open compact subgroup  $K'$  of  $\prod_{w \in \diamond \setminus \{\infty\}} G'_m(\mathbb{Q}_w)$ , there exists an open compact subgroup  $\widetilde{K}$  of  $\prod_{w \in \diamond \setminus \{\infty\}} \widetilde{G}_m(\mathbb{Q}_w)$  containing  $K'$  such that the natural map

$$G'_m(\mathbb{Q}) \backslash G'_m(\mathbb{R})^{\text{ad}} \times G'_m(\mathbb{A}^{\infty}) / K' K'_m{}^{\diamond} \rightarrow \widetilde{G}_m(\mathbb{Q}) \backslash \widetilde{G}_m(\mathbb{R})^{\text{ad}} \times \widetilde{G}_m(\mathbb{A}^{\infty}) / \widetilde{K} \widetilde{K}_m^{\diamond}$$

is injective, and hence an open and closed immersion. In fact, since  $\widetilde{G}_m(\mathbb{Q})$  is discrete in  $\widetilde{G}_m(\mathbb{A}^{\infty})$ , we have

$$\varprojlim_{K' \subseteq \widetilde{K}} \widetilde{G}_m(\mathbb{Q}) \backslash \widetilde{G}_m(\mathbb{R})^{\text{ad}} \times \widetilde{G}_m(\mathbb{A}^{\infty}) / \widetilde{K} \widetilde{K}_m^{\diamond} = \widetilde{G}_m(\mathbb{Q}) \backslash \widetilde{G}_m(\mathbb{R})^{\text{ad}} \times \widetilde{G}_m(\mathbb{A}^{\infty}) / K' \widetilde{K}_m^{\diamond}$$

Then the claim follows from the obvious injectivity of the map

$$G'_m(\mathbb{Q}) \backslash G'_m(\mathbb{R})^{\text{ad}} \times G'_m(\mathbb{A}^{\infty}) / K' K'_m{}^{\diamond} \rightarrow \widetilde{G}_m(\mathbb{Q}) \backslash \widetilde{G}_m(\mathbb{R})^{\text{ad}} \times \widetilde{G}_m(\mathbb{A}^{\infty}) / K' \widetilde{K}_m^{\diamond}.$$

Choose a sufficiently small  $K'$  as above such that  $Z_{\phi,L} \in \mathcal{A}_{r,\text{hol}}^{[r]}(K' K'_m{}^{\diamond}) \otimes_{\mathbb{Q}} \text{H}_{\text{ét}}^{2m}(X_L, \mathbb{Q}_p(m))$ . By the above claim, we may extend  $Z_{\phi,L}$  by zero to obtain an element  $\widetilde{Z}_{\phi,L} \in \widetilde{\mathcal{A}}_{r,\text{hol}}^{[r]}(\widetilde{K} \widetilde{K}_m^{\diamond}) \otimes_{\mathbb{Q}} \text{H}_{\text{ét}}^{2m}(X_L, \mathbb{Q}_p(m))$  for some  $\widetilde{K}$  as above. By Lemma 2.9, we have

$$\widetilde{\mathbf{h}}_r(\widetilde{Z}_{\phi,L}) \in \left( \widetilde{\mathcal{H}}_m^{[r]} \otimes_{\mathbb{Q}} \mathbb{Q}\langle \diamond \rangle \right) \otimes_{\mathbb{Q}_p} \text{H}_{\text{ét}}^{2m}(X_L, \mathbb{Q}_p(m)).$$

It follows from the construction that, in view of (2.6), the element

$$\xi_{r^*} \zeta_r^* \widetilde{\mathbf{h}}_r(\widetilde{Z}_{\phi,L}) \in \left( \text{H}^0(\Sigma_r(K' K'_m{}^{\diamond}), \xi_{r^*}(\omega_r^{\delta})^{\otimes r}) \otimes_{\mathbb{Q}} \mathbb{Q}\langle \diamond \rangle \right) \otimes_{\mathbb{Q}_p} \text{H}_{\text{ét}}^{2m}(X_L, \mathbb{Q}_p(m))$$

belongs to the subspace

$$\left( \text{H}^0(\Sigma_r(K' K'_m{}^{\diamond}), \omega_r^{\otimes r}) \otimes_{\mathbb{Q}} \mathbb{Q}\langle \diamond \rangle \right) \otimes_{\mathbb{Q}_p} \text{H}_{\text{ét}}^{2m}(X_L, \mathbb{Q}_p(m))$$

(along the canonical subbundle  $\omega_r^{\otimes r} \subseteq \xi_{r^*}(\omega_r^{\delta})^{\otimes r}$ ). Then we define  $\mathcal{Z}_{\phi,L}$  to be  $\xi_{r^*} \zeta_r^* \widetilde{\mathbf{h}}_r(\widetilde{Z}_{\phi,L})$ , which satisfies the requirement. The proposition is proved.  $\square$

<sup>9</sup>The uniqueness follows from (4.1), which implies that  $\prod_{v \in \mathbb{V}_F^{\text{fin}} \cap \mathbb{V}_F^{(\diamond)}} G_m(F_v)$  maps surjectively to the double quotient  $G_m(F) \backslash G_m(\mathbb{A}_F^{\infty}) / K_m^{\diamond}$ .

For every finite set  $\blacklozenge$  of places of  $\mathbb{Q}$  containing  $\diamond$ , we have the Hecke characters

$$\chi_\pi^\blacklozenge, \chi_{\hat{\pi}}^\blacklozenge: \mathbb{S}_L^\blacklozenge \rightarrow \mathbb{L}$$

determined by  $\pi$  and  $\hat{\pi}$ , respectively. Put

$$V_{\pi,L} := \varinjlim_{\diamond \subseteq \blacklozenge} H_{\text{ét}}^{2r-1}(X_L \otimes_E \bar{E}, \mathbb{L}(r))[\chi_\pi^\blacklozenge], \quad V_{\hat{\pi},L} := \varinjlim_{\diamond \subseteq \blacklozenge} H_{\text{ét}}^{2r-1}(X_L \otimes_E \bar{E}, \mathbb{L}(r))[\chi_{\hat{\pi}}^\blacklozenge],$$

both being  $\mathbb{L}[\text{Gal}(\bar{E}/E)]$ -submodules of  $H_{\text{ét}}^{2r-1}(X_L \otimes_E \bar{E}, \mathbb{L}(r))$ .<sup>10</sup> The Poincaré duality induces a perfect pairing

$$V_{\hat{\pi},L} \times V_{\pi,L} \rightarrow \mathbb{L}(1).$$

*Remark 4.7.* In both [LL21] and [LL22], the authors mistakenly identified  $\chi_{\pi^\vee}^R$  with  $(\chi_\pi^R)^c$ , where  $\chi_\pi^R: \mathbb{T}_{\mathbb{Q}^{\text{ac}}}^R \rightarrow \mathbb{Q}^{\text{ac}}$  is the Hecke character in [LL21, Definition 6.8] (and similarly for  $\chi_{\pi^\vee}^R$ ); in fact, they only coincide when restricted to  $\mathbb{T}_{(\mathbb{Q}^{\text{ac}})^{c=1}}^R$ . As a consequence, one should replace  $\chi_\pi^R(s)^c$  by  $\chi_{\pi^\vee}^R(s)$  in [LL21, Proposition 6.10(1)]; and whenever one asks for two elements in  $\mathbb{S}_{\mathbb{Q}^{\text{ac}}}^R \setminus \mathfrak{m}_{\pi^\vee}^R$ , the first one should really be in  $\mathbb{S}_{\mathbb{Q}^{\text{ac}}}^R \setminus \mathfrak{m}_{\pi^\vee}^R$ . Such modifications do not affect the proof of the results.

**Lemma 4.8.** *There is a unique up to isomorphism semisimple continuous representation  $\rho_\pi$  of  $\text{Gal}(\bar{E}/E)$  of dimension  $n$  with coefficients in  $\bar{\mathbb{Q}}_p$  such that for every place  $u$  of  $E$  not above  $\diamond$  that is split over  $F$ ,  $\rho_\pi$  is unramified at  $u$  and a geometric Frobenius at  $u$  acts with a characteristic polynomial that coincides with the Satake polynomial of  $\pi_u$ , regarded as an unramified representation of  $\text{GL}_n(E_u)$ . Moreover, we have  $\rho_{\hat{\pi}} \simeq \rho_\pi^c \simeq \rho_\pi^\vee(1-n)$ .*

*Proof.* The uniqueness of  $\rho_\pi$  follows from its property and the Chebotarev density theorem; and the last statement follows from the uniqueness. It remains to show the existence of  $\rho_\pi$ .

Choose an isomorphism  $\iota: \bar{\mathbb{Q}}_p \xrightarrow{\sim} \mathbb{C}$ . By [Mok15], the automorphic base change of  $\otimes_{v \in V_F^{(\infty)}} \pi_v^{[r]} \otimes \iota\pi$  is an isobaric sum of distinct unitary cuspidal automorphic representations  $\Pi_j$  of  $\text{GL}_{n_j}(\mathbb{A}_E)$  for some partition  $n = n_1 + \cdots + n_s$ . By [CH13, Theorem 3.2.3], for each  $1 \leq j \leq s$ , we have a semisimple representation  $\rho_{\Pi_j}$  of  $\text{Gal}(\bar{E}/E)$  such that for every place  $u$  of  $E$  not above  $\diamond$  that is split over  $F$ , the restriction of  $\rho_{\Pi_j}$  to the place  $u$  is unramified and corresponds to the irreducible admissible representation  $\left(\Pi_{j,u} \otimes | \cdot |_{E_u}^{\frac{1-n}{2}}\right) \otimes_{\mathbb{C}, \iota^{-1}} \bar{\mathbb{Q}}_p$  of  $\text{GL}_{n_j}(E_u)$  under the unramified local Langlands correspondence. Then  $\rho_\pi := \bigoplus_{j=1}^s \rho_{\Pi_j}$  does the job.  $\square$

**Hypothesis 4.9.** *For every irreducible  $\mathbb{L}[\text{Gal}(\bar{E}/E)]$ -module  $\rho$  that is a subquotient of  $V_{\pi,L}$ , the base change  $\rho \otimes_{\mathbb{L}} \bar{\mathbb{Q}}_p$  is a direct summand of  $\rho_\pi(r)$ .*

*Remark 4.10.* We have the following remarks concerning Hypothesis 4.9.

- (1) Hypothesis 4.9 implies its dual statement, that is, for every irreducible  $\mathbb{L}[\text{Gal}(\bar{E}/E)]$ -module  $\rho$  that is a subquotient of  $V_{\hat{\pi},L}$ ,  $\rho \otimes_{\mathbb{L}} \bar{\mathbb{Q}}_p$  is a direct summand of  $\rho_\pi^\vee(1-r) = \rho_\pi^c(r) = \rho_{\hat{\pi}}(r)$ .
- (2) In fact, by Matsushima's formula, we have

$$V_{\pi,L} \otimes_{\mathbb{L}} \bar{\mathbb{Q}}_p = \bigoplus_{\pi'} (\pi')^L \otimes \text{Hom}_{\bar{\mathbb{Q}}_p[\mathbb{L} \backslash H(\mathbb{A}_F^\infty)/\mathbb{L}]} \left( (\pi')^L, H_{\text{ét}}^{2r-1}(X_L \otimes_E \bar{E}, \bar{\mathbb{Q}}_p(r)) \right)$$

in which the direct sum is taken over all relevant representations  $\pi'$  (with coefficients in  $\bar{\mathbb{Q}}_p$ ) that are nearly equivalent to  $\pi \otimes_{\mathbb{L}} \bar{\mathbb{Q}}_p$ . For each  $\pi'$ , put

$$\rho[\pi'] := \text{Hom}_{\bar{\mathbb{Q}}_p[\mathbb{L} \backslash H(\mathbb{A}_F^\infty)/\mathbb{L}]} \left( (\pi')^L, H_{\text{ét}}^{2r-1}(X_L \otimes_E \bar{E}, \bar{\mathbb{Q}}_p(r)) \right)$$

as an  $\bar{\mathbb{Q}}_p[\text{Gal}(\bar{E}/E)]$ -module. A precise prediction of which direct summand of  $\rho_\pi(r)$  is the semisimplification of  $\rho[\pi']$  can be found in [LL21, Hypothesis 6.6].

- (3) We understand that Hypothesis 4.9 is expected to be confirmed in a sequel of the work [KSZ].
- (4) It is conjectured that  $\rho[\pi']$  in (2) is irreducible. However, this does not seem reachable at this moment.

<sup>10</sup>In fact, by the strong multiplicity one property [Ram, Theorem A] and the local-global compatibility of base change [KMSW], we have  $V_{\pi,L} = H_{\text{ét}}^{2r-1}(X_L \otimes_E \bar{E}, \mathbb{L}(r))[\chi_\pi^\diamond]$ . However, we define it as a colimit only to emphasize its independence of  $\diamond$ .

**Lemma 4.11.** *Let  $\phi \in \mathcal{S}(V^r \otimes_F \mathbb{A}_F^\infty, \mathbb{Q}\langle \diamond \rangle)^{K_r^\diamond \times L}$  be an element for which Hypothesis 4.4 holds. For every element  $\varphi \in \mathcal{V}_\pi$ ,*

- (1) *the image of  $\langle \varphi^\dagger, \text{pr}_{\hat{\pi}}(\mathcal{Z}_{\phi,L}) \rangle_{\hat{\pi}}$  in  $\mathbb{Q}\langle \diamond \rangle \otimes_{\mathbb{Q}} \mathbb{H}_{\text{ét}}^{2r}(X_L \otimes_E \bar{E}, \mathbb{L}(r))$  vanishes;*
- (2) *assuming Hypothesis 4.9 and that  $\pi_v$  is unramified for every  $v \in \mathbb{V}_F^{(p)}$ , the induced image of  $\langle \varphi^\dagger, \text{pr}_{\hat{\pi}}(\mathcal{Z}_{\phi,L}) \rangle_{\hat{\pi}}$  in  $\mathbb{Q}\langle \diamond \rangle \otimes_{\mathbb{Q}} \mathbb{H}^1(E, \mathbb{H}_{\text{ét}}^{2r-1}(X_L \otimes_E \bar{E}, \mathbb{L}(r)))$  belongs to the subspace  $\mathbb{Q}\langle \diamond \rangle \otimes_{\mathbb{Q}} \mathbb{H}_f^1(E, \mathbb{V}_{\pi,L})$ .*

*Proof.* We may assume that  $\varphi$  is fixed by  $K_r^\diamond$ ; otherwise,  $\langle \varphi^\dagger, \text{pr}_{\hat{\pi}}(\mathcal{Z}_{\phi,L}) \rangle_{\hat{\pi}} = 0$ . By the same argument for [LL21, Proposition 6.10(1)] (see also Remark 4.7),  $\langle \varphi^\dagger, \text{pr}_{\hat{\pi}}(\mathcal{Z}_{\phi,L}) \rangle_{\hat{\pi}}$  is an eigenfunction of  $\mathbb{S}_L^\diamond$  of eigenvalue  $\chi_{\pi^\dagger}^\diamond$ . However, at every place  $v \in \mathbb{V}_F^{\text{spl}}$ ,  $\pi_v^\dagger \simeq \pi_v$ . It follows that  $\langle \varphi^\dagger, \text{pr}_{\hat{\pi}}(\mathcal{Z}_{\phi,L}) \rangle_{\hat{\pi}}$  belongs to  $\mathbb{Q}\langle \diamond \rangle \otimes_{\mathbb{Q}} \mathbb{H}_{\text{ét}}^{2r}(X_L, \mathbb{L}(r))[\chi_{\pi^\dagger}^\diamond]$ .

Part (1) follows from the fact that  $\mathbb{H}_{\text{ét}}^{2r}(X_L \otimes_E \bar{E}, \mathbb{L}(r))[\chi_{\pi^\dagger}^\diamond] = 0$ , which follows from [LL21, Proposition 6.9(1)]. Part (2) follows from Lemma 4.13 below, Lemma 4.14 below and [Nek00, Theorem 3.1].  $\square$

**Definition 4.12.** Suppose that  $\pi_v$  is unramified for every  $v \in \mathbb{V}_F^{(p)}$ . For every  $\phi \in \mathcal{S}(V^r \otimes_F \mathbb{A}_F^\infty, \mathbb{Q}\langle \diamond \rangle)^{K_r^\diamond \times L}$  for which Hypothesis 4.4 holds and every  $\varphi \in \mathcal{V}_\pi$ , we define

$$\Theta_\phi^{\text{Sel}}(\varphi)_L \in \mathbb{H}_f^1(E, \mathbb{V}_{\pi,L}) \otimes_{\mathbb{Q}} \mathbb{Q}\langle \diamond \rangle$$

to be the element induced from  $\langle \varphi^\dagger, \text{pr}_{\hat{\pi}}(\mathcal{Z}_{\phi,L}) \rangle_{\hat{\pi}}$  as in Lemma 4.11, called a *Selmer theta lift* of  $\pi$ . It is clear from the construction that  $\Theta_\phi^{\text{Sel}}(\varphi)_L$  is compatible under pullback with respect to  $L$  (of the form  $L_\diamond L_0^\diamond$ ).

In the rest of this subsection, we study some local properties of  $\mathbb{V}_{\pi,L}$ .

**Lemma 4.13.** *Assume Hypothesis 4.9. Then for every finite place  $u$  of  $E$  not above  $p$ , we have*

$$\mathbb{H}^i(E_u, \mathbb{V}_{\pi,L}) = \mathbb{H}^i(E_u, \mathbb{V}_{\hat{\pi},L}) = 0$$

for every  $i \in \mathbb{Z}$ .

*Proof.* We consider  $\mathbb{V}_{\pi,L}$  and the other case is similar. By Hypothesis 4.9, it suffices to show that  $\mathbb{H}^1(E_u, \rho_\pi(r)) = 0$  for such  $u$ . By [Car12, Theorem 1.1] and [TY07, Lemma 1.4(3)], we know that the associated Weil–Deligne representation of  $\rho_\pi(r)|_{E_u}$  is pure of weight  $-1$ , which implies that  $\mathbb{H}^i(E_u, \rho_\pi(r)) = 0$  by (the proof of) [Nek00, Proposition 2.5].  $\square$

**Lemma 4.14.** *Take  $v \in \mathbb{V}_F^{(p)}$ . If  $\pi_v$  is unramified, then  $\mathbb{V}_{\pi,L}|_{E_u}$  is crystalline for  $u$  above  $v$ .*

*Proof.* By the local-global compatibility of base change [KMSW], for every representation  $\pi'$  appearing in the direct summand in Remark 4.10(2),  $\pi'_v$  is unramified. In particular, it suffices to consider the case where  $L$  is hyperspecial at  $v$ . By [RSZ20, Theorem 4.5] (or a more closely related discussion after [LL21, Proposition 7.1]),  $X_L$  admits a finite étale cover that has smooth reduction at  $u$ . The lemma follows.  $\square$

**Lemma 4.15.** *Assume Hypothesis 4.9 and take  $v \in \mathbb{V}_F^{(p)}$ . If  $\pi_v$  is Panchishkin unramified (Definition 3.20), then  $\mathbb{V}_{\pi,L}|_{E_u}$  satisfies the Panchishkin condition (Definition A.9) and is pure of weight  $-1$  for  $u$  above  $v$ .*

*Proof.* By Lemma 4.14, Lemma A.10 and Hypothesis 4.9, it suffices to show that  $\rho_\pi(r)|_{E_u}$  satisfies the Panchishkin condition and is pure of weight  $-1$  for  $u$  above  $v$ . By [Car14, Theorem 1.1], we know that for every embedding  $\tau: E_u \rightarrow \bar{\mathbb{Q}}_p$ ,

- (1)  $\rho_\pi(r)|_{E_u}$  is crystalline and has Hodge–Tate weights  $\{-r, -r+1, \dots, r-1\}$  at  $\tau$ ;
- (2) the associated Weil–Deligne representation  $\text{WD}(\rho_\pi(r)|_{E_u})_\tau$  (see §A.6) is unramified and its multiset of generalized geometric Frobenius eigenvalues is  $\{\alpha_{v,1} \sqrt{q_v}^{-1}, \dots, \alpha_{v,n} \sqrt{q_v}^{-1}\}$ .

By (2), we know that  $\rho_\pi(r)|_{E_u}$  is pure of weight  $-1$ .

For the Panchishkin condition, by Lemma 3.21, we may assume that the unique subset  $J$  of  $\{1, \dots, n\}$  with  $|J| = r$  such that  $\sqrt{q_v}^{-r} \prod_{j \in J} \alpha_{v,j} \in \mathcal{O}_L^\times$  is  $\{1, \dots, r\}$  without loss of generality. Then  $\alpha_{v,j} \sqrt{q_v}^{-1}$  belongs to  $\bar{\mathbb{Z}}_p$  if and only if  $i \geq r+1$ . It follows that the multiset of generalized geometric Frobenius eigenvalues of  $\mathbb{D}_{\text{dR}}^-(\rho_\pi(r)|_{E_u}) \otimes_{\mathbb{L} \otimes_{\mathbb{Q}_p} E_u, 1 \otimes \tau} \bar{\mathbb{Q}}_p$  is  $\{\alpha_{v,j} \sqrt{q_v}^{-1} \mid 1 \leq j \leq r\}$  for every  $\tau: E_u \rightarrow \bar{\mathbb{Q}}_p$ . By (1), we know that  $F^0 \mathbb{D}_{\text{dR}}(\rho_\pi(r)|_{E_u})$  is a free  $\mathbb{L} \otimes_{\mathbb{Q}} E_u$ -module of rank  $r$ . Thus, it suffices to show that  $F^0 \mathbb{D}_{\text{dR}}(\rho_\pi(r)|_{E_u}) \cap \mathbb{D}_{\text{dR}}^-(\rho_\pi(r)|_{E_u}) = 0$ , which follows from the fact that  $\mathbb{D}_{\text{dR}}(\rho_\pi(r)|_{E_u})$  is (weakly) admissible as  $\rho_\pi(r)|_{E_u}$  is crystalline.

The lemma is proved.  $\square$

*Remark 4.16.* In fact, in the context of Lemma 4.15, if  $\pi_v$  is Panchishkin unramified, then  $V_{\pi,L|E_u}$  also satisfies the Panchishkin condition in [Nek93, 6.7] for  $u$  above  $v$ . Indeed,  $\mathbb{D}_{\text{dR}}(V_{\pi,L|E_u}) = \mathbb{D}_{\text{cris}}(V_{\pi,L|E_u})$  is an admissible filtered  $\varphi$ -modules, so is the  $\varphi$ -submodule  $\mathbb{D}_{\text{dR}}^-(V_{\pi,L|E_u})$  with the induced filtration. Thus, we may find subrepresentations  $X$  of  $V_{\pi,L|E_u}$  such that  $\mathbb{D}_{\text{dR}}(X) = \mathbb{D}_{\text{dR}}^-(V_{\pi,L|E_u})$ . It follows that  $0 \rightarrow X \rightarrow V_{\pi,L|E_u} \rightarrow (V_{\pi,L|E_u})/X \rightarrow 0$  is the desired exact sequence in [Nek93, 6.7].

**4.3. A  $p$ -adic arithmetic inner product formula.** We assume that  $\pi_v$  is Panchishkin unramified for every  $v \in V_F^{(p)}$  and that Hypothesis 4.9 holds for  $\pi$ .

We would like to apply §A.7 to the case where  $K = E$ ,  $X = X_L$ ,  $d = d' = r$ ,  $V = V_{\hat{\pi},L}$  and  $V' = V_{\pi,L}$ . We check the four properties: (V1) is already known; (V2) is due to Lemma 4.13; (V3) and (V4) are due to Lemma 4.14 and Lemma 4.15. As a result, we have a canonical  $p$ -adic height pairing

$$\langle \cdot, \cdot \rangle_{(V_{\hat{\pi},L}, V_{\pi,L}), E} : H_f^1(E, V_{\hat{\pi},L}) \times H_f^1(E, V_{\pi,L}) \rightarrow \Gamma_{E,p} \otimes_{\mathbb{Z}_p} \mathbb{L}.$$

Recall from [LL21, Definition 3.8] that we have a canonical volume  $\text{vol}^{\natural}(L) \in \mathbb{Q}_{>0}$ , which in fact equals the product of the constant  $W_{2r}$  in Lemma 3.2 and the volume of  $L$  under the Siegel–Weil measure in §4.1(H9). For every  $\varphi_1 \in \mathcal{V}_{\hat{\pi}}$ , every  $\varphi_2 \in \mathcal{V}_{\pi}$  and every pair  $\phi_1, \phi_2 \in \mathcal{S}(V^r \otimes_F \mathbb{A}_F^{\infty}, \mathbb{Q}\langle \diamond \rangle)^{K_r^{\diamond} \times L}$  for which Hypothesis 4.4 holds, the height

$$\text{vol}^{\natural}(L) \cdot \langle \Theta_{\phi_1}^{\text{Sel}}(\varphi_1)_L, \Theta_{\phi_2}^{\text{Sel}}(\varphi_2)_L \rangle_{(V_{\hat{\pi},L}, V_{\pi,L}), E} \in \Gamma_{E,p} \otimes_{\mathbb{Z}_p} \mathbb{L} \otimes_{\mathbb{Q}} \mathbb{Q}\langle \diamond \rangle$$

is independent of  $L$  and even  $\diamond$ . We will denote the above canonical value as  $\langle \Theta_{\phi_1}^{\text{Sel}}(\varphi_1), \Theta_{\phi_2}^{\text{Sel}}(\varphi_2) \rangle_{\pi, E}^{\natural}$ . Finally, put

$$\langle \Theta_{\phi_1}^{\text{Sel}}(\varphi_1), \Theta_{\phi_2}^{\text{Sel}}(\varphi_2) \rangle_{\pi, F}^{\natural} := \text{Nm}_{E/F} \langle \Theta_{\phi_1}^{\text{Sel}}(\varphi_1), \Theta_{\phi_2}^{\text{Sel}}(\varphi_2) \rangle_{\pi, E}^{\natural} \in \Gamma_{F,p} \otimes_{\mathbb{Z}_p} \mathbb{L} \otimes_{\mathbb{Q}} \mathbb{Q}\langle \diamond \rangle.$$

Now we can state our  $p$ -adic arithmetic inner product formula.

**Theorem 4.17.** *Suppose that  $n < p$ . Let  $\pi$  be as in Assumption 1.6 with  $r[F : \mathbb{Q}] + |\mathcal{S}_{\pi}|$  odd for which we assume Hypothesis 4.9. We also assume Hypothesis 4.4 for every element in  $\mathcal{S}(V^r \otimes_F \mathbb{A}_F^{\infty}, \mathbb{Q}\langle \diamond \rangle)^{K_r^{\diamond} \times L_0^{\diamond}}$  for every  $\diamond \subseteq \blacklozenge$ . Then for every choice of elements*

- $\varphi_1 = \otimes_v \varphi_{1,v} \in \mathcal{V}_{\hat{\pi}}$  and  $\varphi_2 = \otimes_v \varphi_{2,v} \in \mathcal{V}_{\pi}$  both fixed by  $K_r^{\diamond}$  such that  $\langle \varphi_{1,v}, \varphi_{2,v} \rangle_{\pi_v} = 1$  for every  $v \in V_F \setminus V_F^{(\diamond)}$ ,
- $\phi_1 = \otimes_v \phi_{1,v}, \phi_2 = \otimes_v \phi_{2,v} \in \mathcal{S}(V^r \otimes_F \mathbb{A}_F^{\infty}, \mathbb{Q}\langle \diamond \rangle)$  with  $\phi_{1,v} = \phi_{2,v} = \mathbf{1}_{\Lambda_v}$  for every  $v \in V_F \setminus V_F^{(\diamond)}$ ,

the identity

$$(4.3) \quad \langle \Theta_{\phi_1}^{\text{Sel}}(\varphi_1), \Theta_{\phi_2}^{\text{Sel}}(\varphi_2) \rangle_{\pi, F}^{\natural} = \partial \mathcal{L}_p^{\diamond}(\pi)(\mathbf{1}) \cdot \prod_{v \in V_F^{(p)}} \prod_{u \in P_v} \gamma(\frac{1+r}{2}, \underline{\pi}_u, \psi_{F,v}) \cdot \prod_{v \in V_F^{(\diamond \setminus \{\infty\})}} Z(\varphi_{1,v}^{\dagger} \otimes \varphi_{2,v}, f_{\phi_{1,v} \otimes \phi_{2,v}}^{\text{SW}})$$

holds in  $\Gamma_{F,p} \otimes_{\mathbb{Z}_p} \mathbb{L} \otimes_{\mathbb{Q}} \mathbb{Q}\langle \diamond \rangle$ , where

- $\gamma(\frac{1+r}{2}, \underline{\pi}_u, \psi_{F,v})$  is the unique element in  $\mathbb{L}^{\times}$  satisfying  $\iota \gamma(\frac{1+r}{2}, \underline{\pi}_u, \psi_{F,v}) = \gamma(\frac{1+r}{2}, \iota \underline{\pi}_u, \psi_{F,v})$  for every  $\iota : \mathbb{L} \rightarrow \mathbb{C}$ ;
- the term  $Z(\varphi_{1,v}^{\dagger} \otimes \varphi_{2,v}, f_{\phi_{1,v} \otimes \phi_{2,v}}^{\text{SW}}) \in \mathbb{L} \otimes_{\mathbb{Q}} \mathbb{Q}\langle \diamond \rangle$  is from Lemma 3.29.

*Remark 4.18.* We have the following remarks concerning Theorem 4.17.

- (1) The assumption that  $r[F : \mathbb{Q}] + |\mathcal{S}_{\pi}|$  is odd implies (4.2) and  $\mathcal{L}_p^{\diamond}(\pi)(\mathbf{1}) = 0$ .
- (2) By the interpolation property of  $\mathcal{L}_p^{\diamond}(\pi)$  and Lemma 3.30, the right-hand side of (4.3) does not change when enlarging  $\diamond$ . In particular, we may enlarge  $\diamond$  to prove the theorem.
- (3) Note that when we vary  $\varphi_{1,v}, \varphi_{2,v}, \phi_{1,v}, \phi_{2,v}$  for  $v \in V_F^{(\diamond \setminus \{\infty\})}$ , both sides of (4.3) define elements in the space

$$\bigotimes_{v \in V_F^{(\diamond \setminus \{\infty\})}} \text{Hom}_{G_r(F_v) \times G_r(F_v)} \left( \mathbb{I}_{r,v}^{\square}(\mathbf{1}), \pi_v \boxtimes \pi_v^{\vee} \right),$$

which is 1-dimensional if  $V$  is  $\pi$ -coherent [LL22, Proposition 4.8(1)] and vanishes if not. In particular, when  $V$  is not  $\pi$ -coherent, (4.3) holds trivially as both sides are zero.

The rest of this section is devoted to the proof of Theorem 1.7 and Theorem 4.17. In particular, we will

- assume that Assumption 1.6 holds, so that  $R_{\pi} \subseteq V_F^{(\diamond \setminus \{p\})} \cap V_F^{\text{spl}} \cap V_F^{\heartsuit}$ ,
- assume that  $V$  is  $\pi$ -coherent, so that for  $v \in V_F^{\text{fin}}$ ,  $\epsilon_v = -1$  if and only if  $v \in S_{\pi}$ ,

- assume Hypothesis 4.9,
- fix an (arbitrary) isomorphism  $\iota: \overline{\mathbb{Q}}_p \xrightarrow{\sim} \mathbb{C}$  and suppress  $\iota$  from the notation,
- assume that  $\diamond$  satisfies the following stronger condition

(4.4)

The set of primes of  $E$  above  $\mathbf{V}_F^{(\diamond \setminus \{p\})} \cap \mathbf{V}_F^{\text{spl}} \cap \mathbf{V}_F^\heartsuit$  is nonempty and generate the relative class group of  $E/F$ .

However, we do not assume Hypothesis 4.4 until we say so.

**Notation 4.19.** For every  $v \in \mathbf{V}_F^{(p)}$ , we denote by  $\varepsilon_v \in \mathbb{N}^{\mathbb{P}_v}$  the element that takes value 1 on  $\mathbb{P}_v \cap \mathbb{P}_{\text{CM}}$  (§2.1(F2)) and value 0 on  $\mathbb{P}_v \setminus \mathbb{P}_{\text{CM}}$ . Put  $\varepsilon := (\varepsilon_v)_v \in \mathbb{N}^{\mathbb{P}}$ .

We choose decomposable elements  $\varphi_1 = \otimes_v \varphi_{1,v} \in \mathcal{V}_{\hat{\pi}}$  and  $\varphi_2 = \otimes_v \varphi_{2,v} \in \mathcal{V}_\pi$  satisfying

- (T1)  $\langle \pi_v^\vee(\mathbf{w}_r) \varphi_{1,v}^\dagger, \varphi_{2,v} \rangle_{\pi_v} = 1$  for  $v \in \mathbf{V}_F^{(\diamond \setminus \{p\})} \cap \mathbf{V}_F^{\text{spl}} \cap \mathbf{V}_F^\heartsuit$ ,
- (T2)  $\varphi_{1,v}^\dagger \in (\pi_v^\vee)^-$ ,  $\varphi_{2,v} \in \pi_v^-$  and  $\langle \pi_v^\vee(\mathbf{w}_r) \varphi_{1,v}^\dagger, \varphi_{2,v} \rangle_{\pi_v} = 1$  for  $v \in \mathbf{V}_F^{(p)}$ ,
- (T3)  $\varphi_{1,v}^\dagger \in (\pi_v^\vee)^{K_{r,v}}$ ,  $\varphi_{2,v} \in \pi_v^{K_{r,v}}$  and  $\langle \varphi_{1,v}^\dagger, \varphi_{2,v} \rangle_{\pi_v} = 1$  for  $v \in \mathbf{V}_F^{\text{fin}} \setminus (\mathbf{V}_F^{(\diamond)} \cap \mathbf{V}_F^{\text{spl}} \cap \mathbf{V}_F^\heartsuit \cup \mathbb{S}_\pi)$ ,
- (T4)  $\varphi_{1,v}^\dagger, \varphi_{2,v}$  are new vectors<sup>11</sup> with respect to  $K_{r,v}$  and  $\langle \varphi_{1,v}^\dagger, \varphi_{2,v} \rangle_{\pi_v} = 1$  for  $v \in \mathbb{S}_\pi$ .

We construct some pairs of Schwartz functions in  $\mathcal{S}(V_v^r)$  for every  $v \in \mathbf{V}_F^{\text{fin}}$ .

- (S1) For  $v \in \mathbf{V}_F^{(\diamond \setminus \{p\})} \cap \mathbf{V}_F^{\text{spl}} \cap \mathbf{V}_F^\heartsuit$ , we choose  $\phi_{v,1}, \phi_{v,2} \in \mathcal{S}(V_v^r, \mathbb{Z}\langle p_v \rangle)$  that will be determined later in §4.6(R4).
- (S2) For  $v \in \mathbf{V}_F^{\text{fin}} \setminus (\mathbf{V}_F^{(\diamond)} \cap \mathbf{V}_F^{\text{spl}} \cap \mathbf{V}_F^\heartsuit)$ , put  $\phi_{v,1} = \phi_{v,2} := \mathbf{1}_{\Lambda_v^e} \in \mathcal{S}(V_v^r, \mathbb{Z})$ .
- (S3) For  $v \in \mathbf{V}_F^{(p)}$ , we choose a polarization  $\Lambda_v = \Lambda_{v,1} \oplus \mathfrak{p}_v^{-d_v} \Lambda_{v,2}$  of free  $O_{E_v}$ -modules, namely,  $\Lambda_{v,1}$  and  $\Lambda_{v,2}$  are free isotropic  $O_{E_v}$ -submodules of  $\Lambda_v$  of rank  $r$ . For  $e \in \mathbb{N}^{\mathbb{P}_v}$ , put

$$\Lambda_{v,1}^{[e]} := \left\{ x \in \left( \varpi_v^{-e-\varepsilon_v} \cdot \Lambda_{v,1} \oplus \varpi_v^{-e+\varepsilon_v^c} \cdot \Lambda_{v,2} \right)^r \mid T(x) \in \text{Herm}_r(O_{F_v}) \text{ and } x \bmod \Lambda_{v,2} \otimes \mathbb{Q} \text{ generates } \varpi_v^{-e-\varepsilon_v} \cdot \Lambda_{v,1} \right\},$$

$$\Lambda_{v,2}^{[e]} := \left\{ x \in (\varpi_v^{-e} \cdot \Lambda_{v,1} \oplus \varpi_v^{-e} \cdot \Lambda_{v,2})^r \mid T(x) \in \text{Herm}_r(O_{F_v}) \text{ and } x \bmod \Lambda_{v,1} \otimes \mathbb{Q} \text{ generates } \varpi_v^{-e} \cdot \Lambda_{v,2} \right\}.$$

For  $i = 1, 2$ , let  $\phi_{v,i}^{[e]} \in \mathcal{S}(V_v^r, \mathbb{Z})$  be the characteristic function of  $\Lambda_{v,i}^{[e]}$ .

For  $i = 1, 2$  and  $e \in \mathbb{N}^{\mathbb{P}}$ , we put

$$\phi_i^{[e]} := \left( \bigotimes_{v \in \mathbf{V}_F^{(p)}} \phi_{v,i}^{[e_v]} \right) \otimes \left( \bigotimes_{v \in \mathbf{V}_F^{\text{fin}} \setminus \mathbf{V}_F^{(p)}} \phi_{v,i} \right) \in \mathcal{S}(V^r \otimes_F \mathbb{A}_F^\infty, \mathbb{Z}\langle \diamond \rangle).$$

At last, we choose an open compact subgroup  $L_v \subseteq H(F_v)$  for every  $v \in \mathbf{V}_F^{\text{fin}}$ .

- For  $v \in \mathbf{V}_F^{(\diamond \setminus \{p\})} \cap \mathbf{V}_F^{\text{spl}} \cap \mathbf{V}_F^\heartsuit$ , we choose some  $L_v$  that fixes  $\phi_{v,i}$  for  $i = 1, 2$ .
- For  $v \in \mathbf{V}_F^{\text{fin}} \setminus (\mathbf{V}_F^{(\diamond)} \cap \mathbf{V}_F^{\text{spl}} \cap \mathbf{V}_F^\heartsuit)$ , we define  $L_v$  to be the stabilizer of  $\Lambda_v$ .
- For  $v \in \mathbf{V}_F^{(p)}$ , we define  $L_v$  to be the stabilizer of the lattice chain  $\Lambda_{v,1} \oplus \mathfrak{p}_v \Lambda_{v,2} \subseteq \Lambda_{v,1} \oplus \Lambda_{v,2}$ .

Put  $L := \prod_v L_v \subseteq H(\mathbb{A}_F^\infty)$  which is of the form  $L_\diamond L_0^\diamond$ . We may assume that  $L$  is neat by shrinking  $L_v$  for  $v \in \mathbf{V}_F^{(\diamond \setminus \{p\})} \cap \mathbf{V}_F^{\text{spl}} \cap \mathbf{V}_F^\heartsuit (\neq \emptyset)$ .

**Lemma 4.20.** Take an element  $v \in \mathbf{V}_F^{(p)}$ .

- (1) For  $i = 1, 2$  and  $e, e' \in \mathbb{N}^{\mathbb{P}_v}$ , we have  $U_v^{e'} \phi_{v,i}^{[e]} = \phi_{v,i}^{[e+e']}$ .
- (2) For  $i = 1, 2$  and  $e \in \mathbb{N}^{\mathbb{P}_v}$ ,  $\phi_{v,i}^{[e]}$  is fixed by  $L_v$ .
- (3) For every  $(e_1, e_2) \in \mathbb{N}^{\mathbb{P}_v} \times \mathbb{N}^{\mathbb{P}_v}$ , the support of  $\phi_{v,1}^{[e_1]} \otimes \phi_{v,2}^{[e_2]}$  is contained in  $(V_v^{2r})_{\text{reg}}$  (§4.1(H3)); and we have

$$(4.5) \quad f_{\phi_{v,1}^{[e_1]} \otimes \phi_{v,2}^{[e_2]}}^{\text{SW}} = b_{2r,v}(\mathbf{1})^{-1} \text{vol}(L_v, dh_v) \cdot (\mathbf{f}_{\mathbf{1}_v}^{[e_1^c + \varepsilon_v^c + e_2]})_{\mathbf{1}_v},$$

where  $\text{vol}(L_v, dh_v)$  denotes the volume of  $L_v$  under the Siegel–Weil measure  $dh_v$  in §4.1(H9).

<sup>11</sup>A new vector in an almost unramified representation of  $G_r(F_v)$  is a vector in the (1-dimensional) space in [Liu22, Definition 5.3(2)].

*Proof.* For (1), by induction, it suffices to consider the case where  $e' = 1_u$  for some  $u \in P_v$ . We will prove the case where  $i = 1$  and leave the other similar case to the reader. By definition, we have

$$\begin{aligned}
(\mathbf{U}_v^{1_u} \phi_{v,1}^{[e]})(x) &= \sum_{b \in \text{Herm}_r(O_{F_v}/\varpi_v)} (\omega_{r,v}(n(\varpi_v^{-d_v} b^\#) m(\varpi_v^{1_u})) \phi_{v,1}^{[e]})(x) \\
&= (\omega_{r,v}(m(\varpi_v^{1_u})) \phi_{v,1}^{[e]})(x) \sum_{b \in \text{Herm}_r(O_{F_v}/\varpi_v)} \psi_{F,v}(\text{tr } \varpi_v^{-d_v} b^\# T(x)) \\
(4.6) \quad &= q_v^{-r^2} \phi_{v,1}^{[e]}(\varpi_v^{1_u} x) \sum_{b \in \text{Herm}_r(O_{F_v}/\varpi_v)} \psi_{F,v}(\text{tr } \varpi_v^{-d_v} b^\# T(x)).
\end{aligned}$$

Since

$$\sum_{b \in \text{Herm}_r(O_{F_v}/\varpi_v)} \psi_{F,v}(\varpi_v^{-d_v} \text{tr } b^\# T(x)) = \begin{cases} q_v^{r^2} & \text{if } T(x) \in \text{Herm}_{2r}(O_{F_v}), \\ 0 & \text{if } T(x) \in \varpi_v^{-1} \text{Herm}_{2r}(O_{F_v}) \setminus \text{Herm}_{2r}(O_{F_v}), \end{cases}$$

we have (4.6) =  $\phi_{v,1}^{[e+1_u]}(x)$ .

For (2), by (1), it suffices to consider the case where  $e = 0$ , for which the invariance under  $L_v$  is obvious.

For (3), it is easy to see that the image of  $\Lambda_{v,1}^{[e_1]} \times \Lambda_{v,2}^{[e_2]}$  under the moment map  $T: V_v^{2r} \rightarrow \text{Herm}_{2r}(F_v)$  is contained in the set  $\mathfrak{T}_v^{[e_1^c + e_2^c + e_2]}$  in Construction 3.8, which is contained in  $\text{Herm}_{2r}^\circ(F_v)$ . For (4.5), by (1) and Lemma 3.10, it suffices to consider the case where  $e_1 = e_2 = 0$ . In the definition of  $\Lambda_{v,i}^{[0]}$ , the condition that  $T(x) \in \text{Herm}_{2r}(O_{F_v})$  is automatic. Then it is a straightforward exercise in linear algebra that the image of  $\Lambda_{v,1}^{[0]} \times \Lambda_{v,2}^{[0]}$  under the moment map  $T$  is exactly  $\mathfrak{T}_v^{[e_v^c]}$ ; and that for every  $x \in \Lambda_{v,1}^{[0]} \times \Lambda_{v,2}^{[0]}$ , an element  $h_v \in H(F_v)$  keeps  $x$  in  $\Lambda_{v,1}^{[0]} \times \Lambda_{v,2}^{[0]}$  if and only if  $h_v \in L_v$ . It follows from §4.1(H9) that

$$W_{T^\square}(\int_{\phi_{v,1}^{[0]} \otimes \phi_{v,2}^{[0]}}^{\text{SW}}) = b_{2r,v}(\mathbf{1})^{-1} \text{vol}(L_v, dh_v) \cdot \mathbf{1}_{\mathfrak{T}_v^{[e_v^c]}(T^\square)}$$

for every  $T^\square \in \text{Herm}_{2r}^\circ(F_v)$ , which implies (4.5) (when  $e_1 = e_2 = 0$ ).

The lemma is proved.  $\square$

We denote by  $(\mathbb{S}_{O_L}^\diamond)_{\pi,L}$  and  $(\mathbb{S}_{O_L}^\diamond)_{\hat{\pi},L}$  the ideals of  $\mathbb{S}_{O_L}^\diamond$  that annihilate

$$\mathbf{H}_{\text{ét}}^{2r}(X_L \otimes_E \bar{E}, \mathbb{L}(r)) \oplus \mathbf{H}_{\text{ét}}^{2r-1}(X_L \otimes_E \bar{E}, \mathbb{L}(r))/V_{\pi,L}, \quad \mathbf{H}_{\text{ét}}^{2r}(X_L \otimes_E \bar{E}, \mathbb{L}(r)) \oplus \mathbf{H}_{\text{ét}}^{2r-1}(X_L \otimes_E \bar{E}, \mathbb{L}(r))/V_{\hat{\pi},L},$$

respectively. Then by Lemma 4.13, Lemma 4.14 and [Nek00, Theorem 3.1], for every class  $Z \in \mathbf{H}^{2r}(X_L, \mathbb{L}(r))$ , the class  $s^*Z$  is naturally an element in  $\mathbf{H}_f^1(E, V_{\pi,L})$  (resp.  $\mathbf{H}_f^1(E, V_{\hat{\pi},L})$ ) if  $s \in (\mathbb{S}_{O_L}^\diamond)_{\pi,L}^2$  (resp.  $s \in (\mathbb{S}_{O_L}^\diamond)_{\hat{\pi},L}^2$ ). In particular, for  $(T_1, T_2) \in \text{Herm}_r^\circ(F)^+ \times \text{Herm}_r^\circ(F)^+$ ,  $(s_1, s_2) \in (\mathbb{S}_{O_L}^\diamond)_{\hat{\pi},L}^2 \times (\mathbb{S}_{O_L}^\diamond)_{\pi,L}^2$  and  $(e_1, e_2) \in \mathbb{N}^P \times \mathbb{N}^P$ , we have

$$Z_{T_1}^\star(s_1 \phi_1^{[e_1]})_L = s_1^* Z_{T_1}^\star(\phi_1^{[e_1]})_L \in \mathbf{H}_f^1(E, V_{\hat{\pi},L}) \otimes_{\mathbb{L}} \mathbb{C}, \quad Z_{T_2}^\star(s_2 \phi_2^{[e_2]})_L = s_2^* Z_{T_2}^\star(\phi_2^{[e_2]})_L \in \mathbf{H}_f^1(E, V_{\pi,L}) \otimes_{\mathbb{L}} \mathbb{C},$$

in which the equalities follow from [LL21, Lemma 4.4]. As a result, we may consider the  $p$ -adic height pairing

$$\langle Z_{T_1}^\star(s_1 \phi_1^{[e_1]})_L, Z_{T_2}^\star(s_2 \phi_2^{[e_2]})_L \rangle_{(V_{\hat{\pi},L}, V_{\pi,L}), E} \in \Gamma_{E,p} \otimes_{\mathbb{Z}_p} \mathbb{C}.$$

By [LL21, Lemma 6.4] (in which we may take  $R'$  as  $V_F^{(p)}$  by Lemma 4.20(3)), the algebraic cycles  $Z_{T_1}(s_1 \phi_1^{[e_1]})_L$  and  $Z_{T_2}(s_2 \phi_2^{[e_2]})_L$  do not intersect. Therefore, by the discussion in §A.7, we have a decomposition formula

$$(4.7) \quad \langle Z_{T_1}^\star(s_1 \phi_1^{[e_1]})_L, Z_{T_2}^\star(s_2 \phi_2^{[e_2]})_L \rangle_{(V_{\hat{\pi},L}, V_{\pi,L}), E} = \sum_{u \nmid \infty} \langle Z_{T_1}(s_1 \phi_1^{[e_1]})_L, Z_{T_2}(s_2 \phi_2^{[e_2]})_L \rangle_{(V_{\hat{\pi},L}, V_{\pi,L}), E_u}$$

for our  $p$ -adic height pairing. In what follows, to shorten notation, we will suppress the part  $(V_{\hat{\pi},L}, V_{\pi,L})$  in the subscript of the height pairing.

**Notation 4.21.** For a finite place  $u$  (resp.  $v$ ) of  $E$  (resp.  $F$ ) not above  $p$ , we denote by  $[u]$  (resp.  $[v]$ ) the image of an arbitrary uniformizer at  $u$  (resp.  $v$ ) in  $\Gamma_{E,p}$  (resp.  $\Gamma_{F,p}$ ).



#### 4.4. Local height away from $p$ .

**Lemma 4.22.** *For every  $v \in \mathbb{V}_F^{\text{fin}} \setminus (\mathbb{V}_F^{(\diamond)} \cap \mathbb{V}_F^{\text{spl}} \cap \mathbb{V}_F^{\heartsuit})$  and every  $T^\square \in \text{Herm}_{2r}^\circ(F_v)$ , there exists a unique element  $W_{T^\square, v} \in \mathbb{Z}[X]$  such that*

$$W_{T^\square, v}(\chi_v(\varpi_v)) = b_{2r, v}(\chi) \cdot W_{T^\square}(f_{\chi_v})$$

holds for every finite character  $\chi: \Gamma_{F, p} \rightarrow \mathbb{C}^\times$ , where  $f_{\chi_v} \in \mathbb{I}_{r, v}^\square(\chi_v)$  is the unique section that satisfies  $f_{\chi_v}|_{K_{2r, v}} = f_{\mathbf{1}_{\Lambda^{2r}}}^{\text{SW}}|_{K_{2r, v}}$  and  $\varpi_v$  is an arbitrary uniformizer of  $F_v$ .

*Proof.* When  $v \in \mathbb{V}_F^{\text{ram}}$ , this follows from [LL22, Remark 2.18 & Lemma 2.19]. When  $v \in S_\pi$ , this follows from the discussion in [LZ, Section 9]. The remaining cases have been settled in Lemma 3.5(1) as in these cases  $f_{\chi_v} = f_{\chi_v}^{\text{sph}}$  (Notation 3.4(2)).  $\square$

**Notation 4.23.** For every  $T^\square \in \text{Herm}_{2r}^\circ(F)^+$ , put  $\text{Diff}(T^\square, V) := \{v \in \mathbb{V}_F^{\text{fin}} \mid (V_v^{2r})_{T^\square} = \emptyset\}$ , which is a finite subset of  $\mathbb{V}_F^{\text{fin}} \setminus \mathbb{V}_F^{\text{spl}}$  of odd cardinality. We define  $\text{Herm}_{2r}^\circ(F)_V^+$  to be the subset of  $\text{Herm}_{2r}^\circ(F)^+$  consisting of  $T^\square$  such that  $\text{Diff}(T^\square, V)$  is a singleton, whose unique element we denote by  $v_{T^\square}$ .

**Proposition 4.24.** *There exists a pair  $(t_1, t_2) \in \mathbb{S}_{O_L}^\diamond \times \mathbb{S}_{O_L}^\diamond$  satisfying  $\chi_{\tilde{\pi}}^\diamond(t_1)\chi_\pi^\diamond(t_2) \neq 0$ , such that for every  $(T_1, T_2) \in \text{Herm}_r^\circ(F)^+ \times \text{Herm}_r^\circ(F)^+$ , every  $(s_1, s_2) \in (\mathbb{S}_{O_L}^\diamond)_{\tilde{\pi}, L}^2 \times (\mathbb{S}_{O_L}^\diamond)_{\pi, L}^2$  and every  $(e_1, e_2) \in \mathbb{N}^p \times \mathbb{N}^p$ , we have*

$$\begin{aligned} & \text{Nm}_{E/F} \left( \text{vol}^\natural(L) \sum_{u \nmid \infty p} \langle Z_{T_1}(t_1 s_1 \phi_1^{[e_1]})_L, Z_{T_2}(t_2 s_2 \phi_2^{[e_2]})_L \rangle_{E_u} \right) \\ &= W_{2r} \left( \sum_{\substack{T^\square \in \text{Herm}_{2r}^\circ(F)_V^+ \\ \partial_{r, r} T^\square = (T_1, T_2)}} W'_{T^\square, v_{T^\square}}(1) \cdot I_{T^\square}((t_1 s_1 \phi_1^{[e_1]} \otimes t_2 s_2 \phi_2^{[e_2]})^{v_{T^\square}}) \cdot [v_{T^\square}] \right) \\ &+ W_{2r} \sum_{v \in S_\pi} \frac{2}{q_v^{2r} - 1} \left( \sum_{\substack{T^\square \in \text{Herm}_{2r}^\circ(F)^+ \\ \partial_{r, r} T^\square = (T_1, T_2)}} W_{T^\square, v}^{\text{sph}}(1) \cdot I_{T^\square}((t_1 s_1 \phi_1^{[e_1]} \otimes t_2 s_2 \phi_2^{[e_2]})^v) \right) \cdot [v], \end{aligned}$$

where  $W_{2r}$  is the constant in Lemma 3.2,  $W_{T^\square, v}^{\text{sph}} \in \mathbb{Z}[X]$  is the polynomial in Lemma 3.5(1), and  $I_{T^\square}$  is (the product of) the functional in §4.1(H9).

*Proof.* We first note that by Theorem A.4, the local  $p$ -adic height at  $u \nmid \infty p$  coincides with Beilinson's local index. To compute the local indices at different  $u$ , we have four cases:

Suppose that  $u$  lies over  $\mathbb{V}_F^{\text{spl}}$ . By [LL22, Proposition 4.20] (see also Remark 4.7), we can find a pair  $(t_1^u, t_2^u) \in \mathbb{S}_{O_L}^\diamond \times \mathbb{S}_{O_L}^\diamond$  satisfying  $\chi_{\tilde{\pi}}^\diamond(t_1^u)\chi_\pi^\diamond(t_2^u) \neq 0$  such that  $\langle Z_{T_1}(t_1^u s_1 \phi_1^{[e_1]})_L, Z_{T_2}(t_2^u s_2 \phi_2^{[e_2]})_L \rangle_{E_u} = 0$ . Moreover, we may take  $t_1^u = t_2^u = 1$  for all but finitely many  $u$ .

Suppose that  $u$  lies over an element  $v \in \mathbb{V}_F^{\text{unr}} \setminus S_\pi$ . By [LL21, Proposition 8.1] and Remark A.5, we have

$$\begin{aligned} & \text{vol}^\natural(L) \cdot \langle Z_{T_1}(s_1 \phi_1^{[e_1]})_L, Z_{T_2}(s_2 \phi_2^{[e_2]})_L \rangle_{E_u} \\ &= -W_{2r} \left( \sum_{\substack{T^\square \in \text{Herm}_{2r}^\circ(F)_V^+ \\ \partial_{r, r} T^\square = (T_1, T_2) \\ v_{T^\square} = v}} \frac{b_{2r, v}(\mathbf{1})}{\log q_v} \cdot W'_{T^\square}(0, 1_{4r}, \mathbf{1}_{\Lambda_v^{2r}}) \cdot I_{T^\square}((s_1 \phi_1^{[e_1]} \otimes s_2 \phi_2^{[e_2]})^v) \right) \cdot [u], \end{aligned}$$

where  $W_{T^\square}(s, 1_{4r}, \mathbf{1}_{\Lambda_v^{2r}})$  denotes the usual Siegel–Whittaker function with complex variable  $s$  (see [LL21, (3.3)] for example). In our case, the character  $\chi_v$  plays the role as  $|\cdot|_{F_v}^s$ , which implies that

$$(4.8) \quad W_{T^\square, v}(q_v^{-s}) = \prod_{i=1}^n L(s+i, \eta_{E/F, v}^{n-i}) \cdot W_{T^\square}(s, 1_{4r}, \mathbf{1}_{\Lambda_v^{2r}}).$$

Together with the relation  $\text{Nm}_{E/F}[u] = 2[v]$ , we obtain

(4.9)

$$\text{Nm}_{E/F} \left( \text{vol}^{\natural}(L) \cdot \langle Z_{T_1}(s_1 \phi_1^{[e_1]}), Z_{T_2}(s_2 \phi_2^{[e_2]}) \rangle_{L, E_u} \right) = W_{2r} \left( \sum_{\substack{T^{\square} \in \text{Herm}_{2r}^{\circ}(F)_V^+ \\ \partial_{r,r} T^{\square} = (T_1, T_2) \\ v_{T^{\square}} = v}} \mathbf{W}'_{T^{\square}}(1) \cdot I_{T^{\square}}((s_1 \phi_1^{[e_1]} \otimes s_2 \phi_2^{[e_2]})^v) \right) \cdot [v].$$

Suppose that  $u$  lies over an element  $v \in \mathbf{V}_F^{\text{ram}}$ . By [LL22, Proposition 4.28] and Remark A.5, we have

$$\begin{aligned} & \text{vol}^{\natural}(L) \cdot \langle Z_{T_1}(s_1 \phi_1^{[e_1]}), Z_{T_2}(s_2 \phi_2^{[e_2]}) \rangle_{L, E_u} \\ &= -W_{2r} \left( \sum_{\substack{T^{\square} \in \text{Herm}_{2r}^{\circ}(F)_V^+ \\ \partial_{r,r} T^{\square} = (T_1, T_2) \\ v_{T^{\square}} = v}} \frac{b_{2r,v}(\mathbf{1})}{\log q_v} \cdot \mathbf{W}'_{T^{\square}}(0, 1_{4r}, \mathbf{1}_{\Lambda_v^{2r}}) \cdot I_{T^{\square}}((s_1 \phi_1^{[e_1]} \otimes s_2 \phi_2^{[e_2]})^v) \right) \cdot [u]. \end{aligned}$$

Now we have (4.8) again but  $\text{Nm}_{E/F}[u] = [v]$ , which imply (4.9) as well.

Suppose that  $u$  lies over an element  $v \in \mathbf{S}_{\pi}$ . By [LL21, Proposition 9.1] (see also Remark 4.7) and Remark A.5, we can find a pair  $(t_1^u, t_2^u) \in \mathbb{S}_{O_L}^{\diamond} \times \mathbb{S}_{O_L}^{\diamond}$  satisfying  $\chi_{\pi}^{\diamond}(t_1^u) \chi_{\pi}^{\diamond}(t_2^u) \neq 0$  such that

$$\begin{aligned} & \text{vol}^{\natural}(L) \cdot \langle Z_{T_1}(t_1^u s_1 \phi_1^{[e_1]}), Z_{T_2}(t_2^u s_2 \phi_2^{[e_2]}) \rangle_{L, E_u} \\ &= -W_{2r} \left( \sum_{\substack{T^{\square} \in \text{Herm}_{2r}^{\circ}(F)_V^+ \\ \partial_{r,r} T^{\square} = (T_1, T_2) \\ v_{T^{\square}} = v}} \frac{b_{2r,v}(\mathbf{1})}{\log q_v^2} \cdot \mathbf{W}'_{T^{\square}}(0, 1_{4r}, \mathbf{1}_{\Lambda_v^{2r}}) \cdot I_{T^{\square}}((t_1^u s_1 \phi_1^{[e_1]} \otimes t_2^u s_2 \phi_2^{[e_2]})^v) \right) \cdot [u] \\ &+ W_{2r} \frac{1}{q_v^{2r} - 1} \left( \sum_{\substack{T^{\square} \in \text{Herm}_{2r}^{\circ}(F)^+ \\ \partial_{r,r} T^{\square} = (T_1, T_2)}} \mathbf{W}_{T^{\square},v}^{\text{sph}}(1) \cdot I_{T^{\square}}((t_1^u s_1 \phi_1^{[e_1]} \otimes t_2^u s_2 \phi_2^{[e_2]})^v) \right) \cdot [u]. \end{aligned}$$

Now we have (4.8) and  $\text{Nm}_{E/F}[u] = 2[v]$ , which imply

$$\begin{aligned} & \text{Nm}_{E/F} \left( \text{vol}^{\natural}(L) \cdot \langle Z_{T_1}(t_1^u s_1 \phi_1^{[e_1]}), Z_{T_2}(t_2^u s_2 \phi_2^{[e_2]}) \rangle_{L, E_u} \right) \\ &= W_{2r} \left( \sum_{\substack{T^{\square} \in \text{Herm}_{2r}^{\circ}(F)_V^+ \\ \partial_{r,r} T^{\square} = (T_1, T_2) \\ v_{T^{\square}} = v}} \mathbf{W}'_{T^{\square},v}(1) \cdot I_{T^{\square}}((t_1^u s_1 \phi_1^{[e_1]} \otimes t_2^u s_2 \phi_2^{[e_2]})^v) \right) \cdot [v] \\ &+ W_{2r} \frac{2}{q_v^{2r} - 1} \left( \sum_{\substack{T^{\square} \in \text{Herm}_{2r}^{\circ}(F)^+ \\ \partial_{r,r} T^{\square} = (T_1, T_2)}} \mathbf{W}_{T^{\square},v}^{\text{sph}}(1) \cdot I_{T^{\square}}((t_1^u s_1 \phi_1^{[e_1]} \otimes t_2^u s_2 \phi_2^{[e_2]})^v) \right) \cdot [v]. \end{aligned}$$

Finally, for  $i = 1, 2$ , we take  $t_i = \prod_u t_i^u$  to be the (finite) product of the above auxiliary Hecke operators. The proposition follows by taking the sum over all  $u \nmid \infty p$ , which is a finite sum.  $\square$

**4.5. Local height above  $p$ .** Take an element  $u \in \mathbf{P}$  with  $v \in \mathbf{V}_F^{(p)}$  its underlying place. For technical purposes, we fix an  $E$ -linear isomorphism  $\overline{E}_u \xrightarrow{\sim} \mathbb{C}$ .

**Lemma 4.25.** *Suppose that  $n < p$ . There exists a pair  $(t_1, t_2) \in \mathbb{S}_{O_L}^\diamond \times \mathbb{S}_{O_L}^\diamond$  satisfying  $\chi_\pi^\diamond(t_1)\chi_\pi^\diamond(t_2) \neq 0$ , such that for every  $(T_1, T_2) \in \text{Herm}_r^\circ(F)^+ \times \text{Herm}_r^\circ(F)^+$ , every  $(s_1, s_2) \in (\mathbb{S}_{O_L}^\diamond)_{\pi, L}^2 \times (\mathbb{S}_{O_L}^\diamond)_{\pi, L}^2$  and every  $(e_1, e_2) \in \mathbb{N}^P \times \mathbb{N}^P$ , we have*

$$\langle Z_{T_1}(t_1 s_1 \phi_1^{[e_1]})_L, Z_{T_2}(t_2 s_2 \phi_2^{[e_2]})_L \rangle_{E_u} \in (O_{E_u}^\times)^{\text{fr}} \otimes_{\mathbb{Z}_p} \overline{\mathbb{Q}}_p.$$

*Proof.* In view of Remark A.11, we would like to apply Theorem A.6, for which we need an integral model of  $X_L$  over  $O_{E_u}$ . For this, we need an auxiliary Shimura variety that admits such a model via moduli interpretation. Choose a CM type  $\Phi$  of  $E$  such that the induced  $p$ -adic places of  $E$  of  $\Phi$  via the fixed isomorphism  $\overline{E_u} \xrightarrow{\sim} \mathbb{C}$  form a subset  $\mathbf{P}_\Phi$  of  $\mathbf{P}$  of cardinality  $[F : \mathbb{Q}]$  that contains  $u$ . Then the reflex field  $E_\Phi \subseteq \mathbb{C}$  of  $\Phi$  is contained in  $E_u$ . Recall that we have the  $\mathbb{Q}$ -torus  $T$  from §2.2 and fix a neat open compact subgroup  $K_T$  of  $T(\mathbb{A}^\infty)$  that is maximal at primes not in  $\diamond \setminus \{p\}$ . We have the Shimura variety  $Y_{K_T}$  of  $T$  with respect to the CM type  $\Phi$  at level  $K_T$ , which is finite étale over  $\text{Spec } E_\Phi$ . Put  $X := (X_L \otimes_E E_u) \otimes_{E_\Phi} Y_{K_T}$ , which is a finite étale cover of  $X_L \otimes_E E_u$  and hence a smooth projective scheme over  $E_u$  of pure dimension  $n - 1$ . The ring  $\mathbb{S}^\diamond$  extends naturally to a ring of finite étale correspondences (see §A.1) of  $X$ . For every  $x \in V^r \otimes_F \mathbb{A}_F^\infty$ , we denote by  $Z(x)$  the pullback of  $Z(x)_L$  to  $X$ .

Now for the lemma, it suffices to find elements  $(t_1, t_2) \in \mathbb{S}_{O_L}^\diamond \times \mathbb{S}_{O_L}^\diamond$  satisfying  $\chi_\pi^\diamond(t_1)\chi_\pi^\diamond(t_2) \neq 0$ , such that for every  $x_1, x_2 \in V^r \otimes_F \mathbb{A}_F^\infty$  satisfying

$$(4.10) \quad T(x_i) \in \text{Herm}_r^\circ(F)^+, \quad x_{i,v} \in \bigcup_{e \in \mathbb{N}^{P_v}} \Lambda_{v,i}^{[e]}, \quad i = 1, 2,$$

we have  $\langle t_1^* Z(x_1), t_2^* Z(x_2) \rangle_{X, E_u} \in O_{E_u}^\times \otimes_{\mathbb{Z}_p} \mathbb{L}$ .

Put  $K := E_u$  with the residue field  $\kappa$ . The  $K$ -scheme  $X$  admits an integral model  $\mathcal{X}$  over  $O_K$  such that for every  $S \in \text{Sch}'_{/O_K}$ ,  $\mathcal{X}(S)$  is the set of equivalence classes (given by  $p$ -principal isogenies) of tuples  $(A_0, \lambda_0, \eta_0; A, \lambda, \eta; G_{u^c} \rightarrow G_{A, u^c})$  where

- $A_0$  is an abelian scheme over  $S$  with an action of  $O_E$  of signature type  $\Phi$ , together with a compatible  $p$ -principal polarization  $\lambda_0$  and a level structure  $\eta_0$  away from  $p$ ,
- $A$  is an abelian scheme over  $S$  with an action of  $O_E$  of signature type  $n\Phi - \text{inc} + \text{inc}^c$  (inc being the inclusion  $E \hookrightarrow \mathbb{C}$ ), together with a compatible  $p$ -principal polarization  $\lambda$ , so that  $G_{A, u^c} := A[(u^c)^\infty]$  is an  $O_{F_v}$ -divisible module of dimension 1 and relative height  $n$ ,
- $\eta$  is an  $L^v$ -level structure for the hermitian space  $\text{Hom}_{O_E}(A_0, A) \otimes_F \mathbb{A}_F^{\infty v}$ ,
- $G_{u^c} \rightarrow G_{A, u^c}$  is an isogeny of  $O_{F_v}$ -divisible modules over  $S$  whose kernel is contained in  $G_{u^c}[\varpi_v]$  and has degree  $q_v^r$ .

The reader may consult [LL21, Section 7] for more details about the first three items, which are insensitive to our argument below. By the same argument for [TY07, Proposition 3.4], we know that  $\mathcal{X}$  is a projective strictly semistable scheme over  $O_K$  to which finite étale correspondences in  $\mathbb{S}_{O_L}^\diamond$  naturally extend. Moreover, if we put  $X := \mathcal{X} \otimes_{O_K} \kappa$  and let  $X_1$  (resp.  $X_2$ ) be the closed locus of  $X$  on which the kernel of  $G_{u^c} \rightarrow G_{A, u^c}$  (resp.  $G_{A, u^c} \rightarrow G_{u^c}/G_{u^c}[\varpi_v]$ ) is not étale, then under the notation of §A.5,

$$X^{(1)} = X_1 \bigsqcup X_2, \quad X^{(2)} = X_1 \bigcap X_2, \quad X^{(3)} = X^{(4)} = \dots = \emptyset.$$

We then would like to apply Theorem A.6 with  $\mathbb{T} = \mathbb{S}_{O_L}^\diamond$ ,  $\mathfrak{m} = \text{Ker } \chi_\pi^\diamond$  and  $\mathfrak{m}' = \text{Ker } \chi_\pi^\diamond$ . To check (A.2), we realize that both  $\chi_\pi^\diamond$  and  $\chi_\pi^\diamond$  can be defined over a number field  $\mathbb{E}$  contained in  $\mathbb{L}$ . Thus, by [KM74, Theorem 2], it suffices to show that

$$(4.11) \quad \bigoplus_{q \geq 0} H_{\text{ét}}^q(X^{(2)} \otimes_\kappa \bar{\kappa}, \mathbb{E}_\ell)_\mathfrak{m} = \bigoplus_{q \geq 0} H_{\text{ét}}^q(X^{(2)} \otimes_\kappa \bar{\kappa}, \mathbb{E}_\ell)_{\mathfrak{m}'} = 0$$

where  $\ell$  is an arbitrary prime of  $\mathbb{E}$  not above  $p$ . Indeed, there is a finite flat morphism  $\mathcal{X}_1 \rightarrow \mathcal{X}$  to which finite étale correspondences in  $\mathbb{S}_{O_L}^\diamond$  naturally extend, in which  $\mathcal{X}_1$  is the integral model with a Drinfeld level-1 structure at  $v$  as the one used in [LL21, Section 7]. Then (4.11) follows from claim (2) in the proof of [LL21, Lemma 7.3] with  $m = j = 1$ .

Denote by  $\mathcal{Z}(x)$  the Zariski closure of  $Z(x)$  in  $\mathcal{X}$ . By Theorem A.6 and Remark A.11, it suffices to show the following two claims for the lemma.

- (1) For every  $x_1, x_2 \in V^r \otimes_F \mathbb{A}_F^\infty$  satisfying (4.10) and every  $t_1, t_2 \in \mathbb{S}_{O_L}^\diamond$ , we have  $t_1^* \mathcal{Z}(x_1) \cap t_2^* \mathcal{Z}(x_2) = \emptyset$ .
- (2) For every  $x \in V^r \otimes_F \mathbb{A}_F^\infty$  with  $T(x) \in \text{Herm}_r^\circ(F)^+$ , the dimension of  $\mathcal{Z}(x) \cap X^{(h)}$  is at most  $r - h$  for  $h = 1, 2$ .

For (1), we may assume  $t_1 = t_2 = 1$ . Take an object  $(A_0, \lambda_0, \eta_0; A, \lambda, \eta; G_{u^c} \rightarrow G_{A,u^c})$  of  $\mathcal{X}$ . We put  $G_{A,u} := A[u^\infty]$ , which admits an isomorphism  $\lambda[u^\infty]: G_{A,u} \rightarrow G_{A,u^c}^\vee$ . Put  $G_u := G_{A,u} \times_{G_{A,u^c}} (G_{u^c}/G_{u^c}[\varpi_v])^\vee$ , so that  $G_u \rightarrow G_{A,u}$  is also an isogeny of  $O_{F_v}$ -divisible modules whose kernel is contained in  $G_u[\varpi_v]$  and has degree  $q_v^r$ . Put  $G := G_u \times G_{u^c}$ ,  $G_A := G_{A,u} \times G_{A,u^c}$  and similarly  $G_{A_0} := G_{A_0,u} \times G_{A_0,u^c}$ . We also denote the induced isogeny  $G \rightarrow G_A$  by  $\gamma$  and denote by  $\check{\gamma}: G_A \rightarrow G$  the unique isogeny such that  $\check{\gamma} \circ \gamma = [\varpi_v]_G$ . Finally, we introduce the following two subsets:

- Let  $\text{Hom}(G_{A_0}^{\oplus r}, G)^\circ$  be the subset of  $\text{Hom}(G_{A_0}^{\oplus r}, G)$  consisting of elements  $y$  such that the induced map  $y[\varpi_v]: G_{A_0}^{\oplus r}[\varpi_v] \rightarrow G[\varpi_v]/\text{Ker } \gamma$  is an isomorphism.
- Let  $\text{Hom}(G_{A_0}^{\oplus r}, G_A)^\circ$  be the subset of  $\text{Hom}(G_{A_0}^{\oplus r}, G_A)$  consisting of elements  $y$  such that the induced map  $y[\varpi_v]: G_{A_0}^{\oplus r}[\varpi_v] \rightarrow G_A[\varpi_v]/\text{Ker } \check{\gamma}$  is an isomorphism.

Take  $x_1, x_2 \in V^r \otimes_F \mathbb{A}_F^\infty$  satisfying (4.10). For  $i = 1, 2$ , put  $T_i := T(x_i)$  which belongs to  $\text{Herm}_r^\circ(F)^+ \cap \text{Herm}_r(O_{F_v})$ ; and let  $e_i \in \mathbb{N}^{p_v}$  be the unique element such that  $x_{i,v} \in \Lambda_{v,i}^{[e_i]}$ . We define two moduli schemes  $\mathcal{Y}(x_i)$  finite over  $\mathcal{X}$  for  $i = 1, 2$ , such that for every object  $S = (A_0, \lambda_0, \eta_0; A, \lambda, \eta; G_{u^c} \rightarrow G_{A,u^c})$  of  $\mathcal{X}$ , the set  $\mathcal{Y}(x_i)(S)$  consists of elements  $y_i \in \text{Hom}_{O_E}(A_0^{\oplus r}, A)_\mathbb{Q}$  satisfying  $T(y_i) = T_i$ ,  $y_i^v \in \eta(L^v x^v)$ ,<sup>12</sup> and

$$(*) \quad \varpi_v^{e_1 + \varepsilon_v} y_{1,v} \text{ belongs to } \text{Hom}(G_{A_0}^{\oplus r}, G)^\circ \text{ or } \varpi_v^{e_2} y_{2,v} \text{ belongs to } \text{Hom}(G_{A_0}^{\oplus r}, G_A)^\circ, \text{ where } y_{i,v} := y_i[v^\infty].$$

By [LL21, Lemma 5.4],  $\mathcal{Z}(x_i)$  is contained in the image of  $\mathcal{Y}(x_i)$  in  $\mathcal{X}$  for  $i = 1, 2$ . Thus, it remains to show that there are no  $\bar{\kappa}$ -points of  $\mathcal{X}$  underlying both  $\mathcal{Y}(x_1)$  and  $\mathcal{Y}(x_2)$ . We show this by contradiction. Suppose that  $(A_0, \lambda_0, \eta_0; A, \lambda, \eta; G_{u^c} \rightarrow G_{A,u^c})$  is such a  $\bar{\kappa}$ -point, with  $y_1$  and  $y_2$  above it. We study

$$T(y_1, y_2): A_0^{\oplus 2r} \xrightarrow{(y_1, y_2)} A \xrightarrow{\lambda} A^\vee \xrightarrow{(y_1, y_2)^\vee} (A_0^{\oplus 2r})^\vee$$

regarded as an element in  $\text{Herm}_{2r}(F)$ , via its component at  $v$ . We claim that the composite map

$$T_{12}: G_{A_0}^{\oplus r} \xrightarrow{\varpi_v^{e_2} y_{2,v}} G_A \xrightarrow{\lambda_v} G_A^\vee \xrightarrow{(\varpi_v^{e_1 + \varepsilon_v} y_{1,v})^\vee} (G_{A_0}^{\oplus r})^\vee$$

is an isomorphism. Indeed,  $T_{12}[\varpi_v]$  is the composition of three isomorphisms

$$G_{A_0}^{\oplus r}[\varpi_v] \xrightarrow{\varpi_v^{e_2} y_{2,v}} G_A[\varpi_v]/\text{Ker } \check{\gamma}, \quad G_A[\varpi_v]/\text{Ker } \check{\gamma} \xrightarrow{\lambda_v} G_A^\vee[\varpi_v]/\text{Ker } \gamma^\vee, \quad G_A^\vee[\varpi_v]/\text{Ker } \gamma^\vee \xrightarrow{(\varpi_v^{e_1 + \varepsilon_v} y_{1,v})^\vee} (G_{A_0}^{\oplus r})^\vee[\varpi_v]$$

in which the first and third follow from (\*), which implies that  $T_{12}$  is an isomorphism. Since  $\partial_{r,r} T(y_1, y_2) = (T_1, T_2)$ , we conclude that  $T(y_1, y_2) \in \mathfrak{T}_v^{[e_1 + \varepsilon_v + e_2]}$  and hence  $T(y_1, y_2) \in \text{Herm}_{2r}^\circ(F)$ . Thus,  $(y_1, y_2): A_0^{\oplus 2r} \rightarrow A$  is an isogeny, which is impossible by [RSZ20, Lemma 8.7]. Claim (1) is confirmed.

For (2), since  $\mathcal{Z}(x)$  remains the same if we scale  $x$  by an element in  $F^\times$ , we may assume that  $x_v \in (\Lambda_{v,1} \oplus \Lambda_{v,2})^r$  for every  $v \in \mathbb{V}_F^{(p)}$ . Up to a Hecke translation away from  $p$ , which does not affect the conclusion of (2), we may also assume that  $x \in V^r$ . Similar to the argument for (1), we have a moduli scheme  $\mathcal{Y}(x)$  finite over  $\mathcal{X}$ , such that for every object  $S = (A_0, \lambda_0, \eta_0; A, \lambda, \eta; G_{u^c} \rightarrow G_{A,u^c})$  of  $\mathcal{X}$ , the set  $\mathcal{Y}(x)(S)$  consists of elements  $y \in \text{Hom}_{O_E}(A_0^{\oplus r}, A) \otimes \mathbb{Z}_{(p)}$  satisfying  $T(y) = T(x)$  and  $y^p \in \eta(L^p x)$ . Again by [LL21, Lemma 5.4],  $\mathcal{Z}(x)$  is contained in the image of  $\mathcal{Y}(x)$  in  $\mathcal{X}$ . Thus, it suffices to show that the dimension of  $\mathcal{Y}(x)^{(h)}$  is at most  $r - h$  for  $h = 1, 2$ , where  $\mathcal{Y}(x)^{(h)} := \mathcal{Y}(x) \times_{\mathcal{X}} \mathcal{X}^{(h)}$ .

Let  $V_x \subseteq V$  be the hermitian subspace (of dimension  $r$ ) that is the orthogonal complement of the subspace spanned by  $x$ . Put  $H_x := \text{U}(V_x)$  which is naturally a subgroup of  $H$ , and put  $L_x := L \cap H_x(\mathbb{A}_F^\infty)$ . We have a similar moduli scheme  $\mathcal{X}_x$  over  $O_K$  for  $V_x$  similar to the one for  $V$  but with the hyperspecial level structure at  $p$ . More precisely, for every  $S \in \text{Sch}'_{O_K}$ ,  $\mathcal{X}_x(S)$  is the set of equivalence classes (given by  $p$ -principal isogenies) of tuples  $(A_0, \lambda_0, \eta_0; A_1, \lambda_1, \eta_1)$  where

- $(A_0, \lambda_0, \eta_0)$  is like the one in the definition of  $\mathcal{X}$ ,
- $A_1$  is an abelian scheme over  $S$  with an action of  $O_E$  of signature type  $r\Phi - \text{inc} + \text{inc}^c$ , together with a compatible  $p$ -principal polarization  $\lambda_1$ ,
- $\eta_1$  is an  $L_x^p$ -level structure for the hermitian space  $\text{Hom}_{O_E}(A_0, A_1) \otimes_F \mathbb{A}_F^{\infty p}$ .

In particular,  $\mathcal{X}_x$  is a projective smooth scheme over  $O_K$  of pure relative dimension  $r - 1$ . Put  $X_x := \mathcal{X}_x \otimes_{O_K} \kappa$ ; and for  $h \geq 1$ , denote by  $X_x^{[h]}$  the closed locus of  $X_x$  where the height of the connected part of  $G_{A_1, u^c} := A_1[(u^c)^\infty]$

<sup>12</sup>This is irrelevant for our argument below; but see [LL21, Definition 5.3] for more details.

is at least  $h$ . It is known that  $X_x^{[h]}$  has pure dimension  $r - h$ . Claim (2) will follow if there is a finite morphism  $f: \mathcal{Y}(x) \rightarrow X_x$  that sends  $Y(x)^{(h)}$  into  $X_x^{[h]}$  for  $h = 1, 2$ , which we now construct.

Take a point  $P = (A_0, \lambda_0, \eta_0; A, \lambda, \eta; G_{u^c} \rightarrow G_{A, u^c}; y)$  of  $\mathcal{Y}(x)(S)$ . Put  $A' := (A^\vee / (\lambda \circ y) A_0^{\oplus r})^\vee$ , which inherits an action of  $O_E$  which has signature type  $r\Phi - \text{inc} + \text{inc}^c$  and admits a natural map to  $A$ . Since  $T(x) \in \text{Herm}_r^\circ(F)$ , the induced map  $\lambda': A' \rightarrow A \xrightarrow{\lambda} A^\vee \rightarrow A'^\vee$  is a quasi-polarization such that  $\lambda'[p^\infty]$  is an isogeny. For every  $\tilde{u} \in \mathbf{P}$ , we have the induced isogeny  $\lambda'_{\tilde{u}^c}: G_{A', \tilde{u}^c} \rightarrow G_{A, \tilde{u}}$ . Put

$$A_1 := A' \left/ \bigoplus_{\tilde{u} \in \mathbf{P}_\Phi} \text{Ker } \lambda'_{\tilde{u}^c} \right.$$

and let  $\lambda_1: A_1 \rightarrow A_1^\vee$  be the quasi-polarization induced from  $\lambda'$ , which is in fact  $p$ -principal from the construction. We can also define a natural  $L_x^p$ -level structure  $\eta_1$  for  $A_1$  whose details we leave to the reader. Then we define  $f(P)$  to be  $(A_0, \lambda_0, \eta_0; A_1, \lambda_1, \eta_1)$ . Since the  $O_{F_v}$ -divisible module  $G_{A_0, u^c}$  is étale, the height of the connected part of  $G_{A_1, u^c}$  equals to that of  $G_{A_0, u^c}$ . In particular,  $f$  sends  $Y(x)^{(h)}$  into  $X_x^{[h]}$  for  $h = 1, 2$ . It remains to show that  $f$  is finite. Since  $\mathcal{Y}(x)$  is proper over  $O_K$ , it suffices to show that the fiber of  $f$  over an arbitrary  $\bar{k}$ -point is finite. Indeed, when  $S = \text{Spec } \bar{k}$ ,  $G_{A', \tilde{u}^c}$  has dimension 1 (resp. is étale) if  $\tilde{u} = u$  (resp.  $\tilde{u} \in \mathbf{P}_\Phi \setminus \{u\}$ ), and the degree of  $\lambda'_{\tilde{u}^c}$  is bounded by the moment matrix  $T(x)$ . It follows that up to isomorphism, there are only finitely many such isogenies  $G_{A', \tilde{u}^c} \rightarrow G_{A, \tilde{u}}$  with fixed  $G_{A, \tilde{u}}$  for every  $\tilde{u} \in \mathbf{P}_\Phi$ . Thus,  $f$  is finite and claim (2) is confirmed.

The lemma is finally proved.  $\square$

**Proposition 4.26.** *Suppose that  $n < p$ . There exist an integer  $M \geq 0$  and a pair  $(t_1, t_2) \in \mathbb{S}_{O_L}^\diamond \times \mathbb{S}_{O_L}^\diamond$  satisfying  $\chi_{\bar{\pi}}^\diamond(t_1)\chi_{\bar{\pi}}^\diamond(t_2) \neq 0$ , such that for every  $(T_1, T_2) \in \text{Herm}_r^\circ(F)^+ \times \text{Herm}_r^\circ(F)^+$ , every  $(s_1, s_2) \in (\mathbb{S}_{O_L}^\diamond)_{\bar{\pi}, L}^2 \times (\mathbb{S}_{O_L}^\diamond)_{\bar{\pi}, L}^2$  and every  $(e_1, e_2) \in \mathbb{N}^p \times \mathbb{N}^p$ , we have*

$$\langle Z_{T_1}(t_1 s_1 \phi_1^{[e_1]})_L, Z_{T_2}(t_2 s_2 \phi_2^{[e_2]})_L \rangle_{E_u} \in (O_{E_u}^\times)^{\text{fr}} \otimes_{\mathbb{Z}_p} (p^{|e_2| - M} \overline{\mathbb{Z}}_p).$$

The rest of this subsection is devoted to the proof of this proposition.

Let  $S$  be the kernel of the norm map  $\text{Nm}_{E/F}: \text{Res}_{O_E/O_F} \mathbf{G} \rightarrow \mathbf{G}_{O_F}$ . Consider the reciprocity map

$$(4.12) \quad \text{rec}: \text{Aut}(\mathbb{C}/E) \rightarrow E^\times \backslash \mathbb{A}_E^{\infty, \times} \rightarrow S(F) \backslash S(\mathbb{A}_F^\infty)$$

in which the first one is from the global class field theory and the second (surjective) one sends  $a$  to  $a^c/a$ . For  $d \in \mathbb{N}$ , we

- put  $L_{S, v}^{[d]} := S(O_{F_v}) \cap (1 + \mathfrak{p}_v^d O_{E_v})$ ,
- let  $E^{[d]} \subseteq \mathbb{C}$  be the finite abelian extension of  $E$  such that the map  $\text{rec}$  (4.12) induces an isomorphism

$$\text{Gal}(E^{[d]}/E) \simeq S(F) \backslash S(\mathbb{A}_F^\infty) \left/ L_{S, v}^{[d]} \prod_{\tilde{v} \in \mathbb{V}_F^{\text{fin}} \setminus \{v\}} S(O_{F_{\tilde{v}}}) \right.,$$

- denote by  $Z_{[d]}^r(X_L)$  the image of the norm map

$$\text{Nm}_{E^{[d]}/E}: Z^r(X_L \otimes_E E^{[d]}) \rightarrow Z^r(X_L).$$

**Lemma 4.27.** *For every  $x \in V^r \otimes_F \mathbb{A}_F^\infty$  satisfying  $T(x) \in \text{Herm}_r^\circ(F)^+$  and  $x_v \in \Lambda_{v, 2}^{[e]}$  for some  $e \in \mathbb{N}^p$ , we have*

$$Z(x)_L \in Z_{[|e|]}^r(X_L).$$

*Proof.* Up to a Hecke translation away from  $p$ , which does not affect the conclusion of the lemma, we may assume that  $x \in V^r$ . Let  $V_x \subseteq V$  be the hermitian subspace (of dimension  $r$ ) that is the orthogonal complement of the subspace spanned by  $x$ . Put  $H_x := \text{U}(V_x)$  which is naturally a subgroup of  $H$ , and put  $L_x := L \cap H_x(\mathbb{A}_F^\infty)$ . We have the Shimura variety  $X_{x, L_x}$  for  $H_x$  with level  $L_x$ , similar to  $X_L$ . By definition,  $Z(x)_L$  is the fundamental cycle of the finite unramified morphism  $X_{x, L_x} \rightarrow X_L$  defined over  $E$ .

We have the determinant map  $\det: H_x \rightarrow S \otimes_{O_F} F$  which identifies  $S \otimes_{O_F} F$  with the maximal abelian quotient of  $H_x$ . Then the set of connected components of  $X_{x, L_x} \otimes_E \mathbb{C}$  is canonically parameterized by the set  $S(F) \backslash S(\mathbb{A}_F^\infty) / \det L_x$ . For every  $s \in S(F) \backslash S(\mathbb{A}_F^\infty) / \det L_x$ , we denote by  $X_{x, L_x}^s$  the corresponding connected component. The definition of canonical models of Shimura varieties implies that  $\gamma X_{x, L_x}^s = X_{x, L_x}^{\text{rec}(\gamma)s}$  for every  $\gamma \in \text{Aut}(\mathbb{C}/E)$ , where  $\text{rec}$  is the map (4.12).

We claim that  $\det L_{x,v} \subseteq L_{S,v}^{[e]}$ . Then we have the quotient map

$$S(F) \backslash S(\mathbb{A}_F^\infty) / \det L_x \rightarrow S(F) \backslash S(\mathbb{A}_F^\infty) \Big/ L_{S,v}^{[e]} \prod_{\tilde{v} \in \mathbb{V}_F^{\text{fin}} \setminus \{v\}} S(O_{F_{\tilde{v}}})$$

Let  $\mathfrak{S}$  be the fiber of 1 in the above map. Then  $\sum_{s \in \mathfrak{S}} X_{x,L_x}^s$  is defined over  $E^{[e]}$ ; and  $\text{Nm}_{E^{[e]}/E} \sum_{s \in \mathfrak{S}} X_{x,L_x}^s = X_{x,L_x}$ . The lemma then follows.

It remains to show the claim, which is an exercise in linear algebra. We assume  $e \neq 0$  as the case for  $e = 0$  is trivial. By definition,  $L_{x,v}$  is simply the subgroup of  $L_v$  that fixes  $x_v$ , or equivalently,  $x'_v := \varpi_v^e \cdot x_v$ . By the definition of  $\Lambda_{v,2}^{[e]}$  in §4.3(S3),  $x'_v$  belongs to  $(\Lambda_{v,1} \oplus \Lambda_{v,2})^r$  such that  $T(x'_v) \in \varpi_v^{[e]} \text{Herm}_r(O_{F_v})$  and that  $x'_v \bmod \Lambda_{v,1}$  generates  $\Lambda_{v,2}$ . It follows that the image of  $x'_v$  in  $(\Lambda_{v,1} \oplus \Lambda_{v,2})^r \otimes_{O_{F_v}} O_{F_v}/\mathfrak{p}_v^{[e]}$  generates a Lagrangian  $O_{E_v} \otimes_{O_{F_v}} O_{F_v}/\mathfrak{p}_v^{[e]}$ -submodule of  $(\Lambda_{v,1} \oplus \Lambda_{v,2}) \otimes_{O_{F_v}} O_{F_v}/\mathfrak{p}_v^{[e]}$ . In particular, every element in  $L_{x,v}$ , which stabilizes  $\Lambda_{v,1} \oplus \Lambda_{v,2}$ , has determinant 1 modulo  $\mathfrak{p}_v^{[e]}$ . The claim follows.  $\square$

For  $d \in \mathbb{N}$ , let  $u_d$  be the place of  $E^{[d]}$  induced from the fixed isomorphism  $\overline{E}_u \xrightarrow{\sim} \mathbb{C}$ , which is above  $u$ . Put  $K := E_u$ ,  $K_d := E_{u_d}^{[d]}$  for  $d \in \mathbb{N}$  and  $K_\infty := \bigcup_{d \geq 0} K_d$ . Then  $K_0/K$  is unramified and  $K_d/K_0$  is totally ramified of degree  $(q_v - 1)q_v^{d-1}/|U_E|$  for  $d > 0$ , where  $U_E$  is the torsion subgroup of  $O_E^\times$ .

Recall that we have the subspaces  $V_{\pi,L}, V_{\hat{\pi},L} \subseteq H_{\text{ét}}^{2r-1}(X_L \otimes_E \overline{E}, \mathbb{L}(r))$ . Put

$$T_{\pi,L} := V_{\pi,L} \bigcap H_{\text{ét}}^{2r-1}(X_L \otimes_E \overline{E}, O_{\mathbb{L}}(r))^{\text{fr}}, \quad T_{\hat{\pi},L} := V_{\hat{\pi},L} \bigcap H_{\text{ét}}^{2r-1}(X_L \otimes_E \overline{E}, O_{\mathbb{L}}(r))^{\text{fr}},$$

both being  $O_{\mathbb{L}}[\text{Gal}(\overline{E}/E)]$ -modules. For  $d \in \mathbb{N}$ , we put

$$N_\infty H_f^1(K_d, T_{\hat{\pi},L}) := \bigcap_{d' > d} \text{Im} \left( \text{Cor}_{K_{d'}/K_d} : H_f^1(K_{d'}, T_{\hat{\pi},L}) \rightarrow H_f^1(K_d, T_{\hat{\pi},L}) \right),$$

in which  $\text{Cor}_{K_{d'}/K_d}$  denotes the corresponding corestriction map.

**Lemma 4.28.** *There exists an integer  $M \geq 0$  such that  $p^M$  annihilates  $H_f^1(K_d, T_{\hat{\pi},L})/N_\infty H_f^1(K_d, T_{\hat{\pi},L})$  for every  $d \in \mathbb{N}$ .*

*Proof.* By Lemma 4.15 and Remark 4.16 (for  $\hat{\pi}$ ), we know that  $V_{\hat{\pi},L}|_{K_d}$  satisfies the Panchishkin condition in [Nek93, 6.7] for every  $d \in \mathbb{N}$ . By Lemma 4.14, we may apply [Nek93, Theorem 6.9] to  $V_{\pi,L}|_{K_d}$ .<sup>13</sup> Thus, by the same argument at the end of the proof of [Nek95, Proposition II.5.10], it suffices to show that  $H^0(K_\infty, V_{\pi,L}) = H^0(K_\infty, V_{\hat{\pi},L}) = 0$ .

We follow the strategy in [Shn16, Section 8]. We may choose an element  $\xi \in S(F)$  such that  $\xi = (\varpi_v^{1_u - 1_{u^c}})^{[K_0:K]}$  in  $\text{Gal}(E^{[0]}/E)$ . Then by the same argument for [Shn16, Proposition 8.3],  $K_\infty$  is contained in  $K_\xi$  – the field attached to the Lubin–Tate group relative to the extension  $K_0/K$  with parameter  $\xi$ . Let  $\chi_\xi : \text{Gal}(K_\xi/K_0) \rightarrow O_K^\times$  be the character given by the Galois action on the torsion points of this relative Lubin–Tate group; and let  $K(\chi_\xi)$  be the corresponding 1-dimensional representation. Let  $L$  be the maximal subfield of  $K$  that is unramified over  $\mathbb{Q}_p$ . By the same argument for [Shn16, Proposition 8.4],  $K(\chi_\xi)$  is crystalline, and that the  $q_v$ -Frobenius map (which is  $L$ -linear) acts on  $\mathbb{D}_{\text{cris}}(K(\chi_\xi))$ , which is a free  $K \otimes_{\mathbb{Q}_p} L$ -module of rank 1, by multiplication by  $\xi^{-1}$ .<sup>14</sup> Note that  $\mathbb{L}$  is a subfield of  $\mathbb{C}$  and hence  $\overline{K}$  via the fixed isomorphism  $\overline{K} = \overline{E}_u \xrightarrow{\sim} \mathbb{C}$ . Let  $V$  be either  $V_{\pi,L} \otimes_{\mathbb{L}} \overline{K}$  or  $V_{\hat{\pi},L} \otimes_{\mathbb{L}} \overline{K}$ . Repeating the argument in [Shn16, Proposition 8.9] (which followed an approach in [Nek95]) to  $V$ , we obtain  $H^0(K_\xi, V) = 0$  since  $V$  is crystalline of pure weight  $-1$  by Lemma 4.15.

The lemma is proved.  $\square$

*Proof of Proposition 4.26.* Let  $M \geq 0$  be the integer in Lemma 4.28 and  $(t_1, t_2) \in \mathbb{S}_{O_{\mathbb{L}}}^\diamond \times \mathbb{S}_{O_{\mathbb{L}}}^\diamond$  the pair in Lemma 4.25. We first note that  $Z_{T_i}(t_i s_i \phi_i^{[e_i]})_L \in Z^r(X_L) \otimes \overline{\mathbb{Z}}_p$  for  $i = 1, 2$ . By Lemma 4.27, we may find an element

<sup>13</sup>Though our extension  $K_\infty/K_d$  is in general not a  $\mathbb{Z}_p$ -extension as assumed in [Nek93, 6.2], the argument for [Nek93, Theorem 6.9] works without change.

<sup>14</sup>Note that in this article, we always use the covariant version for  $\mathbb{D}_{\text{dR}}$  and  $\mathbb{D}_{\text{cris}}$ .

$Z \in Z^r(X_L \otimes_E E^{[e_2]}) \otimes \overline{\mathbb{Z}}_p$  such that  $\text{Nm}_{E^{[e_2]}/E} Z = Z_{T_2}(t_2 \phi_2^{[e_2]})_L$ . We may also assume that the support of  $Z$  is contained in the support of  $Z_{T_2}(t_2 \phi_2^{[e_2]})_L$ . Put

$$Z_2 := \text{Nm}_{E^{[e_2]}/E} Z \otimes_E K \in Z^r(X_L \otimes_E K_{[e_2]}) \otimes \overline{\mathbb{Z}}_p,$$

so that  $\text{Nm}_{K_{[e_2]}/K} Z_2 = Z_{T_2}(t_2 \phi_2^{[e_2]})_L \otimes_E K$ . Since the natural map  $T_L/T_{\hat{\pi},L} \rightarrow H_{\text{ét}}^{2r-1}(X_L \otimes_E \overline{E}, \mathbb{L}(r))/V_{\hat{\pi},L}$  is injective, the cycle class of  $Z_{T_1}(t_1 s_1 \phi_1^{[e_1]})_L = s_1^* Z_{T_1}(s_1 \phi_1^{[e_1]})_L$  in  $H^{2r}(X_L \otimes_E K, \overline{\mathbb{Z}}_p(r))$  sits in  $H_f^1(K, T_{\hat{\pi},L})$ . Similarly, the cycle class of  $s_2^* Z_2$  in  $H^{2r}(X_L \otimes_E K_{[e_2]}, \overline{\mathbb{Z}}_p(r))$  sits in  $H_f^1(K_{[e_2]}, T_{\pi,L})$ . By [Nek95, II.(1.9.1)], we have

$$\langle Z_{T_1}(t_1 s_1 \phi_1^{[e_1]})_L, Z_{T_2}(t_2 s_2 \phi_2^{[e_2]})_L \rangle_K = \text{Nm}_{K_{[e_2]}/K} \langle s_1^* Z_{T_1}(t_1 \phi_1^{[e_1]})_L \otimes_E K_{[e_2]}, s_2^* Z_2 \rangle_{K_{[e_2]}}.$$

By Lemma 4.25, we have

$$\langle s_1^* Z_{T_1}(t_1 \phi_1^{[e_1]})_L \otimes_E K_{[e_2]}, s_2^* Z_2 \rangle_{K_{[e_2]}} \in (O_{K_{[e_2]}}^\times)^{\text{fr}} \otimes_{\mathbb{Z}_p} \mathbb{C}.$$

In other words, the corresponding bi-extension is crystalline (Remark A.11). By the argument for [Nek95, Proposition II.1.11], we have

$$\langle s_1^* Z_{T_1}(t_1 \phi_1^{[e_1]})_L \otimes_E K_{[e_2]}, s_2^* Z_2 \rangle_{K_{[e_2]}} \in (O_{K_{[e_2]}}^\times)^{\text{fr}} \otimes_{\mathbb{Z}_p} (p^{-M} \overline{\mathbb{Z}}_p).$$

Finally, since the image of the norm map  $\text{Nm}_{K_d/K} : (O_{K_d}^\times)^{\text{fr}} \rightarrow (O_K^\times)^{\text{fr}}$  is precisely  $p^d (O_K^\times)^{\text{fr}}$  for  $d \in \mathbb{N}$ , we have

$$\langle Z_{T_1}(t_1 s_1 \phi_1^{[e_1]})_L, Z_{T_2}(t_2 s_2 \phi_2^{[e_2]})_L \rangle_K \in (O_K^\times)^{\text{fr}} \otimes_{\mathbb{Z}_p} (p^{|e_2| - M} \overline{\mathbb{Z}}_p).$$

The proposition is proved.  $\square$

**4.6. Proof of main theorems.** To shorten notation, in this subsection we put

$$\mathbf{R} := \mathbf{V}_F^{(\diamond \setminus \{p\})} \cap \mathbf{V}_F^{\text{spl}} \cap \mathbf{V}_F^\heartsuit, \quad \mathbf{T} := \mathbf{V}_F^{\text{fin}} \setminus (\mathbf{V}_F^{(\diamond)} \cap \mathbf{V}_F^{\text{spl}} \cap \mathbf{V}_F^\heartsuit),$$

so that  $\mathbf{V}_F^{(p)} \cup \mathbf{R} \cup \mathbf{T}$  is a partition of  $\mathbf{V}_F^{\text{fin}}$ .

For every  $v \in \mathbf{R}$ , we

- (R1) choose a  $(\varphi_{1,v}^\dagger, \varphi_{2,v})$ -typical element  $\mathbf{f}_v \in \mathcal{S}(\text{Herm}_{2r}(F_v))$  (Definition 3.26) which exists by §4.3(T1) and Remark 3.27,
- (R2) choose an integer  $N_v > 0$  such that  $N_v \widehat{\mathbf{f}}_v$  takes values in  $\mathbb{Z}$  and  $N_v \mathbf{f}_v^{1_v} = f_{\Phi_v}^{\text{SW}}$  for some  $\Phi_v \in \mathcal{S}(V_v^{2r}, \mathbb{Z}\langle p_v \rangle)$ ,
- (R3) put  $f_{\chi_v} := N_v \mathbf{f}_v^{\chi_v} \in I_{r,v}^\square(\chi_v)$  (3.2) for every finite character  $\chi : \Gamma_{F,p} \rightarrow \mathbb{C}^\times$ ,
- (R4) write  $\Phi_v = \sum_{k_i \in \Xi_v} \phi_{v,1}^{(k_i)} \otimes \phi_{v,2}^{(k_i)}$  in which  $\Xi_v$  is a finite set and  $\phi_{v,i}^{(k_i)} \in \mathcal{S}(V_v^r, \mathbb{Z}\langle p_v \rangle)$ ,
- (R5) choose  $L_v \subseteq H(F_v)$  and  $K_v \subseteq G_r(F_v)$  so that they fix every  $\phi_{v,i}^{(k_i)}$ .

Put  $N := \prod_{v \in \mathbf{R}} N_v$  and  $\Xi := \prod_{v \in \mathbf{R}} \Xi_v$ .

For every  $v \in \mathbf{T}$ , we

- put  $\Phi_v := \phi_{v,1} \otimes \phi_{v,2} = \mathbf{1}_{\Lambda_v^{2r}}$  (§4.3(S2)),
- let  $f_{\chi_v} \in I_{r,v}^\square(\chi_v)$  be the section from Lemma 4.22 for every finite character  $\chi : \Gamma_{F,p} \rightarrow \mathbb{C}^\times$ .

For every  $v \in \mathbf{V}_F^{(p)}$  and  $e \in \mathbb{N}$  regarded as a constant tuple in  $\mathbb{N}^{\mathbb{P}_v}$ , we

- put  $\Phi_v^{[e]} := \phi_{v,1}^{[e]} \otimes \phi_{v,2}^{[e]}$  (§4.3(S3)),
- put  $f_{\chi_v}^{[e]} := b_{2r,v}(\mathbf{1})^{-1} \cdot \text{vol}(L_v, dh_v) \cdot (\mathbf{f}_{\chi_v}^{[2e+\varepsilon^c]})^{\chi_v} \in I_{r,v}^\square(\chi_v)$  for every finite character  $\chi : \Gamma_{F,p} \rightarrow \mathbb{C}^\times$ , so that  $\mathbf{f}_{\mathbf{1}_v} = f_{\Phi_v^{[e]}}^{\text{SW}}$  by Lemma 4.20(3).

For every  $e \in \mathbb{N}$ , we

- put

$$\phi_i^{[e](k)} := \left( \bigotimes_{v \in \mathbf{V}_F^{(p)}} \phi_{v,i}^{[e]} \right) \otimes \left( \bigotimes_{v \in \mathbf{R}} \phi_{v,i}^{(k_v)} \right) \otimes \left( \bigotimes_{v \in \mathbf{T}} \phi_{v,i} \right) \in \mathcal{S}(V^r \otimes_F \mathbb{A}_F^\infty, \mathbb{Z}\langle \diamond \rangle)^{K_r^\diamond \times L}$$

for  $i = 1, 2$  and  $k = (k_v)_v \in \Xi$ ,

- put

$$\Phi^{[e]} := \left( \bigotimes_{v \in \mathbb{V}_F^{(p)}} \Phi_v^{[e]} \right) \otimes \left( \bigotimes_{v \in \mathbb{V}_F^{\text{fin}} \setminus \mathbb{V}_F^{(p)}} \Phi_v \right) \in \mathcal{S}(V^r \otimes_F \mathbb{A}_F^\infty, \mathbb{Z}\langle \diamond \rangle)^{K_{2r}^\diamond \times L},$$

- put

$$f_{\chi^\infty}^{[e]} := \left( \bigotimes_{v \in \mathbb{V}_F^{(p)}} f_{\chi_v}^{[e]} \right) \otimes \left( \bigotimes_{v \in \mathbb{V}_F^{\text{fin}} \setminus \mathbb{V}_F^{(p)}} f_{\chi_v} \right) \in \mathbb{I}_r^\square(\chi)^\infty$$

for every finite character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times$ .

In each of the above notation, we suppress the superscript  $[e]$  when  $e = 0$ .

Finally, we set

- $M_1 := \max\{M_u \mid u \in \mathbb{P}\}$  in which  $M_u \geq 0$  is the smallest element for which Proposition 4.26 (for  $u$ ) holds,
- $M_2 \in \mathbb{N}$  to be the smallest element such that both  $p^{M_2} \text{vol}^\natural(L)$  and  $p^{M_2} W_{2r}$  belong to  $\mathbb{Z}_{(p)}$ ,
- $M_3 \in \mathbb{N}$  to be the smallest element such that  $p^{M_3} \prod_{v \in \mathbb{V}_F^{(p)}} \text{vol}(L_v, dh_v) \in \mathbb{Z}_{(p)}$ ,
- $(t_1, t_2) \in \mathbb{S}_{O_L}^\diamond \times \mathbb{S}_{O_L}^\diamond$  to be the product of those chosen pairs from Proposition 4.24 and Proposition 4.26 for every  $u \in \mathbb{P}$ , which satisfies  $\chi_{\hat{\pi}}^\diamond(t_1) \chi_{\hat{\pi}}^\diamond(t_2) \neq 0$ .

These will not be changed in the rest of this subsection. Put  $M = M_1 + M_2 + M_3$ .

*Remark 4.29.* For every  $v \in \mathbb{V}_F^{\text{spl}} \setminus \mathbb{V}_F^{(\diamond)}$ , we have a canonical isomorphism

$$\mathbb{Z}[L_v \setminus H(F_v)/L_v] \simeq \mathbb{Z}[K_{r,v} \setminus G_r(F_v)/K_{r,v}]$$

of rings via Satake isomorphisms. By [Liu11a, Proposition A.5], we know that the action of  $s \in \mathbb{Z}[L_v \setminus H(F_v)/L_v]$  on  $\mathcal{S}(V_v^r)^{K_{r,v} \times L_v}$  via the Weil representation  $\omega_{r,v}$  coincides with that of  $\hat{s} \in \mathbb{Z}[K_{r,v} \setminus G_r(F_v)/K_{r,v}]$ , where  $\hat{s}$  denotes the adjoint of  $s$ .

Note that for  $v \in \mathbb{T} \cap \mathbb{V}_F^{(\diamond)}$ , the section  $f_{\chi_v}$  used here is completely different from the one used in the construction of the  $p$ -adic  $L$ -function in §3.5. We need to study the relation of the doubling integral using these new sections with the  $p$ -adic  $L$ -function we constructed before.

For every  $(s_1, s_2) \in \mathbb{S}_{O_L}^\diamond \times \mathbb{S}_{O_L}^\diamond$ , every  $e \in \mathbb{N}$  and every finite character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times = \overline{\mathbb{Q}}_p^\times$ , we define an Eisenstein series

$$E_{[e]}^{(s_1, s_2)}(\chi) := p^M \cdot b_{2r}^\diamond(\mathbf{1})^{-1} \cdot b_{2r}^\diamond(\chi) \cdot E(-, f_\infty^{[r]} \otimes (\hat{s}_1 \hat{t}_1, \hat{s}_2 \hat{t}_2) f_{\chi^\infty}^{[e]}),$$

which is an element of  $\mathcal{A}_{2r, \text{hol}}^{[r]}$ . Recall that we have the map  $\rho_{r,r}: \mathcal{A}_{2r, \text{hol}}^{[r]} \rightarrow \mathcal{A}_{r, \text{hol}}^{[r]}$  (2.2) and have chosen vectors  $\varphi_1 = \otimes_v \varphi_{1,v} \in \mathcal{V}_{\hat{\pi}}$  and  $\varphi_2 = \otimes_v \varphi_{2,v} \in \mathcal{V}_\pi$  from §4.3(T1–T4). In what follows, we will frequently use Notation 3.35 and its linear extension to  $\overline{\mathbb{Q}}_p \simeq \mathbb{C}$ .

**Lemma 4.30.** *For every  $(s_1, s_2) \in \mathbb{S}_{O_L}^\diamond \times \mathbb{S}_{O_L}^\diamond$ , every  $e \in \mathbb{N}$  and every finite character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times = \overline{\mathbb{Q}}_p^\times$ , we have*

$$\begin{aligned} \left\langle \varphi_1 \otimes \varphi_2, \rho_{r,r} E_{[e]}^{(s_1, s_2)}(\chi) \right\rangle_{\pi, \hat{\pi}} &= N \cdot p^M \cdot \chi_{\hat{\pi}}^\diamond(t_1 s_1) \chi_\pi^\diamond(t_2 s_2) \cdot \left( \prod_{u \in \mathbb{P}} \alpha(\pi_u) \right)^{2e} \cdot \mathcal{L}_p^\diamond(\pi)(\chi) \\ &\quad \times \prod_{v \in \mathbb{V}_F^{(p)}} \prod_{u \in \mathbb{P}_v} \gamma\left(\frac{1+r}{2}, \underline{\pi}_u, \psi_{F,v}\right) \cdot \prod_{v \in \mathbb{V}_F^{(p)}} Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{1,v}) \cdot \prod_{v \in \mathbb{T} \cap \mathbb{V}_F^{(\diamond)}} Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\chi_v}), \end{aligned}$$

where we recall that  $N = \prod_{v \in \mathbb{R}} N_v$  (R2) and the local doubling zeta integral is (3.10) (with  $\iota$  being the restriction of the fixed isomorphism  $\overline{\mathbb{Q}}_p \xrightarrow{\sim} \mathbb{C}$ ).

*Proof.* By the doubling integral expansion, we have

$$\begin{aligned} \left\langle \varphi_1 \otimes \varphi_2, \rho_{r,r} E_{[e]}^{(s_1, s_2)}(\chi) \right\rangle_{\pi, \hat{\pi}} &= p^M \cdot \chi_{\hat{\pi}}^\diamond(t_1 s_1) \chi_\pi^\diamond(t_2 s_2) \cdot \frac{1}{P_\pi} \cdot \frac{Z_r^{[F:\mathbb{Q}]}}{b_{2r}^\diamond(\mathbf{1})} \cdot L\left(\frac{1}{2}, \text{BC}(\pi^\diamond) \otimes (\chi^\diamond \circ \text{Nm}_{E/F})\right) \\ &\quad \times \prod_{v \in \mathbb{V}_F^{(p)}} Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\chi_v}^{[e]}) \cdot \prod_{v \in \mathbb{V}_F^{(\diamond) \setminus \{\infty, p\}}} Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\chi_v}). \end{aligned}$$



By Lemma 4.20(3) and Lemma 3.10, we have for  $v \in \mathbb{V}_F^{(p)}$

$$Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\chi_v}^{[e]}) = \left( \prod_{u \in \mathbb{P}_v} \alpha(\pi_u) \right)^{2e} \cdot Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\chi_v});$$

and together with Proposition 3.31, we have

$$\frac{Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\chi_v})}{Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\mathbf{1}_v})} = \prod_{u \in \mathbb{P}_v} \frac{\gamma(\frac{1+r}{2}, \pi_u, \psi_{F,v})}{\gamma(\frac{1+r}{2}, \pi_u \otimes \chi_v, \psi_{F,v})}.$$

For  $v \in \mathbb{R}$ , we have

$$Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\chi_v}) = N_v \cdot Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, \mathbf{f}_v^{\chi_v}) = N_v.$$

Together with Theorem 3.37, we obtain the lemma.  $\square$

The following nonvanishing result is needed to make Lemma 4.30 useful.

**Lemma 4.31.** *For every  $v \in \mathbb{T}$ , we have*

$$Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\mathbf{1}_v}) \neq 0.$$

*Proof.* By [Liu22, Proposition 5.6], [LL22, Proposition 3.6] and Lemma 3.30 when  $v \in \mathbb{S}_\pi$ , when  $v \in \mathbb{V}_F^{\text{ram}}$  and when  $v \in \mathbb{T} \setminus (\mathbb{S}_\pi \cup \mathbb{V}_F^{\text{ram}})$ , respectively, we have

$$Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\mathbf{1}_v}) = C_v \cdot \frac{L(\frac{1}{2}, \mathbf{BC}(\pi_v))}{b_{2r,v}(\mathbf{1})}$$

for a constant  $C_v \in \mathbb{Q}^\times$ . Then the nonvanishing is clear.  $\square$

In the next two lemmas we study Siegel–Fourier coefficients of  $E_{[e]}^{(s_1, s_2)}(\chi)$  with  $\chi$  varying in terms of  $p$ -adic measures. The readers are recommended to read §4.7 first at this point.

**Lemma 4.32.** *Take  $(s_1, s_2) \in \mathbb{S}_{O_L}^\diamond \times \mathbb{S}_{O_L}^\diamond$ ,  $e \in \mathbb{N}$ ,  $(g_1, g_2) \in P_r(F_{\mathbb{R}}) \times P_r(F_{\mathbb{R}})$  and  $T^\square \in \text{Herm}_{2r}^\circ(F)^+$ .*

(1) *There exists a unique  $\overline{\mathbb{Q}}_p$ -valued  $p$ -adic measure (Definition 4.37)  $\mathfrak{d}_{(g_1, g_2)}^{(s_1, s_2)} \mathscr{W}_{T^\square}^{[e]}$  on  $\Gamma_{F,p}$  such that*

$$(4.13) \quad \mathfrak{d}_{(g_1, g_2)}^{(s_1, s_2)} \mathscr{W}_{T^\square}^{[e]}(\chi) = p^M W_{2r}^\diamond \cdot b_{2r}^\diamond(\chi) \cdot W_{T^\square}((g_1, g_2)(\hat{s}_1 \hat{t}_1, \hat{s}_2 \hat{t}_2) f_{\chi^\diamond}^{[e]})$$

*for every finite character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times = \overline{\mathbb{Q}}_p^\times$ , where  $W_{2r}^\diamond$  is from (3.5).*

(2) *The  $p$ -adic measure  $\mathfrak{d}_{(g_1, g_2)}^{(s_1, s_2)} \mathscr{W}_{T^\square}^{[e]}$  is integral (Definition 4.37).*

(3) *We have  $\mathfrak{d}_{(g_1, g_2)}^{(s_1, s_2)} \mathscr{W}_{T^\square}^{[e]}(\mathbf{1}) = 0$ ; and  $\partial_{(g_1, g_2)}^{(s_1, s_2)} \mathscr{W}_{T^\square}^{[e]}(\mathbf{1}) = 0$  unless  $T^\square \in \text{Herm}_{2r}^\circ(F)_V^+$ .*

*Proof.* For (1), the right-hand side of (4.13) can be rewritten as a product

$$(4.14) \quad \left( p^M W_{2r}^\diamond \prod_{v \in \mathbb{V}_F^{(p)}} b_{2r,v}(\mathbf{1})^{-1} \text{vol}(L_v, dh_v) \right) \times \left( \chi_p(\text{Nm}_{E/F} \det T_{12}^\square) \mathbf{1}_{\mathbb{Z}_p^{[2e+e\epsilon_1]}(T^\square)} \right) \\ \times \left( \prod_{v \in \mathbb{V}_F^{\diamond(\infty, p)}} W_{T^\square}((g_{1,v}, g_{2,v}) f_{\chi_v}) \right) \times \left( b_{2r}^\diamond(\chi) \cdot W_{T^\square}((\hat{s}_1 \hat{t}_1, \hat{s}_2 \hat{t}_2) f_{\chi^\diamond}) \right)$$

of four factors. We show that the four factors are all given by  $\overline{\mathbb{Q}}_p$ -valued  $p$ -adic measures. The first one is a constant. The second one is obviously given by such a measure. By Lemma 3.2(1,2) and Lemma 4.22, the third one is given by such a measure. By Lemma 3.5(1), the last one is given by such a measure as well.

For (2), since

$$p^M W_{2r}^\diamond \prod_{v \in \mathbb{V}_F^{(p)}} b_{2r,v}(\mathbf{1})^{-1} \text{vol}(L_v, dh_v) = p^{M_1} \cdot p^{M_2} W_{2r} \cdot b_{2r, \diamond \setminus \{\infty, p\}}(\mathbf{1}) \cdot p^{M_3} \prod_{v \in \mathbb{V}_F^{(p)}} \text{vol}(L_v, dh_v),$$

the first factor in (4.14) belongs to  $\overline{\mathbb{Z}}_p$  by the definitions of  $M_2$  and  $M_3$ . By Lemma 3.2(1,2), Lemma 4.22 and Lemma 3.5(1), the product of the remaining three factors in (4.14) is given by an integral  $p$ -adic measure. Thus,  $d_{(g_1, g_2)}^{(s_1, s_2)} \mathscr{W}_{T^\square}^{[e]}$  is integral.

For (3), we only need to realize that  $W_{T^\square}(f_{1_v}) = 0$  for every  $v \in \text{Diff}(T^\square, V)$ .

The lemma is proved.  $\square$

**Lemma 4.33.** *For every  $(s_1, s_2) \in \mathbb{S}_{O_L}^\diamond \times \mathbb{S}_{O_L}^\diamond$ , every  $e \in \mathbb{N}$ , every  $(g_1, g_2) \in P_r(F_R) \times P_r(F_R)$  and every finite character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times = \overline{\mathbb{Q}}_p^\times$ , we have*

$$q_{2r}^{\text{an}} \left( (g_1, g_2) \cdot E_{[e]}^{(s_1, s_2)}(\chi) \right) = \sum_{T^\square \in \text{Herm}_{2r}^\circ(F)^+} d_{(g_1, g_2)}^{(s_1, s_2)} \mathscr{W}_{T^\square}^{[e]}(\chi) \cdot q^{T^\square},$$

where  $q_{2r}^{\text{an}}$  is the analytic  $q$ -expansion (2.5).

*Proof.* Note that  $\|2e + \varepsilon^c\| = 4e + 1 > 0$ , which implies that  $f_{\chi_p}^{[2e + \varepsilon^c]}(T^\square) = 0$  for  $T^\square \in \text{Herm}_{2r}(F) \setminus \text{Herm}_{2r}^\circ(F)$ . Then the lemma follows from the discussion in [Liu11b, Section 2B] and Lemma 3.2.  $\square$

**Notation 4.34.** We denote by  $(\mathbb{S}_{O_L}^\diamond)_{\hat{\pi}, L}^\star$  (resp.  $(\mathbb{S}_{O_L}^\diamond)_{\hat{\pi}, L}^\star$ ) the subset of  $(\mathbb{S}_{O_L}^\diamond)_{\hat{\pi}, L}^2$  (resp.  $(\mathbb{S}_{O_L}^\diamond)_{\hat{\pi}, L}^2$ ) consisting of  $s_1$  (resp.  $s_2$ ) such that Hypothesis 4.4 holds for  $s_1 \phi_1^{[e](k)}$  (resp.  $s_2 \phi_2^{[e](k)}$ ) for every  $k \in \Xi$  and every  $e \in \mathbb{N}$ . It is clear that both  $(\mathbb{S}_{O_L}^\diamond)_{\hat{\pi}, L}^\star$  and  $(\mathbb{S}_{O_L}^\diamond)_{\hat{\pi}, L}^\star$  are ideals of  $\mathbb{S}_{O_L}^\diamond$ .

Take a pair  $(s_1, s_2) \in (\mathbb{S}_{O_L}^\diamond)_{\hat{\pi}, L}^\star \times (\mathbb{S}_{O_L}^\diamond)_{\hat{\pi}, L}^\star$  and an element  $e \in \mathbb{N}$ . For every  $k \in \Xi$ , we have

$$\mathcal{Z}_{t_1 s_1 \phi_1^{[e](k)}, L} \in \mathcal{H}_r^{[r]} \otimes_{\mathbb{Q}_p} H_f^1(E, V_{\hat{\pi}, L}) \otimes_{\mathbb{L}} \mathbb{C}, \quad \mathcal{Z}_{t_2 s_2 \phi_2^{[e](k)}, L} \in \mathcal{H}_r^{[r]} \otimes_{\mathbb{Q}_p} H_f^1(E, V_{\pi, L}) \otimes_{\mathbb{L}} \mathbb{C}$$

from Proposition 4.6 and hence the  $p$ -adic height pairing

$$\langle \mathcal{Z}_{t_1 s_1 \phi_1^{[e](k)}, L}, \mathcal{Z}_{t_2 s_2 \phi_2^{[e](k)}, L} \rangle_E \in \Gamma_{E,p} \otimes_{\mathbb{Z}_p} \mathcal{H}_{r,r}^{[r]} \otimes_{\mathbb{Q}_p} \mathbb{C}.$$

Put

$$\mathcal{I}_{[e]}^{(s_1, s_2)} := p^M \sum_{k \in \Xi} \text{Nm}_{E/F} \left( \text{vol}^{\natural}(L) \langle \mathcal{Z}_{t_1 s_1 \phi_1^{[e](k)}, L}, \mathcal{Z}_{t_2 s_2 \phi_2^{[e](k)}, L} \rangle_E \right) \in \Gamma_{F,p} \otimes_{\mathbb{Z}_p} \mathcal{H}_{r,r}^{[r]} \otimes_{\mathbb{Q}_p} \mathbb{C} = \Gamma_{F,p} \otimes_{\mathbb{Z}_p} \mathcal{A}_{r,r, \text{hol}}^{[r]}.$$

Take  $v \in S_\pi$ . For every  $(s_1, s_2) \in \mathbb{S}_{O_L}^\diamond \times \mathbb{S}_{O_L}^\diamond$ , every  $e \in \mathbb{N}$  and every finite character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times = \overline{\mathbb{Q}}_p^\times$ , we define another Eisenstein series

$$v E_{[e]}^{(s_1, s_2)}(\chi) := p^M \cdot b_{2r}^\diamond(\mathbf{1})^{-1} \cdot b_{2r}^\diamond(\chi) \cdot E(-, f_\infty^{[r]} \otimes (\hat{s}_1 \hat{t}_1, \hat{s}_2 \hat{t}_2) \cdot v f_{\chi^\infty}^{[e]}) \in \mathcal{A}_{2r, \text{hol}}^{[r]},$$

where  $v f_{\chi^\infty}^{[e]}$  is obtained from  $f_{\chi^\infty}^{[e]}$  after replacing the component  $f_{\chi_v}$  by  $f_{\chi_v}^{\text{spH}}$  from Notation 3.4. The computation of local heights in §4.4 and §4.5 is integrated into the following proposition.

**Proposition 4.35.** *For every pair  $(s_1, s_2) \in (\mathbb{S}_{O_L}^\diamond)_{\hat{\pi}, L}^\star \times (\mathbb{S}_{O_L}^\diamond)_{\hat{\pi}, L}^\star$ , every  $e \in \mathbb{N}$  and every pair  $(g_1, g_2) \in P_r(F_R) \times P_r(F_R)$ , the  $q$ -expansion*

$$q_{r,r}^{\text{an}} \left( (g_1, g_2) \cdot \left( \mathcal{I}_{[e]}^{(s_1, s_2)} - \sum_{v \in S_\pi} \frac{2}{q_v^{2r} - 1} \cdot \rho_{r,r} \cdot v E_{[e]}^{(s_1, s_2)}(\mathbf{1}) \cdot [v] \right) \right)$$

belongs to  $\Gamma_{F,p}^{\text{fr}} \otimes_{\mathbb{Z}_p} \text{SF}_{r,r}(\overline{\mathbb{Z}}_p)$  and its image in  $\Gamma_{F,p}^{\text{fr}} \otimes_{\mathbb{Z}_p} \text{SF}_{r,r}(\overline{\mathbb{Z}}_p / p^{2e} \overline{\mathbb{Z}}_p)$  equals

$$\sum_{(T_1, T_2) \in \text{Herm}_r(F)^+ \times \text{Herm}_r(F)^+} \left( \sum_{\substack{T^\square \in \text{Herm}_{2r}^\circ(F)^+ \\ \partial_{r,r} T^\square = (T_1, T_2)}} \partial_{(g_1, g_2)}^{(s_1, s_2)} \mathscr{W}_{T^\square}^{[e]}(\mathbf{1}) \right) q^{T_1, T_2}.$$

Note that by Lemma 4.32(2) and Lemma 4.38,  $\partial_{(g_1, g_2)}^{(s_1, s_2)} \mathscr{W}_{T^\square}^{[e]}(\mathbf{1})$  belongs to  $\Gamma_{F,p}^{\text{fr}} \otimes_{\mathbb{Z}_p} \overline{\mathbb{Z}}_p$ .

*Proof.* For simplicity, we assume  $(g_1, g_2) = (1_{2r}, 1_{2r})$  and the general case is similar. For  $(T_1, T_2) \in \text{Herm}_r(F)^+ \times \text{Herm}_r(F)^+$ , denote by  $a_{T_1, T_2}$  and  ${}^v b_{T_1, T_2}$  the  $q^{T_1, T_2}$ -coefficients of  $\mathcal{I}_{[e]}^{(s_1, s_2)}$  and  $\rho_{r, r} {}^v E_{[e]}^{(s_1, s_2)}(\mathbf{1})$ , respectively. By (4.7), we have

$$\begin{aligned} a_{T_1, T_2} &= \mathbf{1}_{(\text{Herm}_r(F)^+)^2}(T_1, T_2) \cdot p^M \text{vol}^{\natural}(L) \cdot \sum_{k \in \Xi} \text{Nm}_{E/F} \langle Z_{T_1}^{\star}(t_1 s_1 \phi_1^{[e](k)})_L, Z_{T_2}^{\star}(t_2 s_2 \phi_2^{[e](k)})_L \rangle_E \\ &= a'_{T_1, T_2} + \mathbf{1}_{(\text{Herm}_r(F)^+)^2}(T_1, T_2) \cdot p^M \text{vol}^{\natural}(L) \cdot \sum_{k \in \Xi} \text{Nm}_{E/F} \sum_{u \nmid \infty p} \langle Z_{T_1}(t_1 s_1 \phi_1^{[e](k)})_L, Z_{T_2}(t_2 s_2 \phi_2^{[e](k)})_L \rangle_{E_u}, \end{aligned}$$

where

$$a'_{T_1, T_2} := \mathbf{1}_{(\text{Herm}_r(F)^+)^2}(T_1, T_2) \cdot p^M \text{vol}^{\natural}(L) \cdot \sum_{k \in \Xi} \text{Nm}_{E/F} \sum_{u|p} \langle Z_{T_1}(t_1 s_1 \phi_1^{[e](k)})_L, Z_{T_2}(t_2 s_2 \phi_2^{[e](k)})_L \rangle_{E_u}$$

which belongs to  $\Gamma_{F, p}^{\text{fr}} \otimes_{\mathbb{Z}_p} (p^{2e} \overline{\mathbb{Z}}_p)$  by Proposition 4.26 and the definition of  $M$ .

On the other hand,

$$\begin{aligned} b_{T_1, T_2} &= p^M W_{2r} \cdot b_{2r}^{\infty}(\mathbf{1}) \left( \sum_{\substack{T^{\square} \in \text{Herm}_{2r}^{\circ}(F)^+ \\ \partial_{r, r} T^{\square} = (T_1, T_2)}} W_{T^{\square}}((\hat{s}_1 \hat{t}_1, \hat{s}_2 \hat{t}_2) {}^v f_{\chi^{\infty}}^{[e]}) \right) \\ &= p^M W_{2r} \sum_{k \in \Xi} \left( \sum_{\substack{T^{\square} \in \text{Herm}_{2r}^{\circ}(F)^+ \\ \partial_{r, r} T^{\square} = (T_1, T_2)}} W_{T^{\square}, v}^{\text{sph}}(\mathbf{1}) \cdot I_{T^{\square}}((t_1 s_1 \phi_1^{[e](k)} \otimes t_2 s_2 \phi_2^{[e](k)})^v) \right) \end{aligned}$$

in which the second equality uses Remark 4.29.

Thus, by Proposition 4.24, it suffices to show that

$$\partial_{(g_1, g_2)}^{(s_1, s_2)} \mathcal{W}_{T^{\square}}^{[e]}(\mathbf{1}) = p^M W_{2r} \sum_{k \in \Xi} W'_{T^{\square}, v_{T^{\square}}}(\mathbf{1}) \cdot I_{T^{\square}}((t_1 s_1 \phi_1^{[e](k)} \otimes t_2 s_2 \phi_2^{[e](k)})^{v_{T^{\square}}}) \cdot [v_{T^{\square}}]$$

for every  $T^{\square} \in \text{Herm}_{2r}^{\circ}(F)^+$ . Indeed, this follows from the  $p$ -adic Leibniz rule and Remark 4.29.

The proposition is proved.  $\square$

We give a formula for the (global)  $p$ -adic height pairing between the Selmer theta lifts with respect to our chosen data in the following proposition.

**Proposition 4.36.** *For every collection of pairs  $(s_1, s_2) \in (\mathbb{S}_{O_L}^{\diamond})_{\hat{\pi}, L}^{\star} \times (\mathbb{S}_{O_L}^{\diamond})_{\pi, L}^{\star}$ , we have*

$$(4.15) \quad \sum_{k \in \Xi} \langle \Theta_{t_1 s_1 \phi_1^{(k)}}^{\text{Sel}}(\varphi_1), \Theta_{t_2 s_2 \phi_2^{(k)}}^{\text{Sel}}(\varphi_2) \rangle_{\pi, F}^{\natural} = \chi_{\hat{\pi}}^{\diamond}(t_1 s_1) \chi_{\pi}^{\diamond}(t_2 s_2) \cdot \partial \mathcal{L}_p^{\diamond}(\pi)(\mathbf{1}) \cdot \prod_{v \in V_F^{(\diamond \setminus \{\infty\})}} Z(\varphi_{1, v}^{\dagger} \otimes \varphi_{2, v}, f_{\Phi_v}^{\text{SW}}).$$

We briefly explain the idea before the proof. We will compare two kernel functions, namely,  $\mathcal{I}_{[0]}^{(s_1, s_2)}$  on the geometric side and the “ $p$ -adic derivative” of  $\rho_{r, r} E_{[0]}^{(s_1, s_2)}$  at  $\mathbf{1}$  on the analytic side. It turns out that they differ by two factors: one consists of pullback Eisenstein series  $\rho_{r, r} {}^v E_{[0]}^{(s_1, s_2)}(\mathbf{1})$  and the other consists of local  $p$ -adic height pairings between generating functions at  $p$ -adic places. For the first one, by the local theta dichotomy,  $\rho_{r, r} {}^v E_{[0]}^{(s_1, s_2)}(\mathbf{1})$  has vanishing pairing with  $\varphi_1 \otimes \varphi_2$ . For the second one, we use an idea originally due to Perrin-Riou [PR87]: the local height pairings converge to 0 in the  $p$ -adic topology when one repeatedly applies the operator  $U_p$  to the Schwartz function; on the other hand, after taking the pairing with  $\varphi_1 \otimes \varphi_2$ , the operator  $U_p$  changes the result by an element in  $O_L^{\times}$ .

*Proof.* We first introduce an open compact subgroup

$$K := \prod_{v \in \mathbb{R}} K_v \prod_{v \in V_F^{(p)}} I_v \prod_{v \in \mathbb{T}} K_{r, v} \subseteq G_r(\mathbb{A}_F^{\infty})$$

in which  $K_v$  is the one chosen in (R5) and  $I_v = \mathcal{G}_r(O_{F_v}) \times_{\mathcal{G}_r(O_{F_v}/\varpi_v)} \mathcal{P}_r(O_{F_v}/\varpi_v)$  is the one used in §3.3. By (4.4), the map

$$\begin{aligned} \mathcal{A}_{r,r,\text{hol}}^{[r]}(K \times K) &\rightarrow \text{SF}_{r,r}(\mathbb{C})^{P_r(F_{\mathbb{R}}) \times P_r(F_{\mathbb{R}})} \\ \Psi &\mapsto \left( \mathbf{q}_{r,r}^{\text{an}}((g_1, g_2) \cdot \Psi) \right)_{(g_1, g_2) \in P_r(F_{\mathbb{R}}) \times P_r(F_{\mathbb{R}})} \end{aligned}$$

is injective. In particular, the subset

$$\mathcal{A}_{r,r,\text{hol}}^{[r]}(K \times K)^+ := \left\{ \Psi \in \mathcal{A}_{r,r,\text{hol}}^{[r]}(K \times K) \mid \mathbf{q}_{r,r}^{\text{an}}((g_1, g_2) \cdot \Psi) \in \text{SF}_{r,r}(\overline{\mathbb{Z}}_p) \text{ for every } (g_1, g_2) \in P_r(F_{\mathbb{R}}) \times P_r(F_{\mathbb{R}}) \right\}$$

is a finitely generated  $\overline{\mathbb{Z}}_p$ -module. Thus, we may choose an integer  $M_0 \geq 0$  such that

$$p^{M_0} \langle \varphi_1 \otimes \varphi_2, \Psi \rangle_{\pi, \hat{\pi}} \in \overline{\mathbb{Z}}_p$$

for every  $\Psi \in \mathcal{A}_{r,r,\text{hol}}^{[r]}(K \times K)^+$ .

We claim that there is a unique integral  $\overline{\mathbb{Q}}_p$ -valued  $p$ -adic measure  $\mathbf{d}\mathcal{M}^{(s_1, s_2)}$  on  $\Gamma_{F,p}$  such that

$$\mathcal{M}^{(s_1, s_2)}(\chi) := p^{M_0} \langle \varphi_1 \otimes \varphi_2, \rho_{r,r} E_{[0]}^{(s_1, s_2)}(\chi) \rangle_{\pi, \hat{\pi}}$$

holds for every finite character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times = \overline{\mathbb{Q}}_p^\times$ . Indeed, take an arbitrary open compact subset  $\Omega \subseteq \Gamma_{F,p}$ . Write  $\mathbf{1}_\Omega = \sum_i c_i \cdot \chi_i$  as a finite sum in a unique way with  $c_i \in \mathbb{C}$  and finite characters  $\chi_i: \Gamma_{F,p} \rightarrow \mathbb{C}^\times$ . By the same argument in the proof of Theorem 3.37, we may choose an element  $e \in \mathbb{N}$  such that  $(U_p^e \times U_p^e) f_{\chi_p}$  is invariant under  $I_v \times I_v$  for every  $i$ . Then  $(U_p^e \times U_p^e) \rho_{r,r} E_{[0]}^{(s_1, s_2)}(\chi_i)$  is invariant under  $K \times K$ . By Lemma 4.20(1), we have

$$p^{M_0} \langle \varphi_1 \otimes \varphi_2, \rho_{r,r} E_{[0]}^{(s_1, s_2)}(\chi) \rangle_{\pi, \hat{\pi}} = p^{M_0} \left( \prod_{u \in \mathbb{P}} \alpha(\pi_u) \right)^{-2e} \left\langle \varphi_1 \otimes \varphi_2, (U_p^e \times U_p^e) \rho_{r,r} \sum_i c_i E_{[0]}^{(s_1, s_2)}(\chi_i) \right\rangle_{\pi, \hat{\pi}}.$$

It remains to show that

$$(U_p^e \times U_p^e) \rho_{r,r} \sum_i c_i E_{[0]}^{(s_1, s_2)}(\chi_i) = \rho_{r,r} \sum_i c_i E_{[e]}^{(s_1, s_2)}(\chi_i) \in \mathcal{A}_{r,r,\text{hol}}^{[r]}(K \times K)^+,$$

where the equality comes from Lemma 4.20(1). However, this follows from Lemma 4.33 and Lemma 4.32(2). The claim is justified.

As  $\mathcal{L}_p^\diamond(\pi)(\mathbf{1}) = 0$  and  $Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{1,v}) \neq 0$  for  $v \in \mathbb{T} \cap \mathbb{V}_F^{(\diamond)}$  by Lemma 4.31, Lemma 4.30 and Theorem 3.37 imply that

$$\partial \mathcal{M}^{(s_1, s_2)}(\mathbf{1}) = p^{M+M_0} \cdot \chi_{\hat{\pi}}^\diamond(t_1 s_1) \chi_{\pi}^\diamond(t_2 s_2) \cdot \partial \mathcal{L}_p^\diamond(\pi)(\mathbf{1}) \cdot \prod_{v \in \mathbb{V}_F^{(\diamond) \setminus \{\infty\}}} Z(\varphi_{1,v}^\dagger \otimes \varphi_{2,v}, f_{\Phi_v}^{\text{SW}})$$

by the  $p$ -adic Leibniz rule. Thus, for (4.15), it remains to show that

$$p^{M_0} \sum_{k \in \Xi} \langle \Theta_{t_1 s_1 \phi_1^{(k)}}^{\text{Sel}}(\varphi_1), \Theta_{t_2 s_2 \phi_2^{(k)}}^{\text{Sel}}(\varphi_2) \rangle_{\pi, F}^{\natural} = \partial \mathcal{M}^{(s_1, s_2)}(\mathbf{1}).$$

For this, it suffices to show that for every  $\lambda \in \text{Hom}_{\mathbb{Z}_p}(\Gamma_{F,p}, \mathbb{Z}_p)$  and every integer  $d \geq 1$ ,

$$(4.16) \quad \partial_\lambda \mathcal{M}^{(s_1, s_2)}(\mathbf{1}) \equiv \lambda \left( p^{M_0} \sum_{k \in \Xi} \langle \Theta_{t_1 s_1 \phi_1^{(k)}}^{\text{Sel}}(\varphi_1), \Theta_{t_2 s_2 \phi_2^{(k)}}^{\text{Sel}}(\varphi_2) \rangle_{\pi, F}^{\natural} \right) \pmod{p^d \overline{\mathbb{Z}}_p}.$$

Take such  $\lambda$  and  $d$ . We choose an element  $e \in \mathbb{N}$  satisfying  $2e \geq d$  and such that  $(U_p^e \times U_p^e) f_{\chi_p}$  is invariant under  $I_v \times I_v$  for every character  $\chi: \Gamma_{F,p} \rightarrow \mathbb{C}^\times$  that factors through  $U_d$ .

Since  $\mathbf{d}\mathcal{M}^{(s_1, s_2)}$  is integral, by Lemma 4.38, we have

$$\partial_\lambda \mathcal{M}^{(s_1, s_2)}(\mathbf{1}) \equiv \sum_{x \in \Gamma_d} \lambda(x) \text{vol}(x U_d, \mathbf{d}\mathcal{M}^{(s_1, s_2)}) \pmod{p^d \overline{\mathbb{Z}}_p}.$$

However,

$$\begin{aligned} \sum_{x \in \Gamma_d} \lambda(x) \operatorname{vol}(xU_d, \mathbf{d} \mathcal{M}^{(s_1, s_2)}) &= \sum_{x \in \Gamma_d} \lambda(x) \frac{\chi(x)^{-1}}{|\Gamma_{F,p}/U_d|} \sum_{\chi: \Gamma_{F,p}/U_d \rightarrow \mathbb{C}^\times} \mathcal{M}^{(s_1, s_2)}(\chi) \\ &= \sum_{x \in \Gamma_d} \lambda(x) \frac{\chi(x)^{-1}}{|\Gamma_{F,p}/U_d|} \sum_{\chi: \Gamma_{F,p}/U_d \rightarrow \mathbb{C}^\times} p^{M_0} \langle \varphi_1 \otimes \varphi_2, \rho_{r,r} E_{[0]}^{(s_1, s_2)}(\chi) \rangle_{\pi, \hat{\pi}}, \end{aligned}$$

which, by Lemma 4.20(1), equals

$$\begin{aligned} &p^{M_0} \left( \prod_{u \in \mathbb{P}} \alpha(\pi_u) \right)^{-2e} \sum_{x \in \Gamma_d} \lambda(x) \frac{\chi(x)^{-1}}{|\Gamma_{F,p}/U_d|} \sum_{\chi: \Gamma_{F,p}/U_d \rightarrow \mathbb{C}^\times} \langle \varphi_1 \otimes \varphi_2, \rho_{r,r} E_{[e]}^{(s_1, s_2)}(\chi) \rangle_{\pi, \hat{\pi}} \\ &= p^{M_0} \left( \prod_{u \in \mathbb{P}} \alpha(\pi_u) \right)^{-2e} \left\langle \varphi_1 \otimes \varphi_2, \sum_{x \in \Gamma_d} \lambda(x) \frac{\chi(x)^{-1}}{|\Gamma_{F,p}/U_d|} \sum_{\chi: \Gamma_{F,p}/U_d \rightarrow \mathbb{C}^\times} \rho_{r,r} E_{[e]}^{(s_1, s_2)}(\chi) \right\rangle_{\pi, \hat{\pi}}. \end{aligned}$$

On the other hand, by Definition 4.12, for every  $e \in \mathbb{N}$ ,

$$(4.17) \quad \sum_{k \in \Xi} \langle \Theta_{\mathfrak{t}_1 s_1 \phi_1^{[e](k)}}^{\operatorname{Sel}}(\varphi_1), \Theta_{\mathfrak{t}_2 s_2 \phi_2^{[e](k)}}^{\operatorname{Sel}}(\varphi_2) \rangle_{\pi, F}^{\natural} = \langle \varphi_1 \otimes \varphi_2, \mathcal{I}_{[e]}^{(s_1, s_2)} \rangle_{\pi, \hat{\pi}}.$$

Since for  $v \in S_\pi$ , the local theta lift of  $\pi_v$  to the split hermitian space over  $E_v$  of rank  $2r$  vanishes, we have

$$\langle \varphi_1 \otimes \varphi_2, \rho_{r,r} {}^v E_{[e]}^{(s_1, s_2)}(\mathbf{1}) \rangle_{\pi, \hat{\pi}} = 0.$$

It follows that

$$(4.18) \quad (4.17) = \left\langle \varphi_1 \otimes \varphi_2, \mathcal{I}_{[e]}^{(s_1, s_2)} - \sum_{v \in S_\pi} \frac{2}{q_v^{2r} - 1} \cdot \rho_{r,r} {}^v E_{[e]}^{(s_1, s_2)}(\mathbf{1}) \cdot [v] \right\rangle_{\pi, \hat{\pi}}.$$

By Lemma 4.20(1) and (4.18),

$$\text{RHS of (4.16)} = p^{M_0} \left( \prod_{u \in \mathbb{P}} \alpha(\pi_u) \right)^{-2e} \left\langle \varphi_1 \otimes \varphi_2, \lambda \left( \mathcal{I}_{[e]}^{(s_1, s_2)} - \sum_{v \in S_\pi} \frac{2}{q_v^{2r} - 1} \cdot \rho_{r,r} {}^v E_{[e]}^{(s_1, s_2)}(\mathbf{1}) \cdot [v] \right) \right\rangle_{\pi, \hat{\pi}}.$$

Put

$$\Psi_d := \sum_{x \in \Gamma_d} \lambda(x) \frac{\chi(x)^{-1}}{|\Gamma_{F,p}/U_d|} \sum_{\chi: \Gamma_{F,p}/U_d \rightarrow \mathbb{C}^\times} \rho_{r,r} E_{[e]}^{(s_1, s_2)}(\chi) - \lambda \left( \mathcal{I}_{[e]}^{(s_1, s_2)} - \sum_{v \in S_\pi} \frac{2}{q_v^{2r} - 1} \cdot \rho_{r,r} {}^v E_{[e]}^{(s_1, s_2)}(\mathbf{1}) \cdot [v] \right).$$

Now by our choice of  $e$ , we have  $\Psi_d \in \mathcal{A}_{r,r, \text{hol}}^{[r]}(K \times K)$ . Thus, (4.16) follows if we can show  $\Psi_d \in p^d \mathcal{A}_{r,r, \text{hol}}^{[r]}(K \times K)^+$ , that is, for every pair  $(g_1, g_2) \in P_r(F_R) \times P_r(F_R)$ ,

$$\mathbf{q}_{r,r}^{\text{an}}((g_1, g_2) \cdot \Psi_d) \in p^d \mathbf{SF}_{r,r}(\overline{\mathbb{Z}}_p).$$

For  $(T_1, T_2) \in \operatorname{Herm}_r(F)^+ \times \operatorname{Herm}_r(F)^+$ , denote by  $c_{T_1, T_2}$  the  $q^{T_1, T_2}$ -coefficient of  $\mathbf{q}_{r,r}^{\text{an}}((g_1, g_2) \cdot \Psi_d)$ . By Lemma 4.33 and Proposition 4.35,

$$\begin{aligned} c_{T_1, T_2} &\equiv \sum_{x \in \Gamma_d} \lambda(x) \frac{\chi(x)^{-1}}{|\Gamma_{F,p}/U_d|} \sum_{\chi: \Gamma_{F,p}/U_d \rightarrow \mathbb{C}^\times} \sum_{\substack{T^\square \in \operatorname{Herm}_{2r}^\circ(F)_V^+ \\ \partial_{r,r} T^\square = (T_1, T_2)}} \mathscr{W}_{T^\square}^{[e]}(\chi) - \lambda \sum_{\substack{T^\square \in \operatorname{Herm}_{2r}^\circ(F)_V^+ \\ \partial_{r,r} T^\square = (T_1, T_2)}} \partial_{(g_1, g_2)} \mathscr{W}_{T^\square}^{[e]}(\mathbf{1}) \\ &\equiv \sum_{\substack{T^\square \in \operatorname{Herm}_{2r}^\circ(F)_V^+ \\ \partial_{r,r} T^\square = (T_1, T_2)}} \left( \sum_{x \in \Gamma_d} \lambda(x) \frac{\chi(x)^{-1}}{|\Gamma_{F,p}/U_d|} \sum_{\chi: \Gamma_{F,p}/U_d \rightarrow \mathbb{C}^\times} \mathscr{W}_{T^\square}^{[e]}(\chi) - \partial_\lambda \mathscr{W}_{T^\square}^{[e]}(\mathbf{1}) \right) \\ &\equiv \sum_{\substack{T^\square \in \operatorname{Herm}_{2r}^\circ(F)_V^+ \\ \partial_{r,r} T^\square = (T_1, T_2)}} \left( \sum_{x \in \Gamma_d} \lambda(x) \operatorname{vol}(xU_d, \mathbf{d} \mathscr{W}_{T^\square}^{[e]}) - \partial_\lambda \mathscr{W}_{T^\square}^{[e]}(\mathbf{1}) \right) \pmod{p^d \overline{\mathbb{Z}}_p}. \end{aligned}$$

Finally, by Lemma 4.32(2) and Lemma 4.38, we have

$$\sum_{x \in \Gamma_d} \lambda(x) \operatorname{vol}(xU_d, \mathbf{d}_{(g_1, g_2)}^{(s_1, s_2)} \mathscr{W}_{T^{\square}}^{[e]}) - \partial \lambda_{(g_1, g_2)}^{(s_1, s_2)} \mathscr{W}_{T^{\square}}^{[e]}(\mathbf{1}) \equiv 0 \pmod{p^d \overline{\mathbb{Z}}_p}.$$

Thus, (4.16) is confirmed. The proposition is proved.  $\square$

*Proof of Theorem 1.7 and Theorem 4.17.* We start from two common ingredients for the proof of both theorems.

- (1) One can find  $(s_1, s_2) \in (\mathbb{S}_{O_L}^{\diamond})_{\pi, L}^2 \times (\mathbb{S}_{O_L}^{\diamond})_{\pi, L}^2$  satisfying  $\chi_{\pi}^{\diamond}(s_1)\chi_{\pi}^{\diamond}(s_2) \neq 0$ . Indeed, by [LL21, Proposition 6.9], the localizations of  $H_{\text{ét}}^{2r}(X_L \otimes_E \overline{E}, \mathbb{L}(r))$  at both  $\operatorname{Ker} \chi_{\pi}^{\diamond}$  and  $\operatorname{Ker} \chi_{\pi}^{\diamond}$  vanish. Also, by Matsushima's formula, the localizations of  $H_{\text{ét}}^{2r-1}(X_L \otimes_E \overline{E}, \mathbb{L}(r))/V_{\pi, L}$  and  $H_{\text{ét}}^{2r-1}(X_L \otimes_E \overline{E}, \mathbb{L}(r))/V_{\hat{\pi}, L}$  at  $\operatorname{Ker} \chi_{\pi}^{\diamond}$  and  $\operatorname{Ker} \chi_{\hat{\pi}}^{\diamond}$  vanish, respectively. Thus, the claim follows from the finite-dimensionality of the spaces involved.
- (2) In Proposition 4.36, we have

$$\prod_{v \in \mathbf{V}_F^{(\diamond) \setminus \{\infty\}}} Z(\varphi_{1, v}^{\dagger} \otimes \varphi_{2, v}, f_{\Phi_v}^{\text{SW}}) \neq 0.$$

Indeed, for  $v \in \mathbf{R}$ ,  $Z(\varphi_{1, v}^{\dagger} \otimes \varphi_{2, v}, f_{\Phi_v}^{\text{SW}}) = N_v \neq 0$ ; for  $v \in \mathbf{T} \cap \mathbf{V}_F^{(\diamond)}$ ,  $Z(\varphi_{1, v}^{\dagger} \otimes \varphi_{2, v}, f_{\Phi_v}^{\text{SW}}) \neq 0$  by Lemma 4.31; for  $v \in \mathbf{V}_F^{(p)}$ ,  $Z(\varphi_{1, v}^{\dagger} \otimes \varphi_{2, v}, f_{\Phi_v}^{\text{SW}}) \neq 0$  by Lemma 4.20(1) and Proposition 3.31.

We now show Theorem 1.7. Assume the converse, that is,  $H_f^1(E, \rho_{\pi}(r)) = 0$ . By [LTX<sup>+</sup>22, Lemma 2.4.5], we also have  $H_f^1(E, \rho_{\hat{\pi}}(r)) = H_f^1(E, \rho_{\pi}(r)) = 0$ . We claim that  $H_f^1(E, \rho_{\pi}(r)) = 0$  and  $H_f^1(E, \rho_{\hat{\pi}}(r)) = 0$  imply  $H_f^1(E, V_{\pi, L}) = 0$  and  $H_f^1(E, V_{\hat{\pi}, L}) = 0$ , respectively. By symmetry, it suffices to consider the former case. Indeed, by Lemma 4.14 and Lemma 4.15,  $V_{\pi, L}$  is crystalline and pure of weight  $-1$ . Thus, by [Nek93, Proposition 1.25] and Lemma 4.13, it suffices to show that for every irreducible  $\mathbb{L}[\operatorname{Gal}(\overline{E}/E)]$ -module  $\rho$  that is a subquotient of  $V_{\pi, L}$ , we have  $H_f^1(E, \rho) = 0$ . However, this follows from Hypothesis 4.9. The claim implies that  $(\mathbb{S}_{O_L}^{\diamond})_{\pi, L}^{\star} = (\mathbb{S}_{O_L}^{\diamond})_{\hat{\pi}, L}^2$  and  $(\mathbb{S}_{O_L}^{\diamond})_{\pi, L}^{\star} = (\mathbb{S}_{O_L}^{\diamond})_{\pi, L}^2$ . We take the pair  $(s_1, s_2)$  as in (1). Then by (2), the right-hand side of (4.15) is nonvanishing, while the left-hand side vanishes. This is a contradiction and hence Theorem 1.7 follows.

We now show Theorem 4.17. We have mentioned in Remark 4.18(2) that one may enlarge  $\diamond$  so that (4.4) is satisfied. It suffices to prove (4.3) along an arbitrary isomorphism  $\overline{\mathbb{Q}}_p \xrightarrow{\sim} \mathbb{C}$ . Then by Remark 4.18(3), it suffices to show (the linear extension of) (4.3) for one element  $\Phi \in \mathbf{I}_r^{\square}(\mathbf{1}^{\infty})$  with  $\Phi_v = \mathbf{1}_{\Lambda_v^{2r}}$  for every  $v \in \mathbf{V}_F \setminus \mathbf{V}_F^{(\diamond)}$  satisfying  $\prod_{v \in \mathbf{V}_F^{(\diamond) \setminus \{\infty\}}} Z(\varphi_{1, v}^{\dagger} \otimes \varphi_{2, v}, f_{\Phi_v}^{\text{SW}}) \neq 0$ . However, in view of (2) and Remark 4.29, (4.15) already gave such a formula with the pair  $(s_1, s_2)$  in (1). Theorem 4.17 is proved.  $\square$

**4.7. Generalities on  $p$ -adic measures.** In this subsection, we review some general facts about derivatives of  $p$ -adic measures. For  $d \geq 1$ , we denote by  $U_d$  the image of  $1 + O_F \otimes (p^d \mathbb{Z}_p)$  in  $\Gamma_{F, p}$ , which is an open subgroup of finite index.

Let  $d\mu$  be an  $\mathbb{M}$ -valued  $p$ -adic measure on  $\Gamma_{F, p}$  for a finite extension  $\mathbb{M}/\mathbb{Q}_p$ . For every continuous character  $\chi: \Gamma_{F, p} \rightarrow R^{\times}$  for a complete  $\mathbb{M}$ -ring  $R$ , we put

$$\mu(\chi) := \int_{\Gamma_{F, p}} \chi \, d\mu := \lim_{d \rightarrow \infty} \sum_{x \in \Gamma_d} \chi(x) \operatorname{vol}(xU_d, d\mu)$$

where  $\Gamma_1 \subseteq \Gamma_2 \subseteq \dots$  is an arbitrary increasing chain of sets of representatives of  $\Gamma_{F, p}/U_d$  for  $d = 1, 2, \dots$ . Then  $\mu(\chi)$  does not depend on  $(\Gamma_d)_d$  and defines a bounded rigid analytic function on  $\mathscr{X}_{F, p} \otimes_{\mathbb{Q}_p} \mathbb{M}$ . We consider its derivative  $\partial \mu(\mathbf{1})$  at  $\mathbf{1}$ , which is an element in  $\Gamma_{F, p} \otimes_{\mathbb{Z}_p} \mathbb{L}$  – the cotangent space of  $\mathscr{X}_{F, p} \otimes_{\mathbb{Q}_p} \mathbb{L}$  at  $\mathbf{1}$ . More precisely,  $\partial \mu(\mathbf{1})$  is the linear functional in  $\operatorname{Hom}_{\mathbb{Z}_p}(\operatorname{Hom}_{\mathbb{Z}_p}(\Gamma_{F, p}, \mathbb{Z}_p), \mathbb{L})$  that sends  $\lambda \in \operatorname{Hom}_{\mathbb{Z}_p}(\Gamma_{F, p}, \mathbb{Z}_p)$  to

$$\partial \lambda \mu(\mathbf{1}) := \lim_{c \rightarrow \infty} \frac{1}{p^c} (\mu(\exp(p^c \lambda)) - \mu(\mathbf{1})) = \lim_{c \rightarrow \infty} \frac{1}{p^c} \lim_{d \rightarrow \infty} \sum_{x \in \Gamma_d} (\exp(p^c \lambda(x)) - 1) \operatorname{vol}(xU_d, d\mu).$$

Since

$$\frac{1}{p^c} (\exp(p^c \lambda(x)) - 1) = \lambda(x) + \frac{p^c \lambda(x)^2}{2!} + \frac{p^{2c} \lambda(x)^3}{3!} + \dots,$$

and  $\text{vol}(xU_d, d\mu)$  is bounded independent of  $x$  and  $d$ , we have

$$(4.19) \quad \partial_\lambda \mu(\mathbf{1}) = \lim_{d \rightarrow \infty} \sum_{x \in \Gamma_d} \lambda(x) \text{vol}(xU_d, d\mu).$$

**Definition 4.37.** By a  $\overline{\mathbb{Q}}_p$ -valued  $p$ -adic measure we mean an  $\mathbb{M}$ -valued  $p$ -adic measure for some finite extension  $\mathbb{M}/\mathbb{Q}_p$  contained in  $\overline{\mathbb{Q}}_p$ . We say that a  $\overline{\mathbb{Q}}_p$ -valued  $p$ -adic measure  $d\mu$  on  $\Gamma_{F,p}$  is *integral* if  $\text{vol}(\Omega, d\mu) \in \overline{\mathbb{Z}}_p$  for every open compact subset  $\Omega \subseteq \Gamma_{F,p}$ .

**Lemma 4.38.** *Let  $d\mu$  be an integral  $\overline{\mathbb{Q}}_p$ -valued  $p$ -adic measure on  $\Gamma_{F,p}$ . Then for every  $\lambda \in \text{Hom}_{\mathbb{Z}_p}(\Gamma_{F,p}, \overline{\mathbb{Z}}_p)$  and every  $d \geq 1$ , we have*

$$\partial_\lambda \mu(\mathbf{1}) - \sum_{x \in \Gamma_d} \lambda(x) \text{vol}(xU_d, d\mu) \in p^d \overline{\mathbb{Z}}_p.$$

In particular,  $\partial_\lambda \mu(\mathbf{1}) \in \Gamma_{F,p}^{\text{fr}} \otimes_{\mathbb{Z}_p} \overline{\mathbb{Z}}_p$ .

*Proof.* Since  $U_d \subseteq p^d \Gamma_{F,p}$ , we have  $\lambda(x) - \lambda(x') \in p^d \overline{\mathbb{Z}}_p$  if  $x = x'$  in  $\Gamma_{F,p}/U_d$ . Then the lemma follows from (4.19) since  $d\mu$  is integral.  $\square$

## APPENDIX A. BI-EXTENSIONS AND $p$ -ADIC HEIGHT PAIRINGS

In this appendix, we develop further the theory of  $p$ -adic heights on general varieties. We fix a prime number  $p$ , a finite extension  $\mathbb{L}/\mathbb{Q}_p$  contained in  $\overline{\mathbb{Q}}_p$ , and an integer  $n \geq 2$ .

**A.1. Étale correspondences.** Let  $X$  be a scheme. An *étale correspondence* on  $X$  is a diagram

$$t: X \xleftarrow{f} X' \xrightarrow{g} X$$

in which both  $f$  and  $g$  are finite étale morphisms.

The collection of all étale correspondences on  $X$  forms a monoidal category  $\text{ÉtCor}(X)$ . See [Liu19, Definition 2.11] for more details. We denote by  $\text{EC}(X)$  the (unital)  $\mathbb{Z}$ -algebra generated by the underlying monoid of isomorphism classes of objects of  $\text{ÉtCor}(X)$ . For every ring  $L$ , an  *$L$ -ring of étale correspondences* on  $X$  is an  $L$ -ring  $\mathbb{T}$  together with a homomorphism  $\mathbb{T} \rightarrow \text{EC}(X)_L$  that is  $L$ -linear and unital. Usually, we only write  $\mathbb{T}$  for the notation when the homomorphism  $\mathbb{T} \rightarrow \text{EC}(X)_L$  is clear.

**Notation A.1.** Let  $S$  be a subset of  $X$ . For an étale correspondence  $t$  as above, we put  $S^t := f(g^{-1}(S))$ . For a finite linear combination  $t = \sum c_i t_i$  with  $c_i \neq 0$ , we put  $S^t := \bigcup_i S^{t_i}$ .

**A.2. Remarks on sheaves.** For a ring  $L$  and a site  $S$ , we denote by

- $\mathbf{M}(S, L)$  the abelian category of sheaves of  $L$ -modules on  $S$ ,
- $\mathbf{C}^+(S, L) = \mathbf{C}^+(\mathbf{M}(S, L))$  the abelian category of bounded below complexes in  $\mathbf{M}(S, L)$ ,
- $\mathbf{D}^+(S, L) = \mathbf{D}^+(\mathbf{M}(S, L))$  the derived category of  $\mathbf{M}(S, L)$  with bounded below cohomology.

Let  $K$  be a field with a fixed algebraic closure  $\overline{K}$ . Let  $G_K$  be the underlying abstract group of the profinite group  $\text{Gal}(\overline{K}/K)$ . For a ring  $L$  and a site  $S$  with an action of  $G_K$ , we denote by

- $\mathbf{M}_K(S, L)$  the abelian category of compatible  $G_K$ -equivariant sheaves of  $L$ -modules on  $S$ ,
- $\mathbf{C}_K^+(S, L) = \mathbf{C}^+(\mathbf{M}_K(S, L))$  the abelian category of bounded below complexes in  $\mathbf{M}_K(S, L)$ ,
- $\mathbf{D}_K^+(S, L) = \mathbf{D}^+(\mathbf{M}_K(S, L))$  the derived category of  $\mathbf{M}_K(S, L)$  with bounded below cohomology.

We suppress  $S$  (with the comma) in the notation when  $S$  is a singleton. In particular,  $\mathbf{M}_K(L)$  is nothing but the category of  $L[G_K]$ -modules, and we have the global section functor  $\Gamma(S, -): \mathbf{M}_K(S, L) \rightarrow \mathbf{M}_K(L)$ . We also

- denote by  $\Phi_K: \mathbf{M}_K(L) \rightarrow \mathbf{M}(L)$  the function taking  $G_K$ -invariants, and
- put  $\Gamma_K(S, -) := \Phi_K \circ \Gamma(S, -): \mathbf{M}_K(S, L) \rightarrow \mathbf{M}(L)$ .

Let  $q$  be an integer.

- We put  $\mathbf{H}^q(K, -) := \mathbf{R}^q \Phi_K(-)$ .
- For  $? \in \{ , K\}$ , we put  $\mathbf{H}_?^q(S, -) := \mathbf{R}^q \Gamma_?(S, -)$ .

- For  $\mathcal{C} \in \mathbf{D}_K^+(S)$ , we put

$$\mathbf{H}_K^q(S, \mathcal{C})^0 := \text{Ker}\left(\mathbf{H}_K^q(S, \mathcal{C}) \rightarrow \mathbf{H}^0(K, \mathbf{H}^q(S, \mathcal{C}))\right).$$

We have a Hochschild–Serre spectral sequence for  $\mathbf{H}_K^\bullet(S, -)$  as the Grothendieck spectral sequence for the composite derived functor  $\mathbf{R}\Gamma_K(S, -) = \mathbf{R}\Phi_K \circ \mathbf{R}\Gamma(S, -)$ . In particular, we have an edge map

$$\mathbf{H}_K^q(S, \mathcal{C})^0 \rightarrow \mathbf{H}^1(K, \mathbf{H}^{q-1}(S, \mathcal{C})).$$

Now we clarify our convention on the  $p$ -adic formalism of étale cohomology. For a scheme  $X$  and an integer  $r$ , we denote by

$$\mathbb{L}(r)_X := \left( \mathbf{R} \lim_{\leftarrow l} \mu_{p^l}^{\otimes r} \right) \otimes_{\mathbb{Z}_p} \mathbb{L} \in \mathbf{D}^+(X_{\text{ét}}, \mathbb{L})$$

For an exact functor  $\Psi: \mathbf{D}^+(X_{\text{ét}}, \mathbb{Z}_p) \rightarrow \mathbf{D}^+(Y_{\text{ét}}, \mathbb{Z}_p)$ , we denote by

$$\Psi \mathbb{L}(r)_X := \left( \mathbf{R} \lim_{\leftarrow l} \Psi \mu_{p^l}^{\otimes r} \right) \otimes_{\mathbb{Z}_p} \mathbb{L} \in \mathbf{D}^+(Y_{\text{ét}}, \mathbb{L}).$$

In particular, by the Grothendieck spectral sequence for composition of derived functors, we have

- For every scheme  $X$ ,  $\mathbf{H}^q(X_{\text{ét}}, \mathbb{L}(r)_X)$  coincides with Jannsen’s continuous étale cohomology [Jan88] of  $X$ , usually denoted by  $\mathbf{H}^q(X, \mathbb{L}(r))$ .
- For a relative scheme  $\pi: X \rightarrow S$  locally of finite type,  $\mathbf{H}^q(S_{\text{ét}}, \pi_! \mathbb{L}(r)_X)$  coincides with the continuous  $p$ -adic cohomology of  $X/S$  with proper support, usually denoted by  $\mathbf{H}_c^q(X, \mathbb{L}(r))$ .

For an object of  $\mathbf{D}^+(X_{\text{ét}}, \mathbb{L})$  of the form  $\Psi \mathbb{L}(r)_Y$  as above, we will simply write

$$\mathbf{H}^q(X, \Psi \mathbb{L}(r)) := \mathbf{H}^q(X_{\text{ét}}, \Psi \mathbb{L}(r)_Y)$$

to be consistent with the common practice, and similarly for cohomology with support. We suppress  $(r)$  in all the notation when  $r = 0$ .

*Remark A.2.* We have

- (1) There is a natural functor  $\mathbf{M}((\text{Spec } K)_{\text{ét}}, L) \rightarrow \mathbf{M}_K(L)$  by regarding a discrete module over the (profinite) group  $\text{Gal}(\bar{K}/K)$  as a module over  $G_K$ . This functor does not commute with projective limits in general.
- (2) For a continuous module  $V$  of the profinite group  $\text{Gal}(\bar{K}/K)$ , we also have the continuous Galois cohomology  $\mathbf{H}_{\text{ct}}^q(K, V)$ , with a canonical map  $\mathbf{H}_{\text{ct}}^q(K, V) \rightarrow \mathbf{H}^q(K, V)$  which is an isomorphism when  $q = 0$  and is injective when  $q = 1$ .
- (3) Suppose that the characteristic of  $K$  is not  $p$ . Let  $X$  be a scheme over  $K$  of finite type and  $\{F_l\}_{l \geq 1}$  a projective system of (torsion) constructible sheaves in  $\mathbf{M}(X_{\text{ét}}, \mathbb{Z}_p)$ . We have a canonical map

$$\mathbf{R}\Gamma\left(X_{\text{ét}}, \mathbf{R} \lim_{\leftarrow l} F_l\right) \rightarrow \mathbf{R}\Phi_K \mathbf{R}\Gamma\left((X_{\bar{K}})_{\text{ét}}, \mathbf{R} \lim_{\leftarrow l} \pi^* F_l\right)$$

in  $\mathbf{D}^+(\mathbb{Z}_p)$ , where  $\pi: X_{\bar{K}} \rightarrow X$  is the natural projection. For every  $q \geq 0$ , it induces a map

$$\mathbf{H}_{\text{ct}}^q(X, (F_l)) \rightarrow \mathbf{H}_K^q\left((X_{\bar{K}})_{\text{ét}}, \mathbf{R} \lim_{\leftarrow l} \pi^* F_l\right)$$

where the left side denotes Jannsen’s continuous étale cohomology. For the source, we have the (continuous) Hochschild–Serre spectral sequence [Jan88, Corollary 3.4]

$$E_{\text{ct}, 2}^{p, q} = \mathbf{H}_{\text{ct}}^p\left(K, \lim_{\leftarrow l} \mathbf{H}^q(X_{\bar{K}}, \pi^* F_l)\right) \Rightarrow \mathbf{H}_{\text{ct}}^{p+q}(X, (F_l));$$

while for the target, we have the (algebraic) Hochschild–Serre spectral sequence

$$E_2^{p, q} = \mathbf{H}^p\left(K, \lim_{\leftarrow l} \mathbf{H}^q(X_{\bar{K}}, \pi^* F_l)\right) \Rightarrow \mathbf{H}_K^{p+q}\left((X_{\bar{K}})_{\text{ét}}, \mathbf{R} \lim_{\leftarrow l} \pi^* F_l\right).$$



It is straightforward to check that the following two diagrams

$$\begin{array}{ccc} \mathbf{H}_{\text{ct}}^q(X, (F_l)) & \longrightarrow & \mathbf{H}_K^q\left((X_{\bar{K}})_{\text{ét}}, \mathbf{R}\lim_{\longleftarrow l} \pi^* F_l\right) \\ \downarrow & & \downarrow \\ \mathbf{H}_{\text{ct}}^0\left(K, \lim_{\longleftarrow l} \mathbf{H}^q(X_{\bar{K}}, \pi^* F_l)\right) & \xlongequal{\quad} & \mathbf{H}^0\left(K, \lim_{\longleftarrow l} \mathbf{H}^q(X_{\bar{K}}, \pi^* F_l)\right) \end{array}$$

and

$$\begin{array}{ccc} \mathbf{H}_{\text{ct}}^q(X, (F_l))^0 & \longrightarrow & \mathbf{H}_K^q\left((X_{\bar{K}})_{\text{ét}}, \mathbf{R}\lim_{\longleftarrow l} \pi^* F_l\right)^0 \\ \downarrow & & \downarrow \\ \mathbf{H}_{\text{ct}}^1\left(K, \lim_{\longleftarrow l} \mathbf{H}^{q-1}(X_{\bar{K}}, \pi^* F_l)\right) & \xrightarrow{\quad} & \mathbf{H}^1\left(K, \lim_{\longleftarrow l} \mathbf{H}^{q-1}(X_{\bar{K}}, \pi^* F_l)\right) \end{array}$$

commute. Here, in the second diagram,  $\mathbf{H}_{\text{ct}}^q(X, (F_l))^0$  denotes the inverse image of  $\mathbf{H}_K^q\left((X_{\bar{K}})_{\text{ét}}, \mathbf{R}\lim_{\longleftarrow l} \pi^* F_l\right)^0$ , and the two vertical arrows are the corresponding edge maps.

**Lemma A.3.** *Let  $L$  be a ring and  $S$  a site with an action of  $G_K$ . Consider a distinguished triangle*

$$\mathcal{C}_0 \xrightarrow{f_0} \mathcal{C}_1 \xrightarrow{f_1} \mathcal{C}_2 \xrightarrow{+1}$$

in  $\mathbf{D}_K^+(S, L)$  and an integer  $q$ , inducing the following commutative diagram

$$\begin{array}{ccccccc} \mathbf{H}_K^{q-1}(S, \mathcal{C}_2) & \xrightarrow{\delta} & \mathbf{H}_K^q(S, \mathcal{C}_0) & \xrightarrow{f_0} & \mathbf{H}_K^q(S, \mathcal{C}_1) & \xrightarrow{f_1} & \mathbf{H}_K^q(S, \mathcal{C}_2) \\ r_2^- \downarrow & & \downarrow r_0 & & \downarrow r_1 & & \downarrow r_2 \\ \mathbf{H}^{q-1}(S, \mathcal{C}_2) & \xrightarrow{\bar{\delta}} & \mathbf{H}^q(S, \mathcal{C}_0) & \xrightarrow{\bar{f}_0} & \mathbf{H}^q(S, \mathcal{C}_1) & \xrightarrow{\bar{f}_1} & \mathbf{H}^q(S, \mathcal{C}_2) \end{array}$$

in which all vertical arrows are restriction maps. Then for every element  $\gamma_1 \in \mathbf{H}_K^q(S, \mathcal{C}_1)$  satisfying  $r_2(f_1(\gamma_1)) = 0$ , the image of  $f_1(\gamma_1)$  under the composite map

$$\mathbf{H}_K^q(S, \mathcal{C}_2)^0 \rightarrow \mathbf{H}^1(K, \mathbf{H}^{q-1}(S, \mathcal{C}_2)) \rightarrow \mathbf{H}^1\left(K, \frac{\mathbf{H}^{q-1}(S, \mathcal{C}_2)}{\mathbf{H}^{q-1}(S, \mathcal{C}_1)}\right)$$

can be represented by the 1-cocycle  $g \mapsto g\tilde{\gamma}_0 - \tilde{\gamma}_0$  for  $g \in G_K$ , where  $\tilde{\gamma}_0$  is an arbitrary element in  $\mathbf{H}^q(S, \mathcal{C}_0)$  satisfying  $\bar{f}_0(\tilde{\gamma}_0) = r_1(\gamma_1)$ .

*Proof.* It is easy to check that the 1-cocycle does not depend on the choice of  $\tilde{\gamma}_0$ . Thus, it suffices to check the statement for one such element.

Take an injective resolution  $0 \rightarrow I_0^\bullet \xrightarrow{i_0} I_1^\bullet \xrightarrow{i_1} I_2^\bullet \rightarrow 0$  in  $\mathbf{C}_K^+(L)$  of the exact triangle

$$\mathbf{R}\Gamma(S, \mathcal{C}_0) \rightarrow \mathbf{R}\Gamma(S, \mathcal{C}_1) \rightarrow \mathbf{R}\Gamma(S, \mathcal{C}_2) \xrightarrow{+1} .$$

Then the diagram in the statement can be replaced by

$$\begin{array}{ccccccc} \mathbf{H}^{q-1}((I_2^\bullet)^{G_K}) & \xrightarrow{\delta} & \mathbf{H}^q((I_0^\bullet)^{G_K}) & \xrightarrow{i_0^q} & \mathbf{H}^q((I_1^\bullet)^{G_K}) & \xrightarrow{i_1^q} & \mathbf{H}^q((I_2^\bullet)^{G_K}) \\ r_2^- \downarrow & & \downarrow r_0 & & \downarrow r_1 & & \downarrow r_2 \\ \mathbf{H}^{q-1}(I_2^\bullet) & \xrightarrow{\bar{\delta}} & \mathbf{H}^q(I_0^\bullet) & \xrightarrow{i_0^q} & \mathbf{H}^q(I_1^\bullet) & \xrightarrow{i_1^q} & \mathbf{H}^q(I_2^\bullet) \end{array}$$

in which all vertical arrows are induced by natural inclusions. We have

$$\mathbf{H}_K^q(S, \mathcal{C}_2)^0 = \mathbf{H}^q((I_2^\bullet)^{G_K})^0 = \frac{\text{Ker}\left((I_2^q)^{G_K} \xrightarrow{d} (I_2^{q+1})^{G_K}\right) \cap \text{Im}\left(I_2^{q-1} \xrightarrow{d} I_2^q\right)}{\text{Im}\left((I_2^{q-1})^{G_K} \xrightarrow{d} (I_2^q)^{G_K}\right)}.$$

For  $\gamma_2 \in \mathbf{H}^q((I_2^\bullet)^{G_K})^0$ , its image in  $\mathbf{H}^1(K, \mathbf{H}^{q-1}(I_2^\bullet))$  under the edge map can be represented by the 1-cocycle that sends  $g \in G_K$  to the (cohomology class) of  $g\beta_2^{q-1} - \beta_2^{q-1}$ , where  $\beta_2^{q-1} \in I_2^{q-1}$  is an arbitrary element whose differential  $d\beta_2^{q-1}$  in  $I_2^q$  represents  $\gamma_2$  (so that  $g\beta_2^{q-1} - \beta_2^{q-1}$  is closed).

Let  $\alpha_1^q \in (I_1^q)^{G_K}$  be a closed element that represents  $\gamma_1 \in \mathbf{H}^d((I_1^\bullet)^{G_K})$ . Since  $i_1^q(\alpha_1^q)$  belongs to  $\mathbf{H}^q((I_2^\bullet)^{G_K})^0$ , we may choose an element  $\beta_2^{q-1} \in I_2^{q-1}$  whose differential  $d\beta_2^{q-1}$  in  $I_2^q$  equals  $i_1^q(\alpha_1^q)$ . To construct an element  $\tilde{\gamma}_0$  that lifts  $r_1(\gamma_1)$ , we take an element  $\beta_1^{q-1} \in I_1^{q-1}$  such that  $i_1^{q-1}(\beta_1^{q-1}) = \beta_2^{q-1}$ . Then  $i_1^q(\alpha_1^q - d\beta_1^{q-1}) = 0$ , which implies that  $\beta_0^q := \alpha_1^q - d\beta_1^{q-1}$  is a closed element of  $I_0^q$ . Thus, we may take  $\tilde{\gamma}_0 \in \mathbf{H}^q(I_0^\bullet)$  to be the cohomology class of  $\beta_0^q$ . Since  $\alpha_1^q$  is fixed by  $G_K$ , we have  $g\beta_0^q - \beta_0^q = gd\beta_1^{q-1} - d\beta_1^{q-1} = d(g\beta_1^{q-1} - \beta_1^{q-1})$ , which is a closed element of  $I_0^q$ . However,  $d(g\beta_1^{q-1} - \beta_1^{q-1})$  exactly represents the class of  $g\beta_2^{q-1} - \beta_2^{q-1}$  under the coboundary map  $\bar{d}: \mathbf{H}^{q-1}(I_2^\bullet) \rightarrow \mathbf{H}^q(I_0^\bullet)$ . The lemma then follows.  $\square$

**A.3. Bi-extensions of cycles.** Let  $K$  be a field of characteristic different from  $p$  with a fixed algebraic closure  $\bar{K}$ . Let  $X$  be a projective smooth scheme over  $K$  of pure dimension  $n - 1$ . For every integer  $d$ , put

$$Z^d(X)_{\mathbb{L}}^0 := \text{Ker}\left(Z^d(X)_{\mathbb{L}} \rightarrow \mathbf{H}^{2d}(X_{\bar{K}}, \mathbb{L}(d))\right).$$

Now we consider two elements  $c \in Z^d(X)_{\mathbb{L}}^0$  and  $c' \in Z^{d'}(X)_{\mathbb{L}}^0$  with  $d + d' = n$ , such that  $c$  and  $c'$  have disjoint supports. Choose disjoint nonempty closed subsets  $Z$  and  $Z'$  of  $X$  of pure codimension  $d$  and  $d'$  containing the supports of  $c$  and  $c'$ , respectively. Denote by  $j: Z \rightarrow X$  and  $j': Z' \rightarrow X$  the closed immersions and put  $U := X \setminus Z$  and  $U' := X \setminus Z'$ . We have the following diagram

$$\begin{array}{ccc} U \cap U' & \xrightarrow{i'} & U \\ \downarrow i & & \downarrow i \\ U' & \xrightarrow{i'} & X \end{array}$$

of open immersions. We have the following induced diagram

$$(A.1) \quad \begin{array}{ccccccc} & & & 0 & & 0 & \\ & & & \downarrow & & \downarrow & \\ \mathbf{H}^{2d-2}(X_{\bar{K}}, \mathbb{L}(d)) & \longrightarrow & \mathbf{H}^{2d-2}(Z'_{\bar{K}}, \mathbb{L}(d)) & \longrightarrow & \mathbf{H}^{2d-1}(X_{\bar{K}}, i'_1 \mathbb{L}(d)) & \longrightarrow & \mathbf{H}^{2d-1}(X_{\bar{K}}, \mathbb{L}(d)) \longrightarrow 0 \\ \downarrow \sim & & \parallel & & \downarrow & & \downarrow \\ \mathbf{H}^{2d-2}(U_{\bar{K}}, \mathbb{L}(d)) & \longrightarrow & \mathbf{H}^{2d-2}(Z'_{\bar{K}}, \mathbb{L}(d)) & \longrightarrow & \mathbf{H}^{2d-1}(U_{\bar{K}}, i'_1 \mathbb{L}(d)) & \longrightarrow & \mathbf{H}^{2d-1}(U_{\bar{K}}, \mathbb{L}(d)) \longrightarrow 0 \\ & & & & \downarrow & & \downarrow \\ & & & & \mathbf{H}^{2d}(Z_{\bar{K}}, j^1 \mathbb{L}(d)) & \xlongequal{\quad} & \mathbf{H}^{2d}(Z_{\bar{K}}, j^1 \mathbb{L}(d)) \\ & & & & \downarrow & & \downarrow \\ & & & & \mathbf{H}^{2d}(X_{\bar{K}}, i'_1 \mathbb{L}(d)) & \xrightarrow{\sim} & \mathbf{H}^{2d}(X_{\bar{K}}, \mathbb{L}(d)) \end{array}$$

in  $\mathbf{M}_K(\mathbb{L})$ .

The element  $c$  gives rise to a map  $\kappa^c: \mathbb{L} \rightarrow \mathbf{H}^{2d}(Z_{\bar{K}}, j^1 \mathbb{L}(d))$  whose image is contained in the kernel of the map  $\mathbf{H}^{2d}(Z_{\bar{K}}, j^1 \mathbb{L}(d)) \rightarrow \mathbf{H}^{2d}(X_{\bar{K}}, \mathbb{L}(d))$ . The element  $c'$  gives rise to a map  $\kappa_{c'}: \mathbf{H}^{2d-2}(Z'_{\bar{K}}, \mathbb{L}(d-1)) \rightarrow \mathbb{L}$  that vanishes

on the image of the map  $H^{2d-2}(X_{\overline{K}}, \mathbb{L}(d)) \rightarrow H^{2d-2}(Z'_{\overline{K}}, \mathbb{L}(d))$ . Applying the pullback along  $\kappa^c$  and the pushforward along  $\kappa_{c'}(1)$  to the diagram (A.1), we obtain the following bi-extension diagram

$$\begin{array}{ccccccc}
& & & 0 & & 0 & \\
& & & \downarrow & & \downarrow & \\
0 & \longrightarrow & \mathbb{L}(1) & \longrightarrow & E_{c'} & \longrightarrow & H^{2d-1}(X_{\overline{K}}, \mathbb{L}(d)) \longrightarrow 0 \\
& & \parallel & & \downarrow & & \downarrow \\
0 & \longrightarrow & \mathbb{L}(1) & \longrightarrow & E_{c'}^c & \longrightarrow & E^c \longrightarrow 0 \\
& & & & \downarrow & & \downarrow \\
& & & & \mathbb{L} & \xlongequal{\quad\quad\quad} & \mathbb{L} \\
& & & & \downarrow & & \downarrow \\
& & & & 0 & & 0
\end{array}$$

in  $\mathbf{M}_K(\mathbb{L})$ . It is easy to see that the above diagram does not depend on the choices of  $Z$  and  $Z'$ . We have three induced extension classes

- $[E^c] \in H_{\text{ct}}^1(K, H^{2d-1}(X_{\overline{K}}, \mathbb{L}(d)))$ ,
- $[E_{c'}^c] = [E_{c'}] \in H_{\text{ct}}^1(K, H^{2d-1}(X_{\overline{K}}, \mathbb{L}(d)))$ ,
- $[E_{c'}^c] \in H_{\text{ct}}^1(K, E_{c'})$ .

**A.4. Relation with Beilinson's local index.** In this subsection, we assume that  $K$  is a non-archimedean local field whose residue field has characteristic different from  $p$ . We study the relation between Nekovář's local  $p$ -adic height and Beilinson's local index.

Assume that the cycle classes of  $c$  and  $c'$  in  $H^{2d}(X, \mathbb{L}(d))$  and  $H^{2d'}(X, \mathbb{L}(d'))$  vanish, respectively.<sup>15</sup> Then the image of  $[E_{c'}^c]$  in  $H_{\text{ct}}^1(K, H^{2d-1}(X_{\overline{K}}, \mathbb{L}(d)))$  vanishes, hence  $[E_{c'}^c]$  belongs to the image of the map  $H_{\text{ct}}^1(K, \mathbb{L}(1)) \rightarrow H_{\text{ct}}^1(K, E_{c'})$  which is injective. Following Nekovář [Nek93], we denote by  $\langle c, c' \rangle_{X, K}^N$  the image of  $[E_{c'}^c]$  under the natural isomorphism  $H_{\text{ct}}^1(K, \mathbb{L}(1)) = \widehat{K}^\times \otimes_{\mathbb{Z}} \mathbb{L}$  given by the Kummer maps.

**Theorem A.4.** *Let the situation be as above. Then*

$$\langle c, c' \rangle_{X, K}^N = \langle c, c' \rangle_{X, K}^B,$$

where  $\langle c, c' \rangle_K^B$  denotes Beilinson's local index which we recall below.

We recall the definition of Beilinson's local index [Bei87, Section 2] (see also [LL21, Appendix B]). We have the refined cycle class  $[c] \in H_Z^{2d}(X, \mathbb{L}(d)) = H^{2d}(Z, j^! \mathbb{L}(d))$ , which is contained in the kernel of the map  $H_Z^{2d}(X, \mathbb{L}(d)) \rightarrow H^{2d}(X, \mathbb{L}(d))$ , hence we may choose an element  $\gamma \in H^{2d-1}(U, \mathbb{L}(d))$  that maps to  $[c]$  under the coboundary map  $H^{2d-1}(U, \mathbb{L}(d)) \rightarrow H_Z^{2d}(X, \mathbb{L}(d))$ . Similarly, we can choose an element  $\gamma' \in H^{2d'-1}(U', \mathbb{L}(d'))$  for  $c'$ . Beilinson's local index, which we denote by  $\langle c, c' \rangle_{X, K}^B$ , is defined as the image of  $\gamma \cup \gamma'$  under the composite map

$$H^{2n-2}(U \cap U', \mathbb{L}(n)) \rightarrow H^{2n-1}(X, \mathbb{L}(n)) \xrightarrow{\text{Tr}_{X/K}} H^1(\text{Spec } K, \mathbb{L}(1)) = H_{\text{ct}}^1(K, \mathbb{L}(1)) = \widehat{K}^\times \otimes_{\mathbb{Z}} \mathbb{L},$$

in which the first map is the coboundary map in the Mayer–Vietoris exact sequence for the open covering  $X = U \cup U'$ .

*Remark A.5.* In fact, in [Bei87, Section 2] and [LL21, Appendix B], the local index  $\langle c, c' \rangle_K^B$  takes value in  $\mathbb{L}$  via the canonical isomorphism  $H^1(\text{Spec } K, \mathbb{Q}_p(1)) \simeq \mathbb{Q}_p$  that is the composition

$$H^1(\text{Spec } K, \mathbb{Q}_p(1)) \rightarrow H_{\text{Spec } \kappa}^2(\text{Spec } O_K, \mathbb{Q}_p(1)) \xrightarrow{\sim} H^0(\text{Spec } \kappa, \mathbb{Q}_p) \simeq \mathbb{Q}_p$$

in which  $\kappa$  is the residue field of  $K$ . By [GD77, 2.1.3] or [Nek95, II.(2.16.1)], the induced isomorphism  $\widehat{K}^\times \otimes_{\mathbb{Z}} \mathbb{Q}_p \rightarrow \mathbb{Q}_p$  sends a uniformizer of  $K$  to  $-1$ , rather than 1.

<sup>15</sup>This is automatic if the monodromy-weight conjecture holds for  $H^{2d-1}(X_{\overline{K}}, \mathbb{L}(d))$ .

*Proof of Theorem A.4.* Denote the image of  $\gamma$  under the restriction map  $H^{2d-1}(U, \mathbb{L}(d)) \rightarrow H^{2d-1}(U_{\bar{K}}, \mathbb{L}(d))$  by  $\bar{\gamma}$ , which belongs to  $E^c$ . Take a preimage  $\tilde{\gamma} \in E_{c'}^c$  of  $\bar{\gamma}$ . Then  $[E_{c'}^c]$ , as an element of  $H_{\text{ct}}^1(K, \mathbb{L}(1))$ , is represented by the 1-cocycle

$$\phi_{\text{N}}: g \mapsto \text{Tr}_{U'_{\bar{K}}/\bar{K}}((g\tilde{\gamma} - \tilde{\gamma}) \cup \tilde{\gamma}'), \quad g \in G_K,$$

where  $\tilde{\gamma}'$  is the image of  $\gamma'$  under the map  $H^{2d'-1}(U', \mathbb{L}(d')) \rightarrow H^{2d'-1}(U'_{\bar{K}}, \mathbb{L}(d'))$ . Here, we note that for  $g \in G_K$ ,  $g\tilde{\gamma} - \tilde{\gamma}$  belongs to  $E_{c'} \subseteq H^{2d-1}(X_{\bar{K}}, \mathbb{L}(d)) = H_{\text{c}}^{2d-1}(U'_{\bar{K}}, \mathbb{L}(d))$ .

To find a 1-cocycle representing  $\langle c, c' \rangle_{X,K}^{\text{B}}$  as an element in  $H_{\text{ct}}^1(K, \mathbb{L}(1))$ , we use the refined cycle class  $[c'] \in H_{\text{Z}}^{2d'}(X, \mathbb{L}(d')) = H^{2d'}(Z', j'^! \mathbb{L}(d'))$ . It follows easily that  $\langle c, c' \rangle_{X,K}^{\text{B}}$  coincides with the image of  $\gamma \cup [c'] \in H^{2n-1}(Z', j'^! \mathbb{L}(n))$  under the composite map

$$H^{2n-1}(Z', j'^! \mathbb{L}(n)) \rightarrow H^{2n-1}(X, \mathbb{L}(n)) \xrightarrow{\text{Tr}_{X/K}} H^1(\text{Spec } K, \mathbb{L}(1)).$$

However, the map

$$H^{2d-1}(U, \mathbb{L}(d)) \times H^{2d'}(Z', j'^! \mathbb{L}(d')) \xrightarrow{\cup} H^{2n-1}(Z', j'^! \mathbb{L}(n)) \rightarrow H^{2n-1}(X, \mathbb{L}(n))$$

can also be computed as the composite map

$$H^{2d-1}(U, \mathbb{L}(d)) \times H^{2d'}(Z', j'^! \mathbb{L}(d')) \rightarrow H^{2d-1}(Z', \mathbb{L}(d)) \times H^{2d'}(Z', j'^! \mathbb{L}(d')) \xrightarrow{\cup} H^{2n-1}(X, \mathbb{L}(n)).$$

Since  $H^{2n-1}(X_{\bar{K}}, \mathbb{L}(n)) = 0$ , the (continuous) Hochschild–Serre spectral sequence induces an isomorphism  $H^{2n-1}(X, \mathbb{L}(n)) \simeq H_{\text{ct}}^1(K, H^{2n-2}(X_{\bar{K}}, \mathbb{L}(n)))$ , under which the trace map  $\text{Tr}_{X/K}$  coincides with  $H_{\text{ct}}^1(K, \text{Tr}_{X_{\bar{K}}/\bar{K}})$ . Since  $H^{2d-1}(Z'_{\bar{K}}, \mathbb{L}(d)) = 0$ , we have the following commutative diagram

$$\begin{array}{ccc} H^{2d-1}(Z', \mathbb{L}(d)) \times H^{2d'}(Z', j'^! \mathbb{L}(d')) & \xrightarrow{\cup} & H^{2n-1}(X, \mathbb{L}(n)) \\ \downarrow & & \downarrow \simeq \\ H_{\text{ct}}^1(K, H^{2d-2}(Z'_{\bar{K}}, \mathbb{L}(d))) \times H^{2d'}(Z'_{\bar{K}}, j'^! \mathbb{L}(d')) & \xrightarrow{\cup} & H_{\text{ct}}^1(K, H^{2n-2}(X_{\bar{K}}, \mathbb{L}(n))) \end{array}$$

of  $\mathbb{L}$ -vector spaces. Suppose that the image of  $\gamma$  under the composite map

$$H^{2d-1}(U, \mathbb{L}(d)) \rightarrow H^{2d-1}(Z', \mathbb{L}(d)) \rightarrow H_{\text{ct}}^1(K, H^{2d-2}(Z'_{\bar{K}}, \mathbb{L}(d)))$$

is represented by a 1-cocycle  $\phi$ . Then from the above discussion,  $\langle c, c' \rangle_{X,K}^{\text{B}}$  is represented by the 1-cocycle

$$\phi_{\text{B}}: g \mapsto \text{Tr}_{X_{\bar{K}}/\bar{K}}(\phi(g) \cup [c']), \quad g \in G_K,$$

where we have regard  $[c']$  its image in  $H^{2d'}(Z'_{\bar{K}}, j'^! \mathbb{L}(d'))$  by abuse of notation.

Note that in the formula for  $\phi_{\text{N}}(g)$ ,  $g\tilde{\gamma} - \tilde{\gamma}$  in fact belongs to  $\mathbb{L}(1)$ , hence  $(g\tilde{\gamma} - \tilde{\gamma}) \cup \tilde{\gamma}' = (g\tilde{\gamma} - \tilde{\gamma}) \cup [c']$ . As the map  $\cup [c']: H^{2d-2}(Z'_{\bar{K}}, \mathbb{L}(d)) \rightarrow \mathbb{L}(1)$  factors through the image of  $H^{2d-2}(U_{\bar{K}}, \mathbb{L}(d)) \rightarrow H^{2d-2}(Z'_{\bar{K}}, \mathbb{L}(d))$ , the proof of Theorem A.4 reduces to the claim that the image of  $\gamma$  under the composite map

$$H^{2d-1}(U, \mathbb{L}(d)) \rightarrow H^{2d-1}(Z', \mathbb{L}(d)) \rightarrow H_{\text{ct}}^1(K, H^{2d-2}(Z'_{\bar{K}}, \mathbb{L}(d))) \rightarrow H_{\text{ct}}^1\left(K, \frac{H^{2d-2}(Z'_{\bar{K}}, \mathbb{L}(d))}{H^{2d-2}(U_{\bar{K}}, \mathbb{L}(d))}\right)$$

can be represented by the 1-cocycle  $g \mapsto g\tilde{\gamma} - \tilde{\gamma}$  for  $g \in G_K$ . Now in view of Remark A.2(3), the proposition follows from Lemma A.3 for  $S = (U_{\bar{K}})_{\text{ét}}$  with the obvious action of  $G_K$ , the distinguished triangle

$$(i'_{\bar{K}})_! \mathbb{L}(d)_{(U \setminus Z')_{\bar{K}}} \rightarrow \mathbb{L}(d)_{U_{\bar{K}}} \rightarrow (j'_{\bar{K}})_* \mathbb{L}(d)_{Z'_{\bar{K}}} \xrightarrow{+1}$$

(with  $j': Z' \rightarrow U$ ), and  $q = 2d - 1$ . □

**A.5. Crystalline property of bi-extensions.** In this subsection, we assume that  $K$  is a finite extension of  $\mathbb{Q}_p$  with residue field  $\kappa$ . Denote by  $W$  the Witt ring of  $\kappa$ , and by  $K_0$  the fraction field of  $W$ , which is canonically a subfield of  $K$ .

We assume that  $X$  admits a proper strictly semistable model  $\mathcal{X}$  over  $O_K$ . Put  $X := \mathcal{X} \otimes_{O_K} \kappa$ . For every integer  $h \geq 1$ , denote by  $X^{(h)}$  the disjoint union of intersections of  $h$  different irreducible components of  $X$ , which is either empty or a proper smooth scheme over  $\kappa$ .

**Theorem A.6.** *Suppose that  $n < p$ . Let  $\mathbb{T}$  be an  $\mathbb{L}$ -ring of étale correspondences on  $\mathcal{X}$  and  $\mathfrak{m}, \mathfrak{m}'$  two maximal ideals of  $\mathbb{T}$  satisfying that*

$$(A.2) \quad \bigoplus_{h>1, q \geq 0} \left( H_{\text{cris}}^q(X^{(h)}/W) \otimes_{\mathbb{Z}_p} \mathbb{L} \right)_{\mathfrak{m}} = \bigoplus_{h>1, q \geq 0} \left( H_{\text{cris}}^q(X^{(h)}/W) \otimes_{\mathbb{Z}_p} \mathbb{L} \right)_{\mathfrak{m}'} = 0.$$

*Then there exist elements  $t \in \mathbb{T} \setminus \mathfrak{m}$  and  $t' \in \mathbb{T} \setminus \mathfrak{m}'$  depending on  $\mathcal{X}$  only such that the following holds: For two arbitrary elements  $c \in Z^d(X)_{\mathbb{L}}^0$  and  $c' \in Z^{d'}(X)_{\mathbb{L}}^0$  with  $d + d' = n$  satisfying that*

- (1)  $(\text{supp } C)^t \cap (\text{supp } C')^{t'} = \emptyset$  for every  $t, t' \in \mathbb{T}$ , where  $C$  and  $C'$  denote the Zariski closures of  $c$  and  $c'$  in  $\mathcal{X}$ , respectively,
- (2) the codimension of  $\text{supp } C'$  in  $X^{(h)}$  is at least  $d'$  for every  $h \geq 1$ ,

*we have*

- $[E^{t^*c}] \in H_f^1(K, H^{2d-1}(X_{\overline{K}}, \mathbb{L}(d)))$ ,
- $[E_{t'^*c'}^{\vee}(1)] \in H_f^1(K, H^{2d'-1}(X_{\overline{K}}, \mathbb{L}(d')))$ ,
- $[E_{t'^*c'}^{t^*c}] \in H_f^1(K, E_{t'^*c'})$ ,

*simultaneously. Here, the bi-extension  $E_{t'^*c'}^{t^*c}$  exists by (1).*

**Remark A.7.** By taking  $\mathbb{T} = \mathbb{L}$ , Theorem A.6 asserts that  $E_{c'}^c$  is crystalline as long as  $\text{supp } C \cap \text{supp } C' = \emptyset$  when  $\mathcal{X}$  is a proper smooth model of  $X$  over  $O_K$ .<sup>16</sup> This confirms the (equivalent) conjecture in the remark after [Shn16, Theorem 8.7] when  $n < p$ .<sup>17</sup>

Our main strategy of proving Theorem A.6 is similar to [Sat13]. The main challenge is to show that the bi-extension  $[E_{t'^*c'}^{t^*c}]$  is a crystalline class. We consider the Abel–Jacobi map from the homologically trivial part of the degree  $2d$  syntomic cohomology of  $\mathcal{X} \setminus (\text{supp } C')$  with proper support to  $H_{\text{ct}}^1(K, H_c^{2d-1}(U'_{\overline{K}}, \mathbb{Q}_p(d)))$ . After we show that  $E_{t'^*c'}$  is crystalline for suitable  $t'$ , it suffices to show that if a syntomic class comes from a (homologically trivial) cycle, then its Abel–Jacobi image vanishes in  $H^1(K, H_c^{2d-1}(U'_{\overline{K}}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\text{cris}})$  “after localization at  $\mathfrak{m}'$ ” (and replace  $K$  by a finite extension in fact), under the conditions in Theorem A.6. However, the main challenge for us is that unlike the situation in [Sat13] where  $U' = X$ , we do not have a comparison theorem for  $H_c^{2d-1}(U'_{\overline{K}}, \mathbb{Q}_p)$  with some cohomology on the special fiber in general. Also, due to the constraint of the conditions in the theorem, we can not reduce the theorem to an alteration. We solve this problem in the following way. First, we show that the kernel of the Abel–Jacobi map from the above syntomic cohomology to  $H^1(K, H_c^{2d-1}(U'_{\overline{K}}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\text{cris}})$  is contained in the kernel of another map to a certain space defined by log rigid cohomology (Proposition B.10). Second, we show that a cycle class will vanish in this space “after localization at  $\mathfrak{m}'$ ”. For the first step, which shall hold more generally without the conditions in the theorem, we pass to a strict semistable alteration.

The full proof of Theorem A.6 will occupy Appendix B.

**A.6. Recollection on  $p$ -adic Galois representation.** Let  $V$  be a finite-dimensional continuous representation of  $\text{Gal}(\overline{K}/K)$  with coefficients in  $\mathbb{L}$ , which we assume to be *de Rham*, that is

$$\mathbb{D}_{\text{dR}}(V) := \left( V \otimes_{\mathbb{Q}_p} \mathbb{B}_{\text{dR}} \right)^{\text{Gal}(\overline{K}/K)}$$

is a free  $\mathbb{L} \otimes_{\mathbb{Q}_p} K$ -module of rank  $\dim_{\mathbb{L}} V$ . We have a decreasing filtration  $F^i \mathbb{D}_{\text{dR}}(V)$  of  $\mathbb{L} \otimes_{\mathbb{Q}_p} K$ -submodules of  $\mathbb{D}_{\text{dR}}(V)$ , known as the de Rham filtration. Moreover, for every embedding  $\tau: K \rightarrow \overline{\mathbb{Q}_p}$ , Fontaine constructed a Weil–Deligne representation  $\text{WD}(V)_{\tau}$  of  $K$  with coefficients in  $\overline{\mathbb{Q}_p}$ , with underlying  $\overline{\mathbb{Q}_p}$ -vector space

<sup>16</sup>We warn the readers that this assertion is wrong if one replaces the word *smooth* by *strictly semistable*.

<sup>17</sup>However, our strategy for the proof of Theorem A.6 is different from the case of (local systems over) curves in [Shn16, Theorem 8.7].

$\mathbb{D}_{\text{dR}}(V) \otimes_{\mathbb{L} \otimes_{\mathbb{Q}_p, K, 1 \otimes \tau} \overline{\mathbb{Q}_p}}$ . The isomorphism class of  $\text{WD}(V)_\tau$  is independent of  $\tau$ . See, for example, [TY07, Section 1] for more details.

**Definition A.8.** Let  $\mu$  be a real number. We say that  $V$  is *pure of weight  $\mu$*  if for some (hence every)  $\tau: K \rightarrow \overline{\mathbb{Q}_p}$ , all geometric Frobenius eigenvalues of  $\text{gr}_i \text{WD}(V)_\tau$  are Weil  $|\kappa|^{\mu+i}$ -numbers for every  $i \in \mathbb{Z}$ , where  $\text{gr}_i \text{WD}(V)_\tau$  denotes the  $i$ -th graded piece of the monodromy filtration on  $\text{WD}(V)_\tau$ .

**Definition A.9.** We denote by  $\mathbb{D}_{\text{dR}}^-(V)$  the maximal  $\mathbb{L} \otimes_{\mathbb{Q}_p} K$ -submodule of  $\mathbb{D}_{\text{dR}}(V)$  such that for every  $\tau: K \rightarrow \overline{\mathbb{Q}_p}$ , the  $\overline{\mathbb{Q}_p}$ -subspace  $\mathbb{D}_{\text{dR}}^-(V) \otimes_{\mathbb{L} \otimes_{\mathbb{Q}_p, K, 1 \otimes \tau} \overline{\mathbb{Q}_p}} \subseteq \text{WD}(V)_\tau$  is stable under the action of the Weil group of  $K$  and that none of its geometric Frobenius eigenvalues belongs to  $\overline{\mathbb{Z}_p}$ .

We say that  $V$  satisfies the *Panchishkin condition* if the natural map

$$\mathbb{D}_{\text{dR}}^-(V) \oplus F^0 \mathbb{D}_{\text{dR}}(V) \rightarrow \mathbb{D}_{\text{dR}}(V)$$

is an isomorphism.

**Lemma A.10.** *Let  $\mu$  be a nonzero integer and  $0 \rightarrow V_1 \rightarrow V \rightarrow V_2 \rightarrow 0$  a short exact sequence of crystalline representations of  $\text{Gal}(\overline{K}/K)$ . If  $V_1$  and  $V_2$  are pure of weight  $\mu$  and satisfy the Panchishkin condition, then so does  $V$ .*

*Proof.* By definition, the induced sequence  $0 \rightarrow \mathbb{D}_{\text{dR}}^-(V_1) \rightarrow \mathbb{D}_{\text{dR}}^-(V) \rightarrow \mathbb{D}_{\text{dR}}^-(V_2) \rightarrow 0$  is exact. Thus it suffices to show that the induced sequence

$$(A.3) \quad 0 \rightarrow \mathbb{D}_{\text{dR}}(V_1)/F^0 \mathbb{D}_{\text{dR}}(V_1) \rightarrow \mathbb{D}_{\text{dR}}(V)/F^0 \mathbb{D}_{\text{dR}}(V) \rightarrow \mathbb{D}_{\text{dR}}(V_2)/F^0 \mathbb{D}_{\text{dR}}(V_2) \rightarrow 0$$

is also exact. Since for  $? \in \{1, 2, \emptyset\}$ ,  $V_?$  is pure of weight  $\mu$  which is nonzero, by [Nek93, Theorem 1.15 & Corollary 1.16], the exponential map  $\mathbb{D}_{\text{dR}}(V_?)/F^0 \mathbb{D}_{\text{dR}}(V_?) \rightarrow H_f^1(K, V_?)$  is an isomorphism. Thus, the exactness of (A.3) follows from the exactness of the following short sequence

$$0 \rightarrow H_f^1(K, V_1) \rightarrow H_f^1(K, V) \rightarrow H_f^1(K, V_2) \rightarrow 0$$

guaranteed by [Nek93, Proposition 1.25].  $\square$

**A.7. Decomposition of  $p$ -adic height pairing.** In this subsection, we take  $K$  to be a number field. Let  $X$  be a proper smooth scheme over  $K$  of pure dimension  $n - 1$  and take two positive integers  $d, d'$  satisfying  $d + d' = n$ .

Consider  $\mathbb{L}[G_K]$ -submodules  $V$  and  $V'$  of  $H^{2d-1}(X_{\overline{K}}, \mathbb{L}(d))$  and  $H^{2d'-1}(X_{\overline{K}}, \mathbb{L}(d'))$ , respectively, satisfying

(V1) The Poincaré duality on  $X_{\overline{K}}$  induces an isomorphism  $V' \simeq V^\vee(1)$ .

(V2) For every nonarchimedean place  $u$  of  $K$  not above  $p$ ,  $H^i(K_u, V) = H^i(K_u, V') = 0$  for  $i \in \mathbb{Z}$ .

(V3) For every place  $u$  of  $K$  above  $p$ ,  $V|_{K_u}$  (hence  $V'|_{K_u}$ ) is semistable and pure of weight  $-1$  (Definition A.8).

(V4) For every place  $u$  of  $K$  above  $p$ ,  $V|_{K_u}$  satisfies the Panchishkin condition (Definition A.9).

We have the canonical  $p$ -adic height pairing

$$\langle \cdot, \cdot \rangle_{(V, V'), K}: H_f^1(K, V) \times H_f^1(K, V') \rightarrow \Gamma_{K, p} \otimes_{\mathbb{Z}_p} \mathbb{L}$$

constructed in [Nek93], using the Hodge splitting map induced from the decomposition in Definition A.9 for places above  $p$ . The pairing is  $\mathbb{L}$ -bilinear.

Recall that we have the Abel–Jacobi map

$$\text{AJ}: Z^d(X)_{\mathbb{L}}^0 \rightarrow H_{\text{ct}}^1(K, H^{2d-1}(X_{\overline{K}}, \mathbb{L}(d))).$$

We denote by  $Z_{\mathbb{V}}^d(X)_{\mathbb{L}}^0$  the subspace of  $Z^d(X)_{\mathbb{L}}^0$  that is the inverse image of  $H_{\text{ct}}^1(K, V)$ . By [Nek00, Theorem 3.1], the image of  $\text{AJ}$  is contained in  $H_{\text{st}}^1(K, H^{2d-1}(X_{\overline{K}}, \mathbb{L}(d)))$ . Moreover, (V2) implies that  $H_{\text{st}}^1(K, V) = H_f^1(K, V)$ . Thus, we have the Abel–Jacobi map

$$\text{AJ}: Z_{\mathbb{V}}^d(X)_{\mathbb{L}}^0 \rightarrow H_f^1(K, V).$$

Similarly, we have  $Z_{\mathbb{V}'}^{d'}(X)_{\mathbb{L}}^0$  and the corresponding Abel–Jacobi map

$$\text{AJ}: Z_{\mathbb{V}'}^{d'}(X)_{\mathbb{L}}^0 \rightarrow H_f^1(K, V').$$

Combining with the two Abel–Jacobi maps, we obtain a pairing

$$(A.4) \quad \langle \cdot, \cdot \rangle_{(V, V'), K}: Z_{\mathbb{V}}^d(X)_{\mathbb{L}}^0 \times Z_{\mathbb{V}'}^{d'}(X)_{\mathbb{L}}^0 \rightarrow \Gamma_{K, p} \otimes_{\mathbb{Z}_p} \mathbb{L}.$$

Take two elements  $c \in Z_{\mathbb{V}}^d(X)_{\mathbb{L}}^0$  and  $c' \in Z_{\mathbb{V}'}^d(X)_{\mathbb{L}}^0$  with disjoint supports. Then according to [Nek93, Section 4], we have a decomposition

$$\langle c, c' \rangle_{(V, V'), K} = \sum_{u \nmid \infty} \langle c, c' \rangle_{(V, V'), K_u}$$

of the pairing (A.4) into local ones  $\langle c, c' \rangle_{(V, V'), K_u} \in \widehat{K_u}^{\times} \otimes_{\widehat{\mathbb{Z}}} \mathbb{L}$  over all *nonarchimedean* places  $u$  of  $K$ , in which  $\langle c, c' \rangle_{(V, V'), K_u} = \langle c, c' \rangle_{X_u, K_u}^N$  for  $u$  not above  $p$ .

*Remark A.11.* For a place  $u$  of  $K$  above  $p$ , the bi-extension class  $[E_{c'}^c] \in H^1(K_u, E_{c'})$  belongs to  $H_f^1(K_u, E_{c'})$  if and only if  $\langle c, c' \rangle_{(V, V'), K_u} \in O_{K_u}^{\times} \otimes_{\mathbb{Z}_p} \mathbb{L}$ .

## APPENDIX B. PROOF OF THEOREM A.6

Let  $K$  be a finite extension of  $\mathbb{Q}_p$  with residue field  $\kappa$ . Denote by  $W$  the Witt ring of  $\kappa$ , and by  $K_0$  the fraction field of  $W$ , which is canonically a subfield of  $K$ . We fix an algebraic closure  $\overline{K}$  of  $K$  with the residue field  $\overline{\kappa}$ . For every finite extension  $K'$  of  $K_0$  contained in  $\overline{K}$ , we let  $G_{K'}$  be the underlying abstract group of  $\text{Gal}(\overline{K}/K')$ .

**B.1. Preparation.** For a scheme  $\mathcal{Z}$  of finite type over  $O_{K'}$  with  $K'$  a finite extension of  $K_0$  contained in  $\overline{K}$ , we

- put  $Z := \mathcal{Z} \otimes_{O_{K'}} K'$  for the generic fiber,
- put  $\mathcal{Z}_l := \mathcal{Z} \otimes \mathbb{Z}/p^l$  for every integer  $l \geq 1$ ,
- put  $Z := \mathcal{Z} \otimes_{O_{K'}} \kappa'$  for its special fiber, where  $\kappa'$  is the residue field of  $K'$ ,
- denote by  $\mathfrak{Z}$  the formal completion of  $\mathcal{Z}$  along  $Z$ ,
- denote by  $\mathfrak{Z}_{\eta}$  the generic fiber of  $\mathfrak{Z}$ , regarded as an analytic space over  $K'$  in the sense of Berkovich,
- put  $\overline{\mathcal{Z}} := \mathcal{Z} \otimes_{O_{K'}} O_{\overline{K}}$ ,  $\overline{Z} := Z \otimes_{K'} \overline{K}$ , and  $\overline{Z} := Z \otimes_{K'} \overline{\kappa}$ .

We apply the similar notational convention to morphisms over  $O_{K'}$ , as well.<sup>18</sup>

Suppose that  $X$  is a subscheme of  $Z$ , we denote by  $]X[_{\mathfrak{Z}_{\eta}}$  its tubular neighbourhood in  $\mathfrak{Z}_{\eta}$ . We have the quasi-étale site  $]X[_{\mathfrak{Z}_{\eta}, \text{qét}}$  [Ber94, §3] with the natural map  $]X[_{\mathfrak{Z}_{\eta}, \text{qét}} \rightarrow (\widehat{\mathfrak{Z}/X})_{\text{ét}}$ . On the other hand, the natural map  $(\widehat{\mathfrak{Z}/X})_{\text{ét}} \rightarrow X_{\text{ét}}$  is an equivalence of sites [Ber96, Proposition 2.1]. Together, we obtain the *specialization* map  $s_{(X, \mathfrak{Z})} : ]X[_{\mathfrak{Z}_{\eta}, \text{qét}} \rightarrow X_{\text{ét}}$ , and will simply write  $s$  when no confusion arises.

**Definition B.1.** Let  $L$  be a ring. In the situation above, suppose that  $X$  is a closed subscheme of  $Z$  and  $U$  an open subscheme of  $X$ , we define a functor

$$f_{(U, X)}^! : \mathbf{M}(]X[_{\mathfrak{Z}_{\eta}, \text{qét}}, L) \rightarrow \mathbf{M}(]X[_{\mathfrak{Z}_{\eta}, \text{qét}}, L)$$

to be the kernel of the unit transform  $\text{id} \rightarrow g_* \circ g^*$ , where  $g$  denotes the open immersion  $]X \setminus U[_{\mathfrak{Z}_{\eta}} \rightarrow ]X[_{\mathfrak{Z}_{\eta}}$ .

*Remark B.2.* The functors  $g^*$ ,  $g_*$ , and  $f_{(U, X)}^!$  are all exact. Moreover, there is in general no functor  $f : \mathbf{D}^+(X_{\text{ét}}, L) \rightarrow \mathbf{D}^+(X_{\text{ét}}, L)$  such that  $f \circ \text{Rs}_{(X, \mathfrak{Z})} \simeq \text{Rs}_{(X, \mathfrak{Z})} \circ f_{(U, X)}^!$ , even when  $X = Z$ .

**Lemma B.3.** *Let the situation be as in Definition B.1. The diagram*

$$\begin{array}{ccc} F_! \circ F^* \circ \text{Rs}_{(X, \mathfrak{Z})} & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ \text{Rs}_{(X, \mathfrak{Z})} & \longrightarrow & \text{Rs}_{(X, \mathfrak{Z})} \circ g_* \circ g^* \end{array}$$

*of functors from  $\mathbf{D}^+(]X[_{\mathfrak{Z}_{\eta}, \text{qét}}, L)$  to  $\mathbf{D}^+(X_{\text{ét}}, L)$  commutes, where  $F : U \rightarrow X$  denotes the open immersion. In particular, there is a canonical natural transform*

$$F_! \circ F^* \circ \text{Rs}_{(X, \mathfrak{Z})} \rightarrow \text{Rs}_{(X, \mathfrak{Z})} \circ f_{(U, X)}^! : \mathbf{D}^+(]X[_{\mathfrak{Z}_{\eta}, \text{qét}}, L) \rightarrow \mathbf{D}^+(X_{\text{ét}}, L).$$

*Proof.* It suffices to notice that the unit transform  $\text{Rs}_{(X, \mathfrak{Z})} \rightarrow \text{Rs}_{(X, \mathfrak{Z})} \circ g_* \circ g^*$  factors through

$$\text{Rs}_{(X, \mathfrak{Z})} \rightarrow G_* \circ G^* \circ \text{Rs}_{(X, \mathfrak{Z})} \rightarrow G_* \circ \text{Rs}_{(X \setminus U, \mathfrak{Z})} \circ g^* \xrightarrow{\sim} \text{Rs}_{(X, \mathfrak{Z})} \circ g_* \circ g^*,$$

where  $G : X \setminus U \rightarrow X$  denotes the closed immersion. □

<sup>18</sup>Later, we will see  $C, D, E, \mathcal{T}, \mathcal{U}, \mathcal{V}, X, \mathcal{Y}, Z$  for schemes and  $\mathcal{A}, \mathcal{F}, \mathcal{G}, \mathcal{H}$  for morphisms over  $O_{K'}$ .

In what follows, we will work with log-schemes, written as  $(X, L)$  with the first variable the underlying scheme and the second variable the log structure. Since the integral model in Theorem A.6 is strictly semistable, we assume that the log structures are defined in the Zariski topology.

For a scheme  $X$  and a closed subset  $Y$ , we denote by  $L_X^Y$  the log structure  $\mathcal{O}_X \cap j_* \mathcal{O}_{X \setminus Y}^\times \rightarrow \mathcal{O}_X$ , where  $j: X \setminus Y \rightarrow X$  denotes the open immersion. For a log-scheme  $(X, L)$  and a morphism  $f: X' \rightarrow X$ , we write  $f^*L$  for the pullback log structure or simply  $L|_{X'}$  when  $f$  is clear from the context.

We write  $W^{\text{triv}}$  for  $(\text{Spec } W, W^\times)$ ,  $W[t]^\circ$  for  $(\text{Spec } W[t], L)$  where  $L$  is the log structure associated with  $1 \mapsto t$ ,  $W^\circ$  for the fiber of  $W[t]^\circ$  at  $t = 0$ , and  $\kappa^\circ$  for the fiber of  $W^\circ$  at  $\bar{p} = 0$ . Note that the natural morphism  $W[t]^\circ \rightarrow W^{\text{triv}}$  is log-smooth. For every extension  $K'/K_0$  contained in  $\bar{K}$  with the residue field  $\kappa'$ , we put  $O_{K'}^{\text{can}} := (\text{Spec } O_{K'}, L_{\text{Spec } O_{K'}}^{\text{Spec } \kappa'})$ .

Let  $(X, M)$  be a fine log-scheme over a fine base log-scheme  $(S, L)$  of finite type. Recall that an *embedding system* for  $(X, M)/(S, L)$  is a projective system  $\{(X^\star, M^\star) \hookrightarrow (Z^\star, N^\star)\}_{\star=0,1,\dots}$  of exact closed immersions of log-schemes over  $(S, L)$  in which  $X^\star$  is a Zariski hypercovering of  $X$ ,  $M^\star = M|_{X^\star}$ , and  $(Z^\star, N^\star)$  is a fine log-scheme log-smooth over  $(S, L)$  of finite type. Note that embedding system always exists.

In the case where  $(S, L) = W[t]^\circ$  and  $(X, M)$  is a strictly semistable log-scheme over  $\kappa^\circ$  [GK05, §2.1] of finite type, we say that an embedding system  $\{(X^\star, M^\star) \hookrightarrow (Z^\star, N^\star)\}$  for  $(X, M)/(S, L)$  is *admissible* if

- $(X^0, M^0) \rightarrow (Z^0, N^0)$  induces an isomorphism  $(X^0, M^0) \simeq (Z^0, N^0) \times_{W[t]^\circ} \kappa^\circ$ ;
- $Z^0$  is flat and generically smooth over  $W[t]$ , and is smooth over  $W$ ;
- $Y^0 := Z^0 \otimes_{W[t]} W$  is a relative strict normal crossings divisor of  $Z^0$  over  $W$ ;
- $N^0 = L_{Z^0}^{Y^0}$ ;
- $(X^\star, M^\star) \rightarrow (Z^\star, N^\star)$  is (isomorphic to the one) induced from  $(X^0, M^0) \rightarrow (Z^0, N^0)$  in the process described in [GK05, §5.1].

**Notation B.4.** In what follows, when we have a projective system of the form  $\{\square_l\}_{l \geq 1}$  in  $\mathbf{D}^+(S, \mathbb{Z}_p)$  or  $\mathbf{D}_K^+(S, \mathbb{Z}_p)$  for a site  $S$ , we put

$$\square_{\mathbb{Q}} := \left( \mathbf{R} \lim_{\leftarrow l} \square_l \right) \otimes \mathbb{Q}$$

which is an object in  $\mathbf{D}^+(S, \mathbb{Q}_p)$  or  $\mathbf{D}_K^+(S, \mathbb{Q}_p)$ .

**B.2. Rigid de Rham–Witt complexes.** Let  $(X, L)$  be a fine log-scheme over  $\kappa^\circ$  of finite type. Let  $F: U \rightarrow X$  be an open subscheme. For every Zariski hypercovering  $X^\star$  of  $X$ , we put  $F^\star: U^\star := U \times_X X^\star \rightarrow X^\star$ .

Choose an embedding system  $\{(X^\star, L^\star) \hookrightarrow (\mathcal{Y}^\star, M^\star)\}$  for  $(X, M)/W^\circ$ . Denote by  $u: X^\star \rightarrow X$  the augmentation map for the hypercovering, and put  $s^\star := s_{(X^\star, \mathcal{Y}^\star)}$ . We define the rigid de Rham–Witt complex of  $(X, L)$  to be<sup>19</sup>

$$\omega_X := \mathbf{R}u_* \left( \mathbf{R}s_*^\star \left( \Omega_{(\mathcal{Y}^\star, M^\star)/W^\circ}^\bullet \otimes_{\mathcal{O}_{\mathcal{Y}^\star}} \mathcal{O}_{|X^\star|_{\mathcal{Y}^\star}} \right) \right) \in \mathbf{D}^+(X_{\text{ét}}, K_0),$$

where  $\Omega_{(\mathcal{Y}^\star, M^\star)/W^\circ}^\bullet$  denotes the complex on  $\mathcal{Y}_{\text{ét}}^\star$  of relative logarithmic differentials of the log-smooth morphism  $(\mathcal{Y}^\star, M^\star)/W^\circ$ . By (the same argument in) [GK05, Lemma 1.4], the complex  $\omega_X$  does not depend on the choice of the embedding system for  $(X, M)/W^\circ$ .<sup>20</sup>

On the other hand, choose an embedding system  $\{(X^\star, L^\star) \hookrightarrow (Z^\star, N^\star)\}$  for  $(X, L)/W[t]^\circ$ , hence for  $(X, L)/W^{\text{triv}}$ . Then  $\{(X^\star, L^\star) \hookrightarrow (\mathcal{Y}^\star, M^\star) := (Z^\star, N^\star) \times_{W[t]^\circ} W^\circ\}$  is an embedding system for  $(X, L)/W^\circ$ . We have the short exact sequence

$$0 \rightarrow \Omega_{(\mathcal{Y}^\star, M^\star)/W^\circ}^{q-1} \rightarrow \Omega_{(Z^\star, N^\star)/W^{\text{triv}}}^q \otimes_{\mathcal{O}_{Z^\star}} \mathcal{O}_{\mathcal{Y}^\star} \rightarrow \Omega_{(\mathcal{Y}^\star, M^\star)/W^\circ}^q \rightarrow 0$$

of coherent sheaves on  $\mathcal{Y}_{\text{ét}}^\star$ , in which the first map is given by  $\wedge d \log t$ , for every  $q \geq 0$  compatible with differentials. We put

$$\tilde{\omega}_X := \mathbf{R}u_* \left( \mathbf{R}s_*^\star \left( \Omega_{(Z^\star, N^\star)/W^{\text{triv}}}^\bullet \otimes_{\mathcal{O}_{Z^\star}} \mathcal{O}_{|X^\star|_{\mathcal{Y}^\star}} \right) \right) \in \mathbf{D}^+(X_{\text{ét}}, K_0).$$

<sup>19</sup>More precisely, it should be called *convergent de Rham–Witt complex* since it gives the log convergent cohomology in general. However, later we will take  $(X, L)$  to be strictly semistable and proper.

<sup>20</sup>Of course,  $\omega_X$  depends on the log structure  $L$ . However, as a common practice for de Rham–Witt complexes, we will not include  $L$  in the notation.



Then there is a distinguished triangle

$$(B.1) \quad \omega_X^\Delta: \quad \omega_X[-1] \rightarrow \widetilde{\omega}_X \rightarrow \omega_X \xrightarrow{N} \omega_X$$

in  $\mathbf{D}^+(\mathcal{X}_{\text{ét}}, K_0)$ , where  $N$  denotes the connecting map; it is independent of the choice of the embedding system for  $(X, L)/W[t]^\circ$ . Moreover, for a morphism  $f: (X', L') \rightarrow (X, L)$  of fine log-schemes over  $\kappa^\circ$  of finite type, we have an induced map  $\omega_X^\Delta \rightarrow \mathbf{R}f_* \widetilde{\omega}_{X'}^\Delta$  of distinguished triangles in  $\mathbf{D}^+(\mathcal{X}_{\text{ét}}, K_0)$ .

We put

$$\begin{aligned} \omega_{(U, X)} &:= \mathbf{R}u_* \left( \mathbf{R}s_*^* \mathbf{f}_{(U^*, X^*)}^! \left( \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^\bullet \otimes_{\mathcal{O}_{\mathcal{Y}^*}} \mathcal{O}_{|X^*|_{\mathcal{Y}^*}} \right) \right), \\ \widetilde{\omega}_{(U, X)} &:= \mathbf{R}u_* \left( \mathbf{R}s_*^* \mathbf{f}_{(U^*, X^*)}^! \left( \Omega_{(\mathcal{Z}^*, N^*)/W^{\text{triv}}}^\bullet \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \mathcal{O}_{|X^*|_{\mathcal{Y}^*}} \right) \right), \end{aligned}$$

both in  $\mathbf{D}^+(\mathcal{X}_{\text{ét}}, K_0)$  (see Definition B.1). Then by definition, we have a distinguished triangle

$$(B.2) \quad \omega_{(U, X)}^\Delta: \quad \omega_{(U, X)}[-1] \rightarrow \widetilde{\omega}_{(U, X)} \rightarrow \omega_{(U, X)} \xrightarrow{N} \omega_{(U, X)}$$

in  $\mathbf{D}^+(\mathcal{X}_{\text{ét}}, K_0)$ , and a distinguished triangle of distinguished triangles

$$(B.3) \quad \omega_{(U, X)}^\Delta \rightarrow \omega_X^\Delta \rightarrow \mathbf{G}_* \omega_{X \setminus U}^\Delta \xrightarrow{+1}$$

where  $\mathbf{G}: X \setminus U \rightarrow X$  denotes the closed immersion.

From now on, we assume that  $(X, L)$  is strictly semistable over  $\kappa^\circ$ . We recall the construction of several crystalline complexes of  $(X, L)$ . For every  $l \geq 1$ , let  $\mathcal{D}_l^\star$  and  $\mathcal{E}_l^\star$  be the (scheme part of the) PD envelops of  $X^\star$  in  $\mathcal{Y}_l^\star$  and  $\mathcal{Z}_l^\star$  (over the base  $W$  equipped with the usual PD structure), respectively.<sup>21</sup> We define

$$\begin{aligned} \mathcal{C}_{l, (X, L)/W^\circ} &:= \mathbf{R}u_* \left( \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^\bullet \otimes_{\mathcal{O}_{\mathcal{Y}^*}} \mathcal{O}_{\mathcal{D}_l^\star} \right), \\ \widetilde{\mathcal{C}}_{l, (X, L)/W^\circ} &:= \mathbf{R}u_* \left( \Omega_{(\mathcal{Z}^*, N^*)/W^{\text{triv}}}^\bullet \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \mathcal{O}_{\mathcal{D}_l^\star} \right), \\ \mathcal{C}_{l, (X, L)/W^{\text{triv}}} &:= \mathbf{R}u_* \left( \Omega_{(\mathcal{Z}^*, N^*)/W^{\text{triv}}}^\bullet \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \mathcal{O}_{\mathcal{E}_l^\star} \right), \\ \mathcal{C}_{l, (X, L)/W[t]^\circ} &:= \mathbf{R}u_* \left( \Omega_{(\mathcal{Z}^*, N^*)/W[t]^\circ}^\bullet \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \mathcal{O}_{\mathcal{E}_l^\star} \right), \end{aligned}$$

all in  $\mathbf{D}^+(\mathcal{X}_{\text{ét}}, W)$ . It is well-known that the above objects do not depend on the choice of the embedding system for  $(X, L)/W[t]^\circ$ . In fact,  $\{\mathcal{C}_{l, (X, L)/W^\circ}\}_{l \geq 1}$  is nothing but the modified de Rham–Witt complex  $\{W_l \omega_X^\bullet\}_{l \geq 1}$  [Hyo91, HK94], and  $\{\widetilde{\mathcal{C}}_{l, (X, L)/W^\circ}\}_{l \geq 1}$  is simply  $\{W_l \widetilde{\omega}_X^\bullet\}_{l \geq 1}$ .

Applying Notation B.4, we obtain a distinguished triangle

$$(B.4) \quad W_{\mathbb{Q}} \omega_X^\bullet[-1] \rightarrow W_{\mathbb{Q}} \widetilde{\omega}_X^\bullet \rightarrow W_{\mathbb{Q}} \omega_X^\bullet \xrightarrow{N} W_{\mathbb{Q}} \omega_X^\bullet$$

in  $\mathbf{D}^+(\mathcal{X}_{\text{ét}}, K_0)$ , similar to (B.1), in which the first arrow is given by  $\wedge d \log t$ , the second arrow is the natural one, and the last arrow is the connecting map.

We would like to compare (B.1) and (B.4). We have a canonical map  $\mathcal{O}_{\mathcal{Y}_l^\star/X^\star} \rightarrow \varprojlim_l \mathcal{O}_{\mathcal{D}_l^\star} = \mathbf{R} \varprojlim_l \mathcal{O}_{\mathcal{D}_l^\star}$  as in [Ber97, (1.9.2)], which induces maps

$$\begin{aligned} \mathbf{R}s_*^* \left( \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^\bullet \otimes_{\mathcal{O}_{\mathcal{Y}^*}} \mathcal{O}_{|X^*|_{\mathcal{Y}^*}} \right) &\simeq \left( \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^\bullet \otimes_{\mathcal{O}_{\mathcal{Y}^*}} \mathcal{O}_{\mathcal{Y}_l^\star/X^\star} \right) \otimes \mathbb{Q} \rightarrow \left( \mathbf{R} \varprojlim_l \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^\bullet \otimes_{\mathcal{O}_{\mathcal{Y}^*}} \mathcal{O}_{\mathcal{D}_l^\star} \right) \otimes \mathbb{Q} \\ \mathbf{R}s_*^* \left( \Omega_{(\mathcal{Z}^*, N^*)/W^{\text{triv}}}^\bullet \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \mathcal{O}_{|X^*|_{\mathcal{Y}^*}} \right) &\simeq \left( \Omega_{(\mathcal{Z}^*, N^*)/W^{\text{triv}}}^\bullet \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \mathcal{O}_{\mathcal{Y}_l^\star/X^\star} \right) \otimes \mathbb{Q} \rightarrow \left( \mathbf{R} \varprojlim_l \Omega_{(\mathcal{Z}^*, N^*)/W^{\text{triv}}}^\bullet \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \mathcal{O}_{\mathcal{D}_l^\star} \right) \otimes \mathbb{Q} \end{aligned}$$

in  $\mathbf{D}^+(\mathcal{X}_{\text{ét}}^\star, K_0)$ . These maps are in fact equivalences since  $(X^\star, L^\star)$  is strictly semistable over  $\kappa^\circ$  by an argument similar to [Ber97, §1.9]. Applying  $\mathbf{R}u_*$ , we obtain equivalences

$$(B.5) \quad \omega_X \xrightarrow{\sim} W_{\mathbb{Q}} \omega_X^\bullet, \quad \widetilde{\omega}_X \xrightarrow{\sim} W_{\mathbb{Q}} \widetilde{\omega}_X^\bullet,$$

<sup>21</sup>The natural morphism  $\mathcal{D}_l^\star \rightarrow \mathcal{E}_l^\star \otimes_{W(t)} W$  is an isomorphism, where  $W(t)$  denotes the PD envelop of  $(W[t], (t))$  over  $W$ .

under which (B.1) is equivalent to (B.4). By Lemma B.3, we have a natural map

$$(B.6) \quad \begin{aligned} F_! F^* W_{\mathbb{Q}} \widetilde{\omega}_X^\bullet &\rightarrow \mathrm{Ru}_* F_! (F^*)^* \left( \left( \mathrm{R} \lim_{\leftarrow} \Omega_{(\mathcal{Z}^*, N^*)/W^{\mathrm{triv}}}^\bullet \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \mathcal{O}_{\mathcal{D}_t^*} \right) \otimes \mathbb{Q} \right) \\ &\xrightarrow{\sim} \mathrm{Ru}_* \left( F_! (F^*)^* \mathrm{Rs}_*^* \left( \Omega_{(\mathcal{Z}^*, N^*)/W^{\mathrm{triv}}}^\bullet \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \mathcal{O}_{|X^*|_{\mathfrak{y}^*}} \right) \right) \\ &\rightarrow \mathrm{Ru}_* \left( \mathrm{Rs}_*^* f_{(U^*, X^*)}^! \left( \Omega_{(\mathcal{Z}^*, N^*)/W^{\mathrm{triv}}}^\bullet \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \mathcal{O}_{|X^*|_{\mathfrak{y}^*}} \right) \right) = \widetilde{\omega}_{(U, X)} \end{aligned}$$

in  $\mathbf{D}^+(X_{\mathrm{ét}}, K_0)$ .

When the model in Theorem A.6 is not smooth, we also need a cohomological variant of the rigid de Rham–Witt complex, which we now introduce. We choose an admissible embedding system  $\{(X^*, L^*) \hookrightarrow (\mathcal{Z}^*, N^*)\}$  for  $(X, L)/W[t]^\circ$ .

For every  $q \geq 0$ , we have a natural subsheaf  $\Omega_{\mathcal{Z}^*/W}^q \subseteq \Omega_{(\mathcal{Z}^*, N^*)/W^{\mathrm{triv}}}^q$ . Put

$$\Xi_{\mathcal{Z}^*}^q := \frac{\Omega_{(\mathcal{Z}^*, N^*)/W^{\mathrm{triv}}}^{q+1}}{\Omega_{\mathcal{Z}^*/W}^{q+1}},$$

which is an  $\mathcal{O}_{\mathfrak{y}^*}$ -module. The map  $\wedge d \log t$  induces a diagram

$$(B.7) \quad \begin{array}{ccc} \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^q & & \\ \downarrow & \searrow & \\ \Omega_{(\mathcal{Z}^*, N^*)/W^{\mathrm{triv}}}^{q+1} \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \mathcal{O}_{\mathfrak{y}^*} & \longrightarrow & \Xi_{\mathcal{Z}^*}^q \end{array}$$

of coherent sheaves on  $\mathcal{Y}_{\mathrm{ét}}^*$ , compatible with differentials. Define

$$\begin{aligned} \omega_X^+ &:= \mathrm{Ru}_* \left( \mathrm{Rs}_*^* \left( \Xi_{\mathcal{Z}^*}^\bullet \otimes_{\mathcal{O}_{\mathfrak{y}^*}} \mathcal{O}_{|X^*|_{\mathfrak{y}^*}} \right) \right), \\ \omega_{(U, X)}^+ &:= \mathrm{Ru}_* \left( \mathrm{Rs}_*^* f_{(U^*, X^*)}^! \left( \Xi_{\mathcal{Z}^*}^\bullet \otimes_{\mathcal{O}_{\mathfrak{y}^*}} \mathcal{O}_{|X^*|_{\mathfrak{y}^*}} \right) \right), \end{aligned}$$

and we have a natural diagram

$$(B.8) \quad \begin{array}{ccc} \omega_{(U, X)}[-1] & & \\ \downarrow & \searrow & \\ \widetilde{\omega}_{(U, X)} & \longrightarrow & \omega_{(U, X)}^+[-1] \end{array}$$

in  $\mathbf{D}^+(X_{\mathrm{ét}}, K_0)$ , in which the vertical arrow is same as the first arrow in the first line in B.3. Let  $\{W_l \Xi_X^\bullet\}_{l \geq 1}$  be the cohomological de Rham–Witt complex defined in [Sat13, Definition 8.3]. Similar to (B.5), we have a natural equivalence

$$\omega_X^+ \simeq W_{\mathbb{Q}} \Xi_X^\bullet$$

by [Sat13, Proposition 8.4]. Similar to (B.6), we have a natural map

$$(B.9) \quad F_! F^* W_{\mathbb{Q}} \Xi_X^\bullet \rightarrow \omega_{(U, X)}^+$$

in  $\mathbf{D}^+(X_{\mathrm{ét}}, K_0)$ , fitting into the following diagram

$$(B.10) \quad \begin{array}{ccc} F_! F^* W_{\mathbb{Q}} \widetilde{\omega}_X^\bullet & \longrightarrow & F_! F^* W_{\mathbb{Q}} \Xi_X^\bullet[-1] \\ (B.6) \downarrow & & \downarrow (B.9) \\ \widetilde{\omega}_{(U, X)} & \longrightarrow & \omega_{(U, X)}^+[-1] \end{array}$$

in which the upper horizontal arrow is induced by the one in [Sat13, Proposition 8.10].

**B.3. Log rigid cohomology with proper support.** Let the situation be as in the previous subsection with  $(X, L)$  strictly semistable over  $\kappa^\circ$ . We also assume that  $X$  is proper of pure dimension  $n - 1$ . For every  $h \geq 1$ , let  $X^{(h)}$  be the disjoint union of intersections of  $h$  different irreducible components of  $X$ , and put  $U^{(h)} := U \times_X X^{(h)}$ .

Recall from [GK05, §1.5] that for a scheme  $Y$  over  $X$  of finite type, we have the log rigid cohomology  $H_{\text{rig}}^\bullet(Y/W^\circ)$  and the log convergent cohomology  $H_{\text{conv}}^\bullet(Y/W^\circ)$  for the log-scheme  $(Y, L|_Y)$ , with a natural map  $H_{\text{rig}}^\bullet(Y/W^\circ) \rightarrow H_{\text{conv}}^\bullet(Y/W^\circ)$ . In particular, we have

$$\mathbf{H}^q(X_{\text{ét}}, \omega_X) = H_{\text{conv}}^q(X/W^\circ), \quad \mathbf{H}^q(X_{\text{ét}}, \mathbf{G}_* \omega_{X|U}) = H_{\text{conv}}^q(X \setminus U/W^\circ).$$

for every  $q \geq 0$ .<sup>22</sup> Moreover, the natural map  $H_{\text{rig}}^\bullet(X/W^\circ) \rightarrow H_{\text{conv}}^\bullet(X/W^\circ)$  is an isomorphism by [GK05, Theorem 5.3(ii)].

**Definition B.5.** We define, in an *ad hoc* way, the *log rigid cohomology of  $U$  with proper support* to be

$$H_{\text{rig}}^q((U, X)/W^\circ) := \mathbf{H}^q(X_{\text{ét}}, \omega_{(U, X)}),$$

which *a priori* depends on the embedding  $U \hookrightarrow X$ . For  $U \subseteq U' \subseteq X$ , we have a natural pushforward map  $H_{\text{rig}}^q((U, X)/W^\circ) \rightarrow H_{\text{rig}}^q((U', X)/W^\circ)$  by construction.

The distinguished triangle (B.3) induces a long exact sequence

$$(B.11) \quad \cdots \rightarrow H_{\text{conv}}^{q-1}(X \setminus U/W^\circ) \rightarrow H_{\text{rig}}^q((U, X)/W^\circ) \rightarrow H_{\text{rig}}^q(X/W^\circ) \rightarrow H_{\text{conv}}^q(X \setminus U/W^\circ) \rightarrow \cdots$$

in  $\mathbf{M}(K_0)$ .

We now review the weight spectral sequence for log rigid cohomology from [GK05, §5], which is the rigid analogue of Mokrane's spectral sequence for log crystalline cohomology [Mok93]. Take an admissible embedding system  $\{(X^*, L^*) \hookrightarrow (Z^*, N^*)\}$  for  $(X, L)/W[t]^\circ$ . For  $j \geq 0$ , put

$$P_j \Omega_{(Z^*, N^*)/W^{\text{triv}}}^q := \text{Im} \left( \Omega_{(Z^*, N^*)/W^{\text{triv}}}^j \otimes \Omega_{Z^*/W}^{q-j} \rightarrow \Omega_{(Z^*, N^*)/W^{\text{triv}}}^q \right).$$

We have the double complex

$$A_{Z^*}^{ij} := \frac{\Omega_{(Z^*, N^*)/W^{\text{triv}}}^{i+j+1}}{P_j \Omega_{(Z^*, N^*)/W^{\text{triv}}}^{i+j+1}}$$

of  $\mathcal{O}_{\mathcal{Y}^*}$ -modules, in which the differential  $A_{Z^*}^{ij} \rightarrow A_{Z^*}^{(i+1)j}$  is given by  $(-1)^j d$  and the differential  $A_{Z^*}^{ij} \rightarrow A_{Z^*}^{i(j+1)}$  is given by  $\wedge d \log t$ , with the filtration

$$P_k A_{Z^*}^{ij} := \frac{P_{2j+k+1} \Omega_{(Z^*, N^*)/W^{\text{triv}}}^{i+j+1}}{P_j \Omega_{(Z^*, N^*)/W^{\text{triv}}}^{i+j+1}}$$

for  $k \geq -j$ . In particular,  $A_{Z^*}^{\bullet 0}$  is nothing but the complex  $\Xi_{Z^*}^\bullet$  from the previous subsection. Let  $A_{Z^*}^\bullet$  be the total complex of  $A_{Z^*}^{\bullet \bullet}$ . It is shown in [GK05, §5.2] that the augmentation map  $\Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^\bullet \rightarrow \Xi_{Z^*}^\bullet = A_{Z^*}^{\bullet 0}$  in (B.7) induces an equivalence  $\Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^\bullet \xrightarrow{\sim} A_{Z^*}^\bullet$  in  $\mathbf{D}^+(\mathcal{Y}_{\text{ét}}^*, K_0)$ . Then the total filtration on  $A_{Z^*}^\bullet$  induces spectral sequences

$$\begin{aligned} E(X)_1^{-k, q+k} &= \bigoplus_{j \geq \max\{0, -k\}} H_{\text{rig}}^{q-2j-k}(X^{(2j+k+1)}/K_0) \Rightarrow H_{\text{rig}}^q(X/W^\circ), \\ E(U)_1^{-k, q+k} &= \bigoplus_{j \geq \max\{0, -k\}} H_{\text{rig}}^{q-2j-k}(U^{(2j+k+1)}/K_0) \Rightarrow H_{\text{rig}}^q(U/W^\circ), \end{aligned}$$

which already appeared in [GK05, (4)], and

$$(B.12) \quad E(X \setminus U)_1^{-k, q+k} = \bigoplus_{j \geq \max\{0, -k\}} H_{\text{rig}}^{q-2j-k}(X^{(2j+k+1)} \setminus U^{(2j+k+1)}/K_0) \Rightarrow H_{\text{conv}}^q(X \setminus U/W^\circ),$$

<sup>22</sup>Here, we use the fact that computing cohomology of coherent sheaves in the quasi-étale topology of analytic spaces is the same as in the  $G$ -topology.

$$(B.13) \quad E(\mathbf{U}, \mathbf{X})_1^{-k, q+k} = \bigoplus_{j \geq \max\{0, -k\}} \mathbf{H}_{\text{rig}}^{q-2j-k}((\mathbf{U}^{(2j+k+1)}, \mathbf{X}^{(2j+k+1)})/K_0) \Rightarrow \mathbf{H}_{\text{rig}}^q((\mathbf{U}, \mathbf{X})/W^\circ).$$

Here,  $\mathbf{H}_{\text{rig}}^\bullet((\mathbf{U}^{(2j+k+1)}, \mathbf{X}^{(2j+k+1)})/K_0)$  is defined similarly as for  $\mathbf{H}_{\text{rig}}^\bullet((\mathbf{U}, \mathbf{X})/W^\circ)$  but without the log structure, which in fact coincides with the rigid cohomology with proper support  $\mathbf{H}_{\text{c,rig}}^\bullet(\mathbf{U}^{(2j+k+1)}/K_0)$  defined by Berthelot since  $\mathbf{X}^{(2j+k+1)}$  is proper. In particular, the spectral sequence (B.13) can also be written as

$$E_c(\mathbf{U})_1^{-k, q+k} = \bigoplus_{j \geq \max\{0, -k\}} \mathbf{H}_{\text{c,rig}}^{q-2j-k}(\mathbf{U}^{(2j+k+1)}/K_0) \Rightarrow \mathbf{H}_{\text{rig}}^q((\mathbf{U}, \mathbf{X})/W^\circ).$$

The following two lemmas will be used later.

**Lemma B.6.** *Let  $d \geq 1$  be an integer. Suppose that  $\dim(\mathbf{X}^{(h)} \setminus \mathbf{U}^{(h)}) \leq d - h$  for every  $h \geq 1$ .*

- (1) *The natural map  $\mathbf{H}_{\text{rig}}^q((\mathbf{U}, \mathbf{X})/W^\circ) \rightarrow \mathbf{H}_{\text{rig}}^q(\mathbf{X}/W^\circ)$  is an isomorphism for  $q \geq 2d$ .*
- (2) *The natural map  $E_c(\mathbf{U})_1^{-k, q+k} \rightarrow E(\mathbf{X})_1^{-k, q+k}$  is an isomorphism for  $q \geq 2d - |k|$ .*
- (3) *For the map  $E_c(\mathbf{U})_1^{0, 2d-1} \rightarrow E(\mathbf{X})_1^{0, 2d-1}$ , the direct summand*

$$\bigoplus_{j \geq 1} \mathbf{H}_{\text{c,rig}}^{2d-1-2j}(\mathbf{U}^{(2j+1)}/K_0) \rightarrow \bigoplus_{j \geq 1} \mathbf{H}_{\text{rig}}^{2d-1-2j}(\mathbf{X}^{(2j+1)}/K_0)$$

*is an isomorphism.*

*Proof.* For (1), it suffices to show that  $\mathbf{H}_{\text{conv}}^q(\mathbf{X} \setminus \mathbf{U}/W^\circ) = 0$  for  $q \geq 2d - 1$ . By the spectral sequence (B.12), it suffices to show that  $\mathbf{H}_{\text{rig}}^{q-2j-k}(\mathbf{X}^{(2j+k+1)} \setminus \mathbf{U}^{(2j+k+1)})/K_0 = 0$  for every  $j, k$ , and  $q \geq 2d - 1$ , which follows from the fact that

$$2d - 1 - 2j - k > 2(d - 2j - k - 1) \geq 2 \dim(\mathbf{X}^{(2j+k+1)} \setminus \mathbf{U}^{(2j+k+1)}).$$

For (2), it follows from the fact that

$$\mathbf{H}_{\text{rig}}^{q-2j-k-1}(\mathbf{X}^{(2j+k+1)} \setminus \mathbf{U}^{(2j+k+1)})/K_0 = \mathbf{H}_{\text{rig}}^{q-2j-k}(\mathbf{X}^{(2j+k+1)} \setminus \mathbf{U}^{(2j+k+1)})/K_0 = 0$$

for every  $j \geq \max\{0, -k\}$  when  $q \geq 2d - |k|$ .

For (3), it follows from the fact that

$$\mathbf{H}_{\text{rig}}^{2d-1-2j-1}(\mathbf{X}^{(2j+1)} \setminus \mathbf{U}^{(2j+1)})/K_0 = \mathbf{H}_{\text{rig}}^{2d-1-2j}(\mathbf{X}^{(2j+1)} \setminus \mathbf{U}^{(2j+1)})/K_0 = 0$$

for  $j \geq 1$ . □

**Lemma B.7.** *There is a spectral sequence  $E_c^+(\mathbf{U})_1^{-k, q+k} \Rightarrow \mathbf{H}^q(\mathbf{X}_{\text{ét}}, \omega_{(\mathbf{U}, \mathbf{X})}^+)$  with*

$$E_c^+(\mathbf{U})_1^{-k, q+k} = \begin{cases} \mathbf{H}_{\text{c,rig}}^{q-k}(\mathbf{U}^{(k+1)}/K_0), & k \geq 0; \\ 0, & k < 0. \end{cases}$$

*Moreover, the map  $\mathbf{H}_{\text{rig}}^q((\mathbf{U}, \mathbf{X})/W^\circ) = \mathbf{H}^q(\mathbf{X}_{\text{ét}}, \omega_{(\mathbf{U}, \mathbf{X})}) \rightarrow \mathbf{H}^q(\mathbf{X}_{\text{ét}}, \omega_{(\mathbf{U}, \mathbf{X})}^+)$  is abutted by the map  $E_c(\mathbf{U})_1^{-k, q+k} \rightarrow E_c^+(\mathbf{U})_1^{-k, q+k}$  given by the obvious projections.*

*Proof.* The spectral sequence follows from the filtration  $P_k A_{\mathcal{Z}^\star}^{i0}$  of  $A_{\mathcal{Z}^\star}^{i0} = \Xi_{\mathcal{Z}^\star}^i$ . Recall that the map  $\omega_{(\mathbf{U}, \mathbf{X})} \rightarrow \omega_{(\mathbf{U}, \mathbf{X})}^+$  is induced by the natural projection map  $A_{\mathcal{Z}^\star}^\bullet \rightarrow A_{\mathcal{Z}^\star}^{00}$ . The lemma follows since  $P_k A_{\mathcal{Z}^\star}^{i0}$  is the image of  $P_k A_{\mathcal{Z}^\star}^i$  under this map. □

**B.4. Abel–Jacobi map via rigid cohomology.** We start to prove Theorem A.6. We fix a uniformizer  $\varpi$  of  $K$ , and regard  $O_K$  as an  $W[t]$ -ring via  $t \mapsto \varpi$ , making  $O_K^{\text{can}}$  an exact closed log-subscheme of  $W[t]^\circ$ . Let  $\mathbb{B}_{\text{cris}}$  be the crystalline period ring and  $\mathbb{B}_{\text{st}}$  the semistable period ring with respect to  $\varpi$ .

We fix a proper strictly semistable scheme  $\mathcal{X}$  over  $O_K$  of pure (absolute) dimension  $n \geq 2$ . Then  $(\mathcal{X}, L_{\mathcal{X}}^X)$  is log-smooth over  $O_K^{\text{can}}$ , and  $(\mathcal{X}, L := L_{\mathcal{X}}^X|_{\mathcal{X}})$  is strictly semistable over  $\kappa^\circ$  of finite type. Consider an open immersion  $\mathcal{F}: \mathcal{U} \rightarrow \mathcal{X}$ . The following definition will be frequently used later, which is related to condition (2) in Theorem A.6.

**Definition B.8.** For an integer  $d \geq 1$ , we say that  $\mathcal{U}$  is *d-dense* if  $\dim(\mathcal{X}^{(h)} \setminus \mathcal{U}^{(h)}) \leq d - h$  for every  $h \geq 1$ .

Put  $i: X \rightarrow \mathcal{X}$  and  $j: X \rightarrow \mathcal{X}$  for the special fiber and the generic fiber of  $\mathcal{X}$ , respectively. Take an integer  $d$  satisfying  $0 \leq d < p - 1$ . Let  $\{\mathcal{S}_l(d)_X\}_{l \geq 1}$  be the projective system of Kato's (log) syntomic complexes (of  $\mathbb{Z}_p$ -modules on  $X_{\text{ét}}$ ) for  $(X, L_X^\times)$ .<sup>23</sup> We have the period map  $\{\mathcal{S}_l(d)_X \rightarrow i^* \mathbf{R}j_* \mu_{p^l}^{\otimes d}\}_{l \geq 1}$  which induces equivalences  $\mathcal{S}_l(d)_X \xrightarrow{\sim} \tau^{\leq d} i^* \mathbf{R}j_* \mu_{p^l}^{\otimes d}$  ([Kat94, Tsu00]). Put  $i_{\mathcal{U}}: \mathbf{U} \rightarrow \mathcal{U}$  and  $j_{\mathcal{U}}: \mathbf{U} \rightarrow \mathcal{U}$  for the special fiber and the generic fiber of  $\mathcal{U}$ , respectively. We have a sequence of maps

$$F_! F^* i^* \mathbf{R}j_* \mu_{p^l}^{\otimes d} \xrightarrow{\sim} F_! i_{\mathcal{U}}^* \mathbf{R}j_{\mathcal{U}} \mu_{p^l}^{\otimes d} \xrightarrow{\sim} i^* \mathcal{F}_! \mathbf{R}j_{\mathcal{U}} \mu_{p^l}^{\otimes d} \rightarrow i^* \mathbf{R}j_* F_! \mu_{p^l}^{\otimes d},$$

in which the last one is given by adjunction, compatible with  $l$ . Then we obtain the maps

$$\begin{aligned} \mathrm{R}\Gamma\left(X_{\text{ét}}, F_! F^* \mathbf{R}\lim_{\leftarrow l} \mathcal{S}_l(d)_X\right) &\rightarrow \mathrm{R}\Gamma\left(X_{\text{ét}}, \mathbf{R}\lim_{\leftarrow l} F_! F^* \mathcal{S}_l(d)_X\right) \\ &\xrightarrow{\sim} \mathbf{R}\lim_{\leftarrow l} \mathrm{R}\Gamma\left(X_{\text{ét}}, F_! F^* \mathcal{S}_l(d)_X\right) \\ &\rightarrow \mathbf{R}\lim_{\leftarrow l} \mathrm{R}\Gamma\left(X_{\text{ét}}, F_! F^* i^* \mathbf{R}j_* \mu_{p^l}^{\otimes d}\right) \\ &\rightarrow \mathbf{R}\lim_{\leftarrow l} \mathrm{R}\Gamma\left(X_{\text{ét}}, i^* \mathbf{R}j_* F_! \mu_{p^l}^{\otimes d}\right) \\ &\xrightarrow{\sim} \mathbf{R}\lim_{\leftarrow l} \mathrm{R}\Gamma\left(X_{\text{ét}}, F_! \mu_{p^l}^{\otimes d}\right) \end{aligned}$$

where we have used the proper base change for the last equivalence. Put

$$\mathrm{R}\Gamma_c(U, \mathbb{Q}_p(d)) := \mathbf{R}\lim_{\leftarrow l} \mathrm{R}\Gamma\left(X_{\text{ét}}, F_! \mu_{p^l}^{\otimes d}\right) \otimes \mathbb{Q},$$

which is nothing but the continuous étale cohomology complex of  $U$  with proper support with coefficients in  $\mathbb{Q}_p$ . Then the composition of the previous sequence gives a map

$$(B.14) \quad \mathrm{R}\Gamma\left(X_{\text{ét}}, F_! F^* \mathcal{S}_Q(d)_X\right) \rightarrow \mathrm{R}\Gamma_c(U, \mathbb{Q}_p(d))$$

(see Notation B.4 for  $\mathcal{S}_Q(d)_X$ ).

The Hochschild–Serre spectral sequence induces an edge map

$$\mathrm{H}_c^q(U, \mathbb{Q}_p(d))^0 \rightarrow \mathrm{H}^1(K, \mathrm{H}_c^{q-1}(\bar{U}, \mathbb{Q}_p(d))),$$

where

$$\mathrm{H}_c^q(U, \mathbb{Q}_p(d))^0 := \mathrm{Ker}\left(\mathrm{H}_c^q(U, \mathbb{Q}_p(d)) \rightarrow \mathrm{H}_c^q(\bar{U}, \mathbb{Q}_p(d))\right).$$

Let  $\mathbf{H}^q(X_{\text{ét}}, F_! F^* \mathcal{S}_Q(d)_X)^\heartsuit$  be the inverse image of  $\mathrm{H}_c^q(U, \mathbb{Q}_p(d))^0$  under the map (B.14) (after taking  $q$ -th cohomology). Then we have the induced composite map

$$(B.15) \quad \alpha_q: \mathbf{H}^q(X_{\text{ét}}, F_! F^* \mathcal{S}_Q(d)_X)^\heartsuit \rightarrow \mathrm{H}^1(K, \mathrm{H}_c^{q-1}(\bar{U}, \mathbb{Q}_p(d))) \rightarrow \mathrm{H}^1(K, \mathrm{H}_c^{q-1}(\bar{U}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\text{cris}}),$$

where in the last map we use the canonical embedding  $\mathbb{Q}_p(d) \hookrightarrow \mathbb{B}_{\text{cris}}$  in  $\mathbf{M}_K(\mathbb{Q}_p)$ .

To study the kernel of  $\alpha_{2d}$ , we need to use crystalline complexes for  $\mathcal{X}$  rather than its special fiber. Let  $\{\mathcal{C}_{l, X/W^{\text{triv}}}\}_{l \geq 1}$  and  $\{\mathcal{C}_{l, X/W[t]^\circ}\}_{l \geq 1}$  be the projective systems of complexes defined similarly as  $\{\mathcal{C}_{l, (X, L)/W^{\text{triv}}}\}_{l \geq 1}$  and  $\{\mathcal{C}_{l, (X, L)/W[t]^\circ}\}_{l \geq 1}$  in §B.2, respectively, for which we use an embedding system for  $(X, L_X^\times)/W[t]^\circ$  and just replace  $\mathcal{E}_l^*$  by the PD envelop of  $X_l$  in  $\mathcal{Z}_l^*$ . We have the following commutative diagram

$$(B.16) \quad \begin{array}{ccccc} \mathcal{S}_Q(d)_X & \longrightarrow & \mathcal{C}_{Q, X/W^{\text{triv}}} & \longrightarrow & \mathcal{C}_{Q, X/W[t]^\circ} \\ & & \downarrow & & \downarrow \\ & & W_{\mathbb{Q}} \tilde{\omega}_X^\bullet & \longrightarrow & W_{\mathbb{Q}} \omega_X^\bullet \end{array}$$

<sup>23</sup>We will recall the construction of many syntomic and crystalline complexes in §B.6 in which  $\mathcal{S}_l(d)_X$  is a special case.

in  $\mathbf{D}^+(\mathcal{X}_{\acute{e}t}, \mathbb{Q}_p)$  (see Notation B.4), in which the first arrow is the natural map from the syntomic complex to the crystalline complex over  $W^{\text{triv}}$ , and the two vertical arrows are given by specialization at  $t = 0$ .

Using (B.6), we obtain a map

$$(B.17) \quad F_! F^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}} \rightarrow \tilde{\omega}_{(U, X)}$$

in  $\mathbf{D}^+(\mathcal{X}_{\acute{e}t}, \mathbb{Q}_p)$ .

**Lemma B.9.** *Suppose that  $\mathcal{U}$  is  $d$ -dense if  $d \geq 1$ . Then the composite map*

$$\mathbf{H}^{2d}(\mathcal{X}_{\acute{e}t}, F_! F^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}}) \rightarrow \mathbf{H}^{2d}(\mathcal{X}_{\acute{e}t}, \tilde{\omega}_{(U, X)}) \rightarrow \mathbf{H}^{2d}(\mathcal{X}_{\acute{e}t}, \omega_{(U, X)}) = \mathbf{H}_{\text{rig}}^{2d}((U, X)/W^\circ)$$

(Definition B.5) vanishes on  $\mathbf{H}^{2d}(\mathcal{X}_{\acute{e}t}, F_! F^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}})^\heartsuit$ .

*Proof.* When  $d = 0$ , the natural map  $\mathbf{H}_{\text{rig}}^0((U, X)/W^\circ) \rightarrow \mathbf{H}_{\text{rig}}^0(X/W^\circ)$  is injective. When  $d \geq 1$ , since  $\mathcal{U}$  is  $d$ -dense, by Lemma B.6(1) and the long exact sequence (B.11), the natural map  $\mathbf{H}_{\text{rig}}^{2d}((U, X)/W^\circ) \rightarrow \mathbf{H}_{\text{rig}}^{2d}(X/W^\circ)$  is an isomorphism, in particular, injective as well.

Thus, in both cases, we may assume  $\mathcal{U} = \mathcal{X}$ . Then the map  $\mathbf{H}^{2d}(\mathcal{X}_{\acute{e}t}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}}) \rightarrow \mathbf{H}_{\text{rig}}^{2d}(X/W^\circ)$  factors through  $\mathbf{H}^{2d}(\mathcal{X}_{\acute{e}t}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}}) \rightarrow \mathbf{H}^{2d}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{X}/W[l]^\circ})$  by (B.16), which vanishes on  $\mathbf{H}^{2d}(\mathcal{X}_{\acute{e}t}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}})^\heartsuit$  by [Sat13, Lemma 9.5 & Proposition A.3.1].<sup>24</sup>  $\square$

The long exact sequence induced by (B.2) gives an isomorphism

$$\frac{\mathbf{H}_{\text{rig}}^{q-1}((U, X)/W^\circ)}{N\mathbf{H}_{\text{rig}}^{q-1}((U, X)/W^\circ)} \xrightarrow{\sim} \text{Ker}(\mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \tilde{\omega}_{(U, X)}) \rightarrow \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \omega_{(U, X)})).$$

Thus, by Lemma B.9, we obtain a map

$$(B.18) \quad \rho_{2d}: \mathbf{H}^{2d}(\mathcal{X}_{\acute{e}t}, F_! F^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}})^\heartsuit \rightarrow \frac{\mathbf{H}_{\text{rig}}^{2d-1}((U, X)/W^\circ)}{N\mathbf{H}_{\text{rig}}^{2d-1}((U, X)/W^\circ)}.$$

For every finite extension  $K^\dagger/K$  contained in  $\overline{K}$  and every object  $V \in \mathbf{M}_K(\mathbb{Q}_p)$ , we denote by

$$\text{res}_{K^\dagger}: \mathbf{H}^1(K, V) \rightarrow \mathbf{H}^1(K^\dagger, V)$$

the restriction map.

The following proposition is the key to the proof of Theorem A.6, whose proof will be given in §B.6.

**Proposition B.10.** *Suppose that  $n < p$ ,  $1 \leq d < p - 1$ , and  $\mathcal{U}$  is  $d$ -dense. There exists a finite extension  $K_U/K$  (depending on  $U$ ) contained in  $\overline{K}$  such that*

$$\text{Ker}(\rho_{2d}) \subseteq \text{Ker}(\text{res}_{K_U} \circ \alpha_{2d})$$

*holds. Moreover, we may take  $K_X$  to be  $K$ .*

Now we bring the  $\mathbb{L}$ -ring  $\mathbb{T}$  of étale correspondences on  $\mathcal{X}$  in Theorem A.6. In what follows, we write  $V_{\mathbb{L}} := V \otimes_{\mathbb{Q}_p} \mathbb{L}$  for a  $\mathbb{Q}_p$ -vector space  $V$ . In the situation of Theorem A.6, we may assume  $1 \leq d \leq n - 1$  without loss of generality.

For every  $t \in \mathbb{T}$ , put

$$\mathcal{U}_t := \mathcal{X} \setminus ((\mathcal{X} \setminus \mathcal{U})^{t^\vee}), \quad \mathcal{F}_t: \mathcal{U}_t \rightarrow \mathcal{X}$$

<sup>24</sup>In fact, we will prove a similar statement in a more general context in §B.6 (when  $n < p$ ), so the readers can just admit this claim for now.

(see Notation A.1), where  $t^\vee$  denotes the transpose of  $t$ . Then we have  $(\mathcal{U}_t)^t \subseteq \mathcal{U}$ . The element  $t$  acts on various cohomology and spectral sequences compatibly,<sup>25</sup> giving a commutative diagram

(B.19)

$$\begin{array}{ccccc}
 \mathbf{H}_{\text{rig}}^{2d-1}((U_t, X)/W^\circ)_{\mathbb{L}} & \xrightarrow{t^*} & \mathbf{H}_{\text{rig}}^{2d-1}((U, X)/W^\circ)_{\mathbb{L}} & & \\
 \downarrow N & & \downarrow N & & \\
 \mathbf{H}_{\text{rig}}^{2d-1}((U_t, X)/W^\circ)_{\mathbb{L}} & \xrightarrow{t^*} & \mathbf{H}_{\text{rig}}^{2d-1}((U, X)/W^\circ)_{\mathbb{L}} & & \\
 & \searrow & & \searrow & \\
 & & \mathbf{H}^{2d-1}(X_{\text{ét}}, \omega_{(U_t, X)}^+)_{\mathbb{L}} & \xrightarrow{t^*} & \mathbf{H}^{2d-1}(X_{\text{ét}}, \omega_{(U, X)}^+)_{\mathbb{L}} \\
 & \nearrow & & \nearrow & \\
 \mathbf{H}^{2d}(X_{\text{ét}}, \tilde{\omega}_{(U_t, X)})_{\mathbb{L}} & \xrightarrow{t^*} & \mathbf{H}^{2d}(X_{\text{ét}}, \tilde{\omega}_{(U, X)})_{\mathbb{L}} & & \\
 \uparrow (B.17) & & \uparrow (B.17) & & \\
 \mathbf{H}^{2d}(X_{\text{ét}}, F_t! F_t^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}})_{\mathbb{L}} & \xrightarrow{t^*} & \mathbf{H}^{2d}(X_{\text{ét}}, F! F^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}})_{\mathbb{L}} & & \\
 \downarrow & & \downarrow & & \\
 \mathbf{H}_{\text{rig}}^{2d}((U_t, X)/W^\circ)_{\mathbb{L}} & \xrightarrow{t^*} & \mathbf{H}_{\text{rig}}^{2d}((U, X)/W^\circ)_{\mathbb{L}} & & 
 \end{array}$$

in which the two triangles are induced from (B.8).

Let  $\mathbb{I} \subseteq \mathbb{T}$  be the annihilator of

$$\bigoplus_{h>1, q \geq 0} \mathbf{H}_{\text{cris}}^q(X^{(h)}/W) \otimes_{\mathbb{Z}_p} \mathbb{L} = \bigoplus_{h>1, q \geq 0} \mathbf{H}_{\text{rig}}^q(X^{(h)}/K_0)_{\mathbb{L}}.$$

**Lemma B.11.** *Suppose that  $\mathcal{U}$  is  $d$ -dense. Then for every  $t \in \mathbb{I}^{4n-5}$ , the kernel of the map*

$$\frac{\mathbf{H}_{\text{rig}}^{2d-1}((U_t, X)/W^\circ)_{\mathbb{L}}}{\mathbf{NH}_{\text{rig}}^{2d-1}((U_t, X)/W^\circ)_{\mathbb{L}}} \rightarrow \mathbf{H}^{2d-1}(X_{\text{ét}}, \omega_{(U_t, X)}^+)_{\mathbb{L}}$$

is annihilated by  $t^*$ .

*Proof.* Since  $\mathcal{U}$  is  $d$ -dense,  $\mathcal{U}_t$  is  $d$ -dense as well. Let

$$0 = F^{-1} \subseteq F^0 \subseteq \dots \subseteq F^{4d-2} = \mathbf{H}_{\text{rig}}^{2d-1}((U_t, X)/W^\circ)$$

be the filtration induced by the spectral sequence  $\mathbf{E}_c(U_t)_1^{-k, q+k}$ . Let  $V$  be the kernel of the map in the lemma. For every  $i$ , we regard  $F_{\mathbb{L}}^i \cap V$  as the intersection of  $V$  and the image of  $F_{\mathbb{L}}^i$  in the target of the map in the lemma.

By Lemma B.6(2,3) and Lemma B.7, we know that  $F_{\mathbb{L}}^i \cap V/F_{\mathbb{L}}^{i-1} \cap V$  is a subquotient of  $\bigoplus_{h>1, q} \mathbf{H}_{\text{rig}}^q(X^{(h)}/K_0)_{\mathbb{L}}$  for every  $0 \leq i \leq 4d-2$ . Thus, every element in  $\mathbb{I}$  annihilates  $F_{\mathbb{L}}^i \cap V/F_{\mathbb{L}}^{i-1} \cap V$  for  $0 \leq i \leq 4d-2$ , which implies that  $t^*$  annihilates  $V$ .  $\square$

**Proposition B.12.** *Suppose that  $n < p$  and  $\mathcal{U}$  is  $d$ -dense. Let  $t$  be an element in  $\mathbb{I}^{4n-5}$ . Then for every  $c \in \mathbf{Z}^d(X)_{\mathbb{L}}^0$  such that the Zariski closure of its support in  $X$  is contained in  $\mathcal{U}_t$ , we have*

$$\text{res}_{K_U}(t^* \beta_c) \in \mathbf{H}_f^1(K_U, \mathbf{H}_c^{2d-1}(\overline{U}, \mathbb{L}(d))),$$

where  $\beta_c \in \mathbf{H}_{\text{ct}}^1(K, \mathbf{H}_c^{2d-1}(\overline{U}_t, \mathbb{L}(d)))$  is the image of the cycle class of  $c$  in  $\mathbf{H}_c^{2d}(U_t, \mathbb{L}(d))^0$  under the edge map, and  $K_U/K$  is the finite extension in Proposition B.10.

*Proof.* Let  $\{\mathcal{T}_l(d)_{\mathcal{X}}\}_{l \geq 1}$  be the projective system of objects in  $\mathbf{D}^+(X_{\text{ét}}, \mathbb{Z}_p)$  defined in [Sat07, Definition 4.2.4], which fits into a distinguished triangle

$$(B.20) \quad v_{l, X}^{d-1}[-d-1] \rightarrow \mathcal{T}_l(d)_{\mathcal{X}} \rightarrow \tau^{\leq d} i^* \mathbf{R}j_* \mu_{p^l}^{\otimes d} \xrightarrow{+1} v_{l, X}^{d-1}[-d]$$

where  $\{v_{l, X}^{d-1}\}_{l \geq 1}$  is the logarithmic Hodge–Witt sheaves on  $X_{\text{ét}}$  defined in [Sat07, §2.2].

<sup>25</sup>We note that for a finite étale morphism  $f: X_0 \rightarrow X$ , one can choose admissible embedding systems  $\{(X_0^*, L_0^*) \hookrightarrow (Z_0^*, N_0^*)\}$  and  $\{(X^*, L^*) \hookrightarrow (Z^*, N^*)\}$  for  $(X_0, L_0)/W[t]^\circ$  and  $(X, L)/W[t]^\circ$ , respectively, with an étale morphism  $(Z_0^*, N_0^*) \rightarrow (Z^*, N^*)$  that is compatible with  $f$ . We have the similar statement for embedding systems for  $(X_0, L_{X_0}^0)/W[t]^\circ$  and  $(X, L_X^0)/W[t]^\circ$ .

Let  $C$  be the Zariski closure of  $c$  in  $\mathcal{X}$ . Let  $\{\mathcal{H}_i: C_i \rightarrow \mathcal{X}\}$  be the (finite) set of irreducible components of  $\text{supp } C$ , which are projective flat schemes over  $O_K$  of pure (absolute) dimension  $n-d$ . The construction of the refined cycle class of  $C_i$  in [Sat07, Definition 5.1.2] induces a map

$$\bigoplus_i (\mathbb{Z}_p/p^l)_{C_i} \rightarrow H_i^! \mathcal{T}(d)_{\mathcal{X}}[2d],$$

and hence

$$\bigoplus_i (\mathbb{Q}_p)_{C_i} \rightarrow H_i^! \mathcal{T}_{\mathbb{Q}}(d)_{\mathcal{X}}[2d].$$

As  $\text{supp } C$  is contained in  $\mathcal{U}_t$ , the natural map  $H_i^! F_{t!} F_t^* \mathcal{T}_{\mathbb{Q}}(d)_{\mathcal{X}} \rightarrow H_i^! \mathcal{T}_{\mathbb{Q}}(d)_{\mathcal{X}}$  is an equivalence. Thus, we obtain a Gysin map

$$\bigoplus_i H^0(C_i, \mathbb{L}) \rightarrow \mathbf{H}^{2d}(\mathcal{X}_{\text{ét}}, F_{t!} F_t^* \mathcal{T}_{\mathbb{Q}}(d)_{\mathcal{X}})_{\mathbb{L}}.$$

Let  $\tau_c \in \mathbf{H}^{2d}(\mathcal{X}_{\text{ét}}, F_{t!} F_t^* \mathcal{T}_{\mathbb{Q}}(d)_{\mathcal{X}})_{\mathbb{L}}$  be the image of the cycle  $C$  under the above map.

Since  $d < n < p$ , the period map induces equivalences  $\mathcal{S}_l(d)_{\mathcal{X}} \xrightarrow{\sim} \tau^{\leq d} i^* \mathbf{R}j_* \mu_{p^l}^{\otimes d}$ . Replacing  $\tau^{\leq d} i^* \mathbf{R}j_* \mu_{p^l}^{\otimes d}$  by  $\mathcal{S}_l(d)_{\mathcal{X}}$  in (B.20) and taking limit, we obtain a distinguished triangle

$$v_{\mathbb{Q}, \mathcal{X}}^{d-1}[-d-1] \rightarrow \mathcal{T}_{\mathbb{Q}}(d)_{\mathcal{X}} \rightarrow \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}} \xrightarrow{+1} v_{\mathbb{Q}, \mathcal{X}}^{d-1}[-d].$$

Denote by  $\sigma_c \in \mathbf{H}^{2d}(\mathcal{X}_{\text{ét}}, F_{t!} F_t^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}})_{\mathbb{L}}$  the image of  $\tau_c$  under the above sequence, which then belongs to  $\mathbf{H}^{2d}(\mathcal{X}_{\text{ét}}, F_{t!} F_t^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}})_{\mathbb{L}}^{\vee}$  since the cycle class of  $c$  vanishes in  $H^{2d}(\bar{X}, \mathbb{L}(d))$  and hence in  $H_c^{2d}(\bar{U}_t, \mathbb{L}(d))$ . Now we compute  $\rho_{2d}(\sigma_c)$ , where  $\rho_{2d}$  is defined in (B.18).

In the following diagram

$$\begin{array}{ccccc} F_{t!} F_t^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}} & \xrightarrow{\text{(B.16)}} & F_{t!} F_t^* W_{\mathbb{Q}} \tilde{\omega}_{\mathcal{X}}^{\bullet} & \xrightarrow{\text{(B.6)}} & \tilde{\omega}_{(\mathcal{U}, \mathcal{X})} \\ \downarrow & & \downarrow & & \downarrow \text{(B.8)} \\ F_{t!} F_t^* v_{\mathbb{Q}, \mathcal{X}}^{d-1}[-d] & \longrightarrow & F_{t!} F_t^* W_{\mathbb{Q}} \Xi_{\mathcal{X}}^{\bullet}[-1] & \xrightarrow{\text{(B.9)}} & \omega_{(\mathcal{U}, \mathcal{X})}^+[-1] \end{array}$$

in  $\mathbf{D}^+(\mathcal{X}_{\text{ét}}, \mathbb{Q}_p)$ , the left square commutes as shown in the proof of [Sat13, Proposition 9.10] (followed by taking limit and applying  $F_{t!} F_t^*$ ), and the right square is (B.10). It follows that

$$\rho_{2d}(\sigma_c) \in \ker \left( \frac{H_{\text{rig}}^{2d-1}((\mathcal{U}_t, \mathcal{X})/W^{\circ})_{\mathbb{L}}}{NH_{\text{rig}}^{2d-1}((\mathcal{U}_t, \mathcal{X})/W^{\circ})_{\mathbb{L}}} \rightarrow \mathbf{H}^{2d-1}(\mathcal{X}_{\text{ét}}, \omega_{(\mathcal{U}, \mathcal{X})}^+)_{\mathbb{L}} \right).$$

By Lemma B.11 and the diagram (B.19), we have  $\rho_{2d}(t^* \sigma_c) = t^* \rho_{2d}(\sigma_c) = 0$ . Finally, by Proposition B.10, we have  $\text{res}_{K_U}(\alpha_{2d}(t^* \sigma_c)) = 0$ . Since  $\text{res}_{K_U}(\alpha_{2d}(t^* \sigma_c))$  coincides with the image of  $\text{res}_{K_U}(t^* \beta_c)$  under the map

$$H^1(K_U, H_c^{2d-1}(\bar{U}, \mathbb{L}(d))) \rightarrow H^1(K_U, H_c^{2d-1}(\bar{U}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\text{cris}})_{\mathbb{L}},$$

$\text{res}_{K_U}(t^* \beta_c)$  belongs to  $H_f^1(K_U, H_c^{2d-1}(\bar{U}, \mathbb{L}(d)))$  in view of Remark A.2(2,3). The proposition is proved.  $\square$

*Proof of Theorem A.6.* We consider the localized cohomology  $H^{2d'-1}(\bar{X}, \mathbb{L}(d'))_{m'}$ , which is a direct summand of  $H^{2d'-1}(\bar{X}, \mathbb{L}(d'))$  in the category  $\mathbf{M}_K(\mathbb{L})$ . By the  $C_{\text{st}}$ -comparison theorem for  $\mathcal{X}$ , Mokrane's weight spectral sequence [Mok93] and (A.2), we know that  $H^{2d'-1}(\bar{X}, \mathbb{L}(d'))_{m'}$  is either zero or a semistable representation of  $\text{Gal}(\bar{K}/K)$  pure of weight  $-1$  (Definition A.8). In particular, [Nek93, Proposition 1.25] implies the following

(\*) For every short exact sequence  $0 \rightarrow \mathbb{L}(1) \rightarrow E \rightarrow H^{2d-1}(\bar{X}, \mathbb{L}(d)) \rightarrow 0$  such that  $[E^{\vee}(1)]$  belongs to  $H_f^1(K, H^{2d'-1}(\bar{X}, \mathbb{L}(d'))) \cap H_{\text{ct}}^1(K, H^{2d'-1}(\bar{X}, \mathbb{L}(d'))_{m'})$ , we have a short exact sequence

$$0 \rightarrow H_f^1(K^{\dagger}, \mathbb{L}(1)) \rightarrow H_f^1(K^{\dagger}, E) \rightarrow H_f^1(K^{\dagger}, H^{2d-1}(\bar{X}, \mathbb{L}(d))) \rightarrow 0$$

for every finite extension  $K^{\dagger}/K$  contained in  $\bar{K}$ .



Let  $\mathbb{J}' \subseteq \mathbb{T}$  be the annihilator of

$$\bigoplus_{d'=1}^{n-1} \text{Ker} \left( \text{H}_{\text{ct}}^1(K, \text{H}^{2d'-1}(\overline{X}, \mathbb{L}(d'))) \rightarrow \text{H}_{\text{ct}}^1(K, \text{H}^{2d'-1}(\overline{X}, \mathbb{L}(d'))_{m'}) \right).$$

In particular, we have  $[E_{t_1^{*c}}] \in \text{H}_{\text{ct}}^1(K, \text{H}^{2d'-1}(\overline{X}, \mathbb{L}(d'))_{m'})$  for every  $t_1' \in \mathbb{J}'$ . We need to apply Proposition B.12 three times.

First, we apply Proposition B.12 to the case  $c = t_1^{*c}$ ,  $d = d'$ , and  $\mathcal{U} = \mathcal{X}$ . Then for every  $t_2' \in \mathbb{I}^{4n-5}$ , the class  $\beta_{t_2 t_1^{*c}} = t_2^* \beta_{t_1^{*c}}$  belongs to  $\text{H}_f^1(K, \text{H}^{2d'-1}(\overline{X}, \mathbb{L}(d')))$ . In other words, we have

$$[E_{t_2 t_1^{*c}}] \in \text{H}_f^1(K, \text{H}^{2d'-1}(\overline{X}, \mathbb{L}(d'))) \cap \text{H}_{\text{ct}}^1(K, \text{H}^{2d'-1}(\overline{X}, \mathbb{L}(d'))_{m'}),$$

where  $t' := t_1' t_2'$ . By (\*), we have a short exact sequence

$$0 \rightarrow \text{H}_f^1(K^\dagger, \mathbb{L}(1)) \rightarrow \text{H}_f^1(K^\dagger, E_{t'^*c'}) \rightarrow \text{H}_f^1(K^\dagger, \text{H}^{2d-1}(\overline{X}, \mathbb{L}(d))) \rightarrow 0$$

for every finite extension  $K^\dagger/K$  contained in  $\overline{K}$ . Now we denote by  $\text{H}_\#^1(K^\dagger, E_{t'^*c'})$  the inverse image of the subspace  $\text{H}_f^1(K^\dagger, \text{H}^{2d-1}(\overline{X}, \mathbb{L}(d)))$  under the map  $\text{H}_{\text{ct}}^1(K^\dagger, E_{t'^*c'}) \rightarrow \text{H}_{\text{ct}}^1(K^\dagger, \text{H}^{2d-1}(\overline{X}, \mathbb{L}(d)))$ .<sup>26</sup> Then the diagram

$$(B.21) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \text{H}_f^1(K^\dagger, \mathbb{L}(1)) & \longrightarrow & \text{H}_f^1(K^\dagger, E_{t'^*c'}) & \longrightarrow & \text{H}_f^1(K^\dagger, \text{H}^{2d-1}(\overline{X}, \mathbb{L}(d))) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \text{H}_{\text{ct}}^1(K^\dagger, \mathbb{L}(1)) & \longrightarrow & \text{H}_\#^1(K^\dagger, E_{t'^*c'}) & \longrightarrow & \text{H}_f^1(K^\dagger, \text{H}^{2d-1}(\overline{X}, \mathbb{L}(d))) \longrightarrow 0 \end{array}$$

is a pushout of extensions.

Second, we apply Proposition B.12 to the case  $c = c$ ,  $d = d$ , and  $\mathcal{U} = \mathcal{X}$ . Then for every  $t_1 \in \mathbb{I}^{4n-5}$ , the class  $\beta_{t_1^c} = t_1^* \beta_c$  belongs to  $\text{H}_f^1(K, \text{H}^{2d-1}(\overline{X}, \mathbb{L}(d)))$ . In other words,  $[E_{t_1^c}] \in \text{H}_f^1(K, \text{H}^{2d-1}(\overline{X}, \mathbb{L}(d)))$  and hence  $[E_{t_2 t_1^c}] \in \text{H}_\#^1(K, E_{t'^*c'})$ .

Third, we apply Proposition B.12 to the case  $c = t_1^*c$ ,  $d = d$ , and  $\mathcal{U} = \mathcal{X} \setminus (\text{supp } C)'$ . Then for every  $t_2 \in \mathbb{I}^{4n-5}$ , the class  $\text{res}_{K_U}(\beta_{t_2 t_1^*c}) = \text{res}_{K_U}(t_2^* \beta_{t_1^*c})$  belongs to  $\text{H}_f^1(K_U, \text{H}_c^{2d-1}(\overline{U}_t, \mathbb{L}(d)))$ , where  $t := t_1 t_2$ . Note that the fact that  $\mathcal{U}$  is  $d$ -dense follows from condition (2), and that  $\text{supp } C \subseteq \mathcal{U}_t$  follows from condition (1). Since  $[E_{t_2 t_1^*c}]$  is the image of  $\beta_{t_2 t_1^*c}$  under the pushout map

$$\text{H}_{\text{ct}}^1(K, \text{H}_c^{2d-1}(\overline{U}_t, \mathbb{L}(d))) \rightarrow \text{H}_{\text{ct}}^1(K, E_{t'^*c'}),$$

we have  $\text{res}_{K_U}([E_{t_2 t_1^*c}]) \in \text{H}_f^1(K_U, E_{t'^*c'})$ . Since the inverse image of  $\text{H}_f^1(K_U, \mathbb{L}(1))$  under  $\text{res}_{K_U}$  coincides with  $\text{H}_f^1(K, \mathbb{L}(1))$ , we conclude that  $[E_{t_2 t_1^*c}] \in \text{H}_f^1(K, E_{t'^*c'})$  by the diagram (B.21) (for  $K^\dagger = K, K_U$ ).

From the above discussion, the conclusion of the theorem holds for every pair of elements  $t \in (\mathbb{I}^{4n-5} \setminus \mathfrak{m})^2$  and  $t' \in (\mathbb{J}' \setminus \mathfrak{m}') \cdot (\mathbb{I}^{4n-5} \setminus \mathfrak{m}')$ . It is clear that  $\mathbb{J}' \setminus \mathfrak{m}' \neq \emptyset$ . By (A.2), we also have  $\mathbb{I} \setminus \mathfrak{m} \neq \emptyset$  and  $\mathbb{I} \setminus \mathfrak{m}' \neq \emptyset$ .

The theorem is proved.  $\square$

**B.5. Further preparation.** Let  $\mathbf{A}$  be a  $\mathbb{Z}_p$ -linear additive category. Following Fontaine, we say that a  $(W, \varphi, N)$ -module or simply  $(\varphi, N)$ -module in  $\mathbf{A}$  is a (compatible)  $W$ -module object  $C$  in  $\mathbf{A}$  with a  $W$ -semi-linear endomorphism  $\varphi_C : C \rightarrow C$  (called the Frobenius operator) and a  $W$ -linear endomorphism  $N_C : C \rightarrow C$  (called the monodromy operator) satisfying that  $N_C \circ \varphi_C = p \cdot \varphi_C \circ N_C$ . A map between  $(\varphi, N)$ -modules is a  $W$ -linear map that commutes with both Frobenius operators and monodromy operators.

Suppose that  $\mathbf{A}$  admits  $\mathbb{Z}_p$ -linear tensors and quotients.

- For an object  $A$  and a  $(\varphi, N)$ -module  $C$  in  $\mathbf{A}$ , we define a  $(\varphi, N)$ -module structure on  $A \otimes_{\mathbb{Z}_p} C$  through  $C$  via linear extension.
- For two  $(\varphi, N)$ -modules  $C$  and  $D$  in  $\mathbf{A}$ , we equip  $C \otimes_W D$  with a  $(\varphi, N)$ -module structure with the obvious  $W$ -action, together with

$$\varphi_{C \otimes_W D} := \varphi_C \otimes \varphi_D, \quad N_{C \otimes_W D} := N_C \otimes 1 + 1 \otimes N_D.$$

<sup>26</sup>In fact,  $\text{H}_\#^1(K^\dagger, E_{t'^*c'}) = \text{H}_{\text{st}}^1(K^\dagger, E_{t'^*c'})$ .

When  $\mathbf{A}$  is an abelian category, we say that a  $(\varphi, N)$ -module is *nilpotent* if it has finite length and the monodromy operator is nilpotent.

**Definition B.13.** Let  $\mathbf{D}$  be a  $\mathbb{Z}_p$ -linear triangulated category. For a finite complex

$$\mathcal{C}_0 \rightarrow \mathcal{C}_1 \rightarrow \cdots \rightarrow \mathcal{C}_m$$

in  $\mathbf{D}$ , we have a canonical diagram

$$\begin{array}{ccccccc}
 \mathcal{C}_{-1} & \longrightarrow & 0 & & & & \\
 \downarrow & & \downarrow & & & & \\
 \mathcal{C}_0 & \longrightarrow & \bullet & \longrightarrow & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \mathcal{C}_1 & \longrightarrow & \cdots & & \\
 & & \downarrow & & \downarrow & & \\
 & & \cdots & \longrightarrow & \bullet & \longrightarrow & 0 \\
 & & & & \downarrow & & \\
 & & & & \mathcal{C}_{m-2} & \longrightarrow & \bullet & \longrightarrow & 0 \\
 & & & & \downarrow & & \downarrow & & \\
 & & & & & & \mathcal{C}_{m-1} & \longrightarrow & \mathcal{C}_m
 \end{array}$$

in  $\mathbf{D}$ , in which every square is a homotopy fiber. We call  $\mathcal{C}_{-1}$  the *successive homotopy fiber* of the complex  $\mathcal{C}_0 \rightarrow \mathcal{C}_1 \rightarrow \cdots \rightarrow \mathcal{C}_m$ .

If  $\mathcal{C}_0 \rightarrow \mathcal{C}_1 \rightarrow \cdots \rightarrow \mathcal{C}_m$  is a complex of  $(\varphi, N)$ -modules, then  $\mathcal{C}_{-1}$  is canonically a  $(\varphi, N)$ -module.

Next, we review some period rings. For every  $l \geq 1$ , let  $R_l$  be the PD envelop of  $O_K/p^l$  in  $W[t]/p^l$ , and let  $P_l$  be the  $R_l$ -ring in  $\mathbf{M}_K(\mathbb{Z}_p)$  defined in [Kat94, (3.2)] or [Tsu99, §1.6]. Put

$$\mathbb{K} := \left( \varprojlim_l R_l \right) \otimes \mathbb{Q} \in \mathbf{M}(\mathbb{Q}_p), \quad \widehat{\mathbb{B}}_{\text{st}}^+ := \left( \varprojlim_l P_l \right) \otimes \mathbb{Q} \in \mathbf{M}_K(\mathbb{Q}_p).$$

Both  $\mathbb{K}$  and  $\widehat{\mathbb{B}}_{\text{st}}^+$  are  $(\varphi, N)$ -modules in relevant categories. See [Bre97, §2 & §4] for an explicit description.<sup>27</sup>

**Lemma B.14.** *The following holds.*

- (1) *There is a canonical isomorphism  $\widehat{\mathbb{B}}_{\text{st}}^+ \simeq (\widehat{\mathbb{B}}_{\text{st}}^+)^{N\text{-nilp}}$ .*
- (2) *The ring  $\widehat{\mathbb{B}}_{\text{st}}^+$  is flat over  $\mathbb{K}$ .*
- (3) *The monodromy operator  $N: \widehat{\mathbb{B}}_{\text{st}}^+ \rightarrow \widehat{\mathbb{B}}_{\text{st}}^+$  is surjective.*
- (4) *We have  $\mathbb{K} = (\widehat{\mathbb{B}}_{\text{st}}^+)^{G_K}$ .*

*Proof.* For (1), this is [Tsu99, Proposition 4.1.3].

For (2), this is [Tsu99, Proposition 4.1.5].

For (3), we have  $N = \left( \varprojlim_l N_l \right) \otimes \mathbb{Q}$ , where  $N_l$  is the monodromy operator on  $P_l$  [Kat94, Definition 3.4]. Then (3) follows from the fact that  $N_l$  is surjective and  $\varprojlim_l \text{Ker } N_l = 0$  [Kat94, Corollary 3.6].

For (4), this is [Bre97, Corollaire 4.1.3]. □

For two  $(\varphi, N)$ -modules  $C$  and  $D$  in  $\mathbf{M}(\mathbb{Q}_p)$  or  $\mathbf{M}_K(\mathbb{Q}_p)$ . The  $(\varphi, N)$ -module structure on  $C \otimes_W D$  clearly descends to  $C \otimes_{K_0} D$ . Now we consider a nilpotent  $(\varphi, N)$ -module  $D$  in  $\mathbf{M}(\mathbb{Q}_p)$ . By Lemma B.14, we have a Frobenius

<sup>27</sup>Our  $\mathbb{K}$  and  $\widehat{\mathbb{B}}_{\text{st}}^+$  are denoted as  $S_{\min}$  and  $\widehat{B}_{\text{st}}^+$  in [Bre97], respectively.

equivariant diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & (D \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0} & \longrightarrow & D \otimes_{K_0} \mathbb{B}_{\text{st}}^+ & \xrightarrow{N} & D \otimes_{K_0} \mathbb{B}_{\text{st}}^+ \longrightarrow 0 \\
& & \parallel & & \downarrow & & \downarrow \\
0 & \longrightarrow & (D \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0} & \longrightarrow & D \otimes_{K_0} \widehat{\mathbb{B}}_{\text{st}}^+ & \xrightarrow{N} & D \otimes_{K_0} \widehat{\mathbb{B}}_{\text{st}}^+ \longrightarrow 0
\end{array}$$

in  $\mathbf{M}_K(K_0)$ , in which the two rows are short exact sequences. It induces a diagram

$$\begin{array}{ccc}
\frac{D}{ND} \hookrightarrow \mathbf{H}^1\left(K, (D \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0}\right) & & \\
\downarrow & & \parallel \\
\frac{D \otimes_{K_0} \mathbb{K}}{N(D \otimes_{K_0} \mathbb{K})} \hookrightarrow \mathbf{H}^1\left(K, (D \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0}\right) & & 
\end{array}$$

of edge maps as  $(\mathbb{B}_{\text{st}}^+)^{G_K} = K_0$ .

**Lemma B.15.** *For every integer  $r$ , the restricted edge map*

$$\left( \frac{D \otimes_{K_0} \mathbb{K}}{N(D \otimes_{K_0} \mathbb{K})} \right)^{\varphi=p^r} \rightarrow \mathbf{H}^1\left(K, (D \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0}\right)$$

factors through the map

$$\frac{D \otimes_{K_0} \mathbb{K}}{N(D \otimes_{K_0} \mathbb{K})} \rightarrow \frac{D}{ND}$$

induced by the specialization map  $\mathbb{K} \rightarrow K_0$  at  $t = 0$ .

*Proof.* This follows from the same proof of [Lan99, Lemma 2.6].<sup>28</sup>  $\square$

**Remark B.16.** Lemma B.15 is certainly wrong without the restriction to the part  $\varphi = p^r$  since the specialization map  $\mathbb{K}/N\mathbb{K} \rightarrow K_0$  has a large kernel. However, we do not know whether the edge map  $D \otimes_{K_0} \mathbb{K} \rightarrow \mathbf{H}^1(K, (D \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0})$  factors through the quotient  $D/ND$ , which is equivalent to the inclusion  $\text{Ker}(\mathbb{K} \rightarrow K_0) \subseteq N(\widehat{\mathbb{B}}_{\text{st}})^{G_K}$  where  $\widehat{\mathbb{B}}_{\text{st}} := \widehat{\mathbb{B}}_{\text{st}}^+ \otimes_{\mathbb{B}_{\text{cris}}^+} \mathbb{B}_{\text{cris}}$ . See [Bre97, §5] for the mystery of  $(\widehat{\mathbb{B}}_{\text{st}})^{G_K}$ .

**B.6. Proof of Proposition B.10.** This subsection is devoted to the proof of Proposition B.10, for which we use de Jong's alterations. We may assume  $1 \leq d \leq n - 1$ .

By [dJ96, Theorem 6.5], we may find a finite extension  $K'$  of  $K$  (depending on  $U$ ) contained in  $\overline{K}$ , a projective strictly semistable scheme  $\mathcal{X}'$  over  $O_{K'}$  and a generically finite morphisms  $\mathcal{A}: \mathcal{X}' \rightarrow \mathcal{X}$  over  $O_K$ ,<sup>29</sup> such that  $(\mathcal{X}', A^{-1}U)$  is a strict semistable pair [dJ96, §6.3]. Let  $\mathcal{F}': \mathcal{U}' \rightarrow \mathcal{X}'$  be the open subscheme with  $U' = A^{-1}U$  and such that  $\mathcal{X}' \setminus \mathcal{U}'$  is flat over  $O_{K'}$ . Note that  $\mathcal{U}'$  may strictly contain  $\mathcal{A}^{-1}\mathcal{U}$ . We have the similar constructions for  $(\mathcal{X}', \mathcal{U}')$  and we put a *prime* for the notation.

**Lemma B.17.** *Suppose that  $n < p$ . Then for every  $q \geq 0$ , the composite map*

$$\mathbf{H}^q(\mathcal{X}'_{\text{ét}}, F'_! F'^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}'}) \rightarrow \mathbf{H}^q(\mathcal{X}'_{\text{ét}}, \widetilde{\omega}_{(\mathcal{U}', \mathcal{X}')} ) \rightarrow \mathbf{H}^q(\mathcal{X}'_{\text{ét}}, \omega_{(\mathcal{U}', \mathcal{X}')} ) = \mathbf{H}_{\text{rig}}^q((\mathcal{U}', \mathcal{X}')/W'^{\circ})$$

(Definition B.5) vanishes on  $\mathbf{H}^q(\mathcal{X}'_{\text{ét}}, F'_! F'^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}'})^{\heartsuit}$ .

Note that this lemma does not follow from Lemma B.9 even for  $q = 2d$ , since  $\mathcal{U}'$  is not necessarily  $d$ -dense anymore. By Lemma B.17, we have the map

$$(\text{B.22}) \quad \rho'_q: \mathbf{H}^q(\mathcal{X}'_{\text{ét}}, F'_! F'^* \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}'})^{\heartsuit} \rightarrow \frac{\mathbf{H}_{\text{rig}}^{q-1}((\mathcal{U}', \mathcal{X}')/W'^{\circ})}{N\mathbf{H}_{\text{rig}}^{q-1}((\mathcal{U}', \mathcal{X}')/W'^{\circ})}$$

<sup>28</sup>In [Lan99], the author works with  $\widehat{K}\langle t \rangle$ , which is different from our  $\mathbb{K}$  when  $K/\mathbb{Q}_p$  is ramified. However, such difference will not affect the proof in view of the explicit description of  $\mathbb{K}$  in [Bre97, §4].

<sup>29</sup>The letter  $\mathcal{A}$  stands for *alteration*.

similar to (B.18) for every  $q \geq 0$ .

**Lemma B.18.** *Suppose that  $n < p$ . Then*

$$\mathrm{Ker}(\rho'_q) \subseteq \mathrm{Ker}(\alpha'_q)$$

holds for every  $q \geq 0$ .

We now prove Proposition B.10 assuming the above two lemmas, whose proofs are postponed later.

*Proof of Proposition B.10.* Take  $K_U = K'$ . It is also clear that we may take  $K_X$  to be  $K$ . We have the commutative diagram

$$\begin{array}{ccc} \frac{H_{\mathrm{rig}}^{2d-1}((U', X')/W'^{\circ})}{NH_{\mathrm{rig}}^{2d-1}((U', X')/W'^{\circ})} & \longleftarrow \mathbf{H}^{2d}(X'_{\acute{\mathrm{e}}\mathrm{t}}, F'_! F'^* \mathcal{S}_{\mathbb{Q}}(d)_{X'})^{\heartsuit} & \longrightarrow H^1(K', H_c^{2d-1}(\overline{U}', \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\mathrm{cris}}) \\ & \uparrow A^* & \uparrow A^* \\ \frac{H_{\mathrm{rig}}^{2d-1}((U, X)/W^{\circ})}{NH_{\mathrm{rig}}^{2d-1}((U, X)/W^{\circ})} & \longleftarrow \mathbf{H}^{2d}(X_{\acute{\mathrm{e}}\mathrm{t}}, F_! F^* \mathcal{S}_{\mathbb{Q}}(d)_X)^{\heartsuit} & \longrightarrow H^1(K, H_c^{2d-1}(\overline{U}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\mathrm{cris}}) \end{array}$$

in  $\mathbf{M}(\mathbb{Q}_p)$ .<sup>30</sup> By Lemma B.18, to prove the proposition, it suffices to show that the map

$$H^1(K', H_c^{2d-1}(\overline{U}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\mathrm{cris}}) \rightarrow H^1(K', H_c^{2d-1}(\overline{U}', \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\mathrm{cris}})$$

is injective. However, this follows from the fact that the map  $H_c^q(\overline{U}, \mathbb{Q}_p) \rightarrow H_c^q(\overline{U}', \mathbb{Q}_p)$  in the category  $\mathbf{M}_{K'}(\mathbb{Q}_p)$  admits a section, which is a consequence of the usual Poincaré duality for étale cohomology of  $\overline{U}$  and  $\overline{U}'$ . The proposition is proved.  $\square$

It remains to show Lemma B.17 and Lemma B.18. Since we will only study  $(X', \mathcal{U}')$  from now on, we will suppress the *prime* from all notation to release some burden. Put  $\mathcal{V} := X \setminus \mathcal{U}$ , and for every  $h \geq 1$ , let  $\mathcal{V}^{(h)}$  be the disjoint union of intersections of  $h$  different irreducible components of  $\mathcal{V}$ , which is either empty or a strictly semistable scheme over  $O_K$  of pure (absolute) dimension  $n - h$ . For notational convenience, we also put  $\mathcal{V}^{(0)} := X$  and  $\mathcal{V}^{(-1)} := (\mathcal{U}, X)$ . Denote by  $\mathcal{G}^{(h)}: \mathcal{V}^{(h)} \rightarrow X$  the obvious morphism for  $h \geq 0$ .

**Lemma B.19.** *For every  $h \geq 0$ , the pullback of the log structure  $L_X^X$  for  $X$  to  $\mathcal{V}^{(h)}$  coincides with  $L_{\mathcal{V}^{(h)}}^{\mathcal{V}^{(h)}}$ .*

*Proof.* The question is local in the Zariski topology. By [dJ96, §6.4], Zariski locally  $X$  is smooth over  $O_K[t_1, \dots, t_i, s_1, \dots, s_j]/(t_1 \cdots t_i - \varpi)$ . We may assume  $j \geq h$  since otherwise  $\mathcal{V}^{(h)}$  is empty in this chart. It suffices to consider the open and closed subscheme  $\mathcal{T}$  of  $\mathcal{V}^{(h)}$  defined by  $s_{j-h+1} = \cdots = s_j = 0$ . Now locally  $L_X^X$  and  $L_{\mathcal{T}}^{\mathcal{T}}$  are the log structures associated with the pre-log structures  $\mathbb{N}^i \rightarrow \mathcal{O}_X$  and  $\mathbb{N}^i \rightarrow \mathcal{O}_{\mathcal{T}}$  sending 1 in the  $i'$ -th factor to the pullback of  $t_{i'}$  for  $1 \leq i' \leq i$ , respectively. The lemma follows immediately.  $\square$

Our first step is to construct, for every  $r \geq 0$ , a projective system of syntomic complexes  $\{\mathcal{S}_l(r)_{\overline{\mathcal{V}^{(-1)}}}\}_{l \geq 1}$  in  $\mathbf{D}_K^+(\overline{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathbb{Z}_p)$ , together with a period map

$$(B.23) \quad \mathcal{S}_l(r)_{\overline{\mathcal{V}^{(-1)}}} \rightarrow \bar{i}^* \mathbf{R} \bar{j}_* \bar{F}_! \mu_{p^l}^{\otimes r}$$

when  $0 \leq r < p - 1$ , that becomes an equivalence when  $n - 1 \leq r < p - 1$ . The construction is inspired by the observation in the following remark.

*Remark B.20.* After choosing an order on the (finite) set of irreducible components of  $\mathcal{V}$ , we have an exact sequence

$$0 \rightarrow \bar{F}_! \mu_{p^l}^{\otimes r} \rightarrow \overline{G^{(0)}}_* \mu_{p^l}^{\otimes r} \rightarrow \overline{G^{(1)}}_* \mu_{p^l}^{\otimes r} \rightarrow \overline{G^{(2)}}_* \mu_{p^l}^{\otimes r} \rightarrow \cdots \rightarrow \overline{G^{(n-1)}}_* \mu_{p^l}^{\otimes r} \rightarrow 0$$

<sup>30</sup>Note that  $U'$  may properly contain  $A^{-1}U$ . By  $A^*$ , we mean the restriction map with a possible composition of the pushforward map along the inclusion  $A^{-1}U \subseteq U'(\subseteq X')$  of log rigid cohomology (Definition B.5) or étale cohomology with proper support.

in  $\mathbf{M}_K(\overline{X}_{\acute{e}t}, \mathbb{Z}_p)$ , compatible in  $l$ . In particular,  $\overline{F}_! \mu_{p^l}^{\otimes r}$  is canonically equivalent to the successive homotopy fiber of the complex

$$\overline{G}^{(0)} *_\mu_{p^l}^{\otimes r} \rightarrow \overline{G}^{(1)} *_\mu_{p^l}^{\otimes r} \rightarrow \overline{G}^{(2)} *_\mu_{p^l}^{\otimes r} \rightarrow \cdots \rightarrow \overline{G}^{(n-1)} *_\mu_{p^l}^{\otimes r}.$$

In order to unify the discussion, we put

$$\mathcal{N}_l(r)_{\overline{\mathcal{V}^{(-1)}}} := \overline{i}^* \mathbf{R} \overline{j}_* \overline{F}_! \mu_{p^l}^{\otimes r} \in \mathbf{D}_K^+(\overline{X}_{\acute{e}t}, \mathbb{Z}_p), \quad \mathcal{N}_l(r)_{\mathcal{V}^{(-1)}} := i^* \mathbf{R} j_* F_! \mu_{p^l}^{\otimes r} \in \mathbf{D}^+(X_{\acute{e}t}, \mathbb{Z}_p),$$

and

$$\mathcal{N}_l(r)_{\overline{\mathcal{V}^{(h)}}} := \overline{i}^* \mathbf{R} \overline{j}_* \overline{G}^{(h)} *_\mu_{p^l}^{\otimes r} \in \mathbf{D}_K^+(\overline{X}_{\acute{e}t}, \mathbb{Z}_p), \quad \mathcal{N}_l(r)_{\mathcal{V}^{(h)}} := i^* \mathbf{R} j_* G_*^{(h)} \mu_{p^l}^{\otimes r} \in \mathbf{D}^+(X_{\acute{e}t}, \mathbb{Z}_p),$$

for  $h \geq 0$ .<sup>31</sup>

To define  $\mathcal{S}_l(r)_{\overline{\mathcal{V}^{(-1)}}$ , we need to consider all extensions of  $K$  in  $\overline{K}$ . A *Galois embedding system* for  $O_{\overline{K}}^{\text{can}}/W[t]^\circ$  consists of

- an increasing tower  $K = K_1 \subseteq K_2 \subseteq K_3 \subseteq \cdots$  of finite Galois extensions of  $K$  with  $\bigcup_m K_m = \overline{K}$  (and we regard  $O_{K_m}^{\text{can}}$  as a log-scheme over  $O_K^{\text{can}}$ ),
- for every  $m \geq 1$ , an embedding system  $\{O_{K_m}^{\text{can}} \hookrightarrow (\mathcal{Z}_m^b, N_m^b)\}$  for  $O_{K_m}^{\text{can}}/W[t]^\circ$  with a compatible action of  $\text{Gal}(K_m/K)$  that fits into a commutative diagram

$$\begin{array}{ccc} \vdots & & \vdots \\ \downarrow & & \downarrow \\ O_{K_{m+1}}^{\text{can}} & \hookrightarrow & (\mathcal{Z}_{m+1}^b, N_{m+1}^b) \\ \downarrow & & \downarrow \\ O_{K_m}^{\text{can}} & \hookrightarrow & (\mathcal{Z}_m^b, N_m^b) \\ \downarrow & & \downarrow \\ \vdots & & \vdots \end{array}$$

of log-schemes over  $W[t]^\circ$  that is  $G_K$ -equivariant.

It is clear that Galois embedding system for  $O_{\overline{K}}^{\text{can}}/W[t]^\circ$  exists.

We now choose a Galois embedding system for  $O_{\overline{K}}^{\text{can}}/W[t]^\circ$  as above, and write  $\kappa = \kappa_1 \subseteq \kappa_2 \subseteq \kappa_3 \subseteq \cdots$  for the induced tower of residue fields. We also choose an embedding system  $\{(\mathcal{X}^\star, L^\star) \hookrightarrow (\mathcal{Z}^\star, N^\star)\}$  for  $(\mathcal{X}, L_{\mathcal{X}}^\star)/W[t]^\circ$ . For  $m \geq 1$ , put

$$(\mathcal{Z}_m^\star, N_m^\star) := (\mathcal{Z}^\star, N^\star) \times_{W[t]^\circ} (\mathcal{Z}_m^b, N_m^b).$$

Let  $\mathcal{T}$  be an irreducible component of  $\mathcal{V}^{(h)}$  for some  $h \geq 0$ . Lemma B.19 implies that

$$\left\{ (\mathcal{T}, L_{\mathcal{T}}^\top) \times_{(\mathcal{X}, L_{\mathcal{X}}^\star)} (\mathcal{X}^\star, L^\star) \hookrightarrow (\mathcal{Z}^\star, N^\star) \right\}$$

is an embedding system for  $(\mathcal{T}, L_{\mathcal{T}}^\top)/W[t]^\circ$ . For every  $m \geq 1$ , let  $\mathcal{E}_{m,l}^\star$  be the PD envelop of  $\mathcal{T}^\star \otimes_{O_K} O_{K_m}/p^l$  in  $\mathcal{Z}_m^\star \otimes \mathbb{Z}/p^l$ . For  $i \geq 1$ , let  $\mathcal{J}^{[i]} \subseteq \mathcal{O}_{\mathcal{E}_{m,l}^\star}$  be the  $i$ -th divided power of the ideal  $\mathcal{J} := \text{Ker}(\mathcal{O}_{\mathcal{E}_{m,l}^\star} \rightarrow \mathcal{O}_{\mathcal{T}^\star \otimes_{O_K} O_{K_m}/p^l})$ . For  $i \leq 0$ , we put  $\mathcal{J}^{[i]} := \mathcal{O}_{\mathcal{E}_{m,l}^\star}$ . We have the complex

$$(B.24) \quad B_l(r)_{\mathcal{T},m} : \mathcal{J}^{[r]} \rightarrow \mathcal{J}^{[r-1]} \otimes_{\mathcal{O}_{\mathcal{Z}_m^\star}} \Omega_{(\mathcal{Z}_m^\star, N_m^\star)/W^{\text{triv}}}^1 \rightarrow \mathcal{J}^{[r-2]} \otimes_{\mathcal{O}_{\mathcal{Z}_m^\star}} \Omega_{(\mathcal{Z}_m^\star, N_m^\star)/W^{\text{triv}}}^2 \rightarrow \cdots$$

in  $\mathbf{M}_K((\mathbb{T} \times_{\mathcal{X}} \mathcal{X}^\star \otimes_{\kappa} \kappa_m)_{\acute{e}t}, \mathbb{Z}_p)$ , where  $\mathcal{J}^{[r]}$  is put in degree 0. Put

$$(B.25) \quad B_l(r)_{\overline{\mathcal{T}}} := \varinjlim_m B_l(r)_{\mathcal{T},m}|_{\mathbb{T} \times_{\mathcal{X}} \mathcal{X}^\star},$$

<sup>31</sup>The letter  $\mathcal{N}$  stands for *nearby*.

where the colimit is taken in the abelian category  $\mathbf{C}_K^+(\overline{\mathbb{T}} \times_X X^*)_{\acute{e}t, \mathbb{Z}_p}$ . For every  $h \geq 0$ , we put

$$B_l(r)_{\overline{\mathcal{V}^{(h)}}} := \bigoplus_{\mathcal{T}} B_l(r)_{\overline{\mathcal{T}}},$$

where the direct sum is taken over all irreducible components of  $\mathcal{V}^{(h)}$ , regarded as an element in  $\mathbf{C}_K^+(\overline{X^*}_{\acute{e}t}, \mathbb{Z}_p)$  via pushforward along closed immersions  $\mathcal{T} \rightarrow X$ . Then parallel to Remark B.20, we have a complex

$$B_l(r)_{\overline{\mathcal{V}^{(0)}}} \rightarrow B_l(r)_{\overline{\mathcal{V}^{(1)}}} \rightarrow B_l(r)_{\overline{\mathcal{V}^{(2)}}} \rightarrow \cdots \rightarrow B_l(r)_{\overline{\mathcal{V}^{(n-1)}}}$$

in  $\mathbf{C}_K^+(\overline{X^*}_{\acute{e}t}, \mathbb{Z}_p)$ .

Take  $h \geq 0$ . Put  $C_{l, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}} := B_l(0)_{\overline{\mathcal{V}^{(h)}}}$ . Then we have the canonical map  $B_l(r)_{\overline{\mathcal{V}^{(h)}}} \rightarrow C_{l, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}$  given by the inclusion  $\mathcal{I}^{[s]} \rightarrow \mathcal{I}^{[0]}$ . We also have the crystalline complex  $C_{l, \overline{\mathcal{V}^{(h)}}/W[t]^\circ}$ , which is obtained in the same way as  $C_{l, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}$  except that in  $B_l(0)_{\mathcal{T}, m}$  (B.24), we replace  $\Omega_{(\mathcal{Z}_m^*, N_m^*)/W^{\text{triv}}}^q$  by  $\Omega_{(\mathcal{Z}_m^*, N_m^*)/W[t]^\circ}^q$ . We have natural Frobenius operators on both  $C_{l, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}$  and  $C_{l, \overline{\mathcal{V}^{(h)}}/W[t]^\circ}$ , and a distinguished triangle

$$(B.26) \quad C_{l, \overline{\mathcal{V}^{(h)}}/W[t]^\circ}^\Delta : \quad C_{l, \overline{\mathcal{V}^{(h)}}/W[t]^\circ}[-1] \rightarrow C_{l, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}} \rightarrow C_{l, \overline{\mathcal{V}^{(h)}}/W[t]^\circ} \xrightarrow{N} C_{l, \overline{\mathcal{V}^{(h)}}/W[t]^\circ}$$

in  $\mathbf{D}_K^+(\overline{X^*}_{\acute{e}t}, \mathbb{Z}_p)$ , where the first arrow is given by  $\wedge d \log t$ , the second arrow is the canonical one (which is Frobenius equivariant), and the third arrow is the connecting map, such that  $C_{l, \overline{\mathcal{V}^{(h)}}/W[t]^\circ}$  becomes a  $(\varphi, N)$ -module in  $\mathbf{D}_K^+(\overline{X^*}_{\acute{e}t}, \mathbb{Z}_p)$ . Let  $B_l(r)_{\overline{\mathcal{V}^{(l-1)}}$  be the successive homotopy fiber (Definition B.13) of the complex

$$B_l(r)_{\overline{\mathcal{V}^{(0)}}} \rightarrow B_l(r)_{\overline{\mathcal{V}^{(1)}}} \rightarrow B_l(r)_{\overline{\mathcal{V}^{(2)}}} \rightarrow \cdots \rightarrow B_l(r)_{\overline{\mathcal{V}^{(n-1)}}$$

in  $\mathbf{D}_K^+(\overline{X^*}_{\acute{e}t}, \mathbb{Z}_p)$ , and similarly for  $C_{l, \overline{\mathcal{V}^{(l-1)}}/W^{\text{triv}}}$  and  $C_{l, \overline{\mathcal{V}^{(l-1)}}/W[t]^\circ}$ . Then  $C_{l, \overline{\mathcal{V}^{(l-1)}}/W^{\text{triv}}}$  is a  $(\varphi, N)$ -module and we have a similar distinguished triangle (B.26) for  $h = -1$ . Now for every  $h \geq -1$  and every  $l \geq 1$ , we define  $S_l(r)_{\overline{\mathcal{V}^{(h)}}}$  to be the homotopy fiber of the map

$$1 - p^{-r} \varphi_{l+r} : B_l(r)_{\overline{\mathcal{V}^{(h)}}} \rightarrow C_{l, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}$$

(see [Tsu00, Page 540] for more details). Finally, put

$$\mathcal{S}_l(r)_{\overline{\mathcal{V}^{(h)}}} := \mathbf{R}\overline{U}_* S_l(r)_{\overline{\mathcal{V}^{(h)}}}, \quad \mathcal{C}_{l, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}} := \mathbf{R}\overline{U}_* C_{l, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}, \quad \mathcal{C}_{l, \overline{\mathcal{V}^{(h)}}/W[t]^\circ} := \mathbf{R}\overline{U}_* C_{l, \overline{\mathcal{V}^{(h)}}/W[t]^\circ}$$

in  $\mathbf{D}_K^+(\overline{X}_{\acute{e}t}, \mathbb{Z}_p)$ , so we have the canonical map

$$(B.27) \quad \overline{\xi}_r : \mathcal{S}_l(r)_{\overline{\mathcal{V}^{(h)}}} \rightarrow \mathcal{C}_{l, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}$$

and a distinguished triangle

$$(B.28) \quad \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W[t]^\circ}^\Delta : \quad \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W[t]^\circ}[-1] \rightarrow \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}} \rightarrow \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W[t]^\circ} \xrightarrow{N} \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W[t]^\circ}$$

in  $\mathbf{D}_K^+(\overline{X}_{\acute{e}t}, \mathbb{Q}_p)$ .

When  $0 \leq r < p - 1$ , the usual period maps for  $\mathcal{V}^{(h)}$  with  $h \geq 0$  give a commutative diagram

$$\begin{array}{ccccccc} \mathcal{S}_l(r)_{\overline{\mathcal{V}^{(0)}}} & \longrightarrow & \mathcal{S}_l(r)_{\overline{\mathcal{V}^{(1)}}} & \longrightarrow & \mathcal{S}_l(r)_{\overline{\mathcal{V}^{(2)}}} & \longrightarrow & \cdots \longrightarrow \mathcal{S}_l(r)_{\overline{\mathcal{V}^{(n-1)}}} \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \overline{\mathbf{G}}^{(0)} \ast \overline{i}^{(0)*} \mathbf{R}j^{(0)} \ast \mu_{p^l}^{\otimes r} & \longrightarrow & \overline{\mathbf{G}}^{(1)} \ast \overline{i}^{(1)*} \mathbf{R}j^{(1)} \ast \mu_{p^l}^{\otimes r} & \longrightarrow & \overline{\mathbf{G}}^{(2)} \ast \overline{i}^{(2)*} \mathbf{R}j^{(2)} \ast \mu_{p^l}^{\otimes r} & \longrightarrow & \cdots \longrightarrow \overline{\mathbf{G}}^{(n-1)} \ast \overline{i}^{(n-1)*} \mathbf{R}j^{(n-1)} \ast \mu_{p^l}^{\otimes r} \end{array}$$

in  $\mathbf{D}_K^+(\overline{X}_{\acute{e}t}, \mathbb{Z}_p)$ , where  $i^{(h)} : V^{(h)} \rightarrow \mathcal{V}^{(h)}$  and  $j^{(h)} : V^{(h)} \rightarrow \mathcal{V}^{(h)}$  denote the special and generic fibers, respectively, for  $h \geq 0$ . However, since the natural map

$$\mathcal{N}_l(r)_{\overline{\mathcal{V}^{(h)}}} = \overline{i}^* \mathbf{R}\overline{j}_* \overline{\mathbf{G}}^{(h)} \ast \mu_{p^l}^{\otimes r} \rightarrow \overline{\mathbf{G}}^{(h)} \ast \overline{i}^{(h)*} \mathbf{R}j^{(h)} \ast \mu_{p^l}^{\otimes r}$$

is an equivalence for every  $h \geq 0$ , we obtain the period map

$$(B.29) \quad \overline{\pi}_r : \mathcal{S}_l(r)_{\overline{\mathcal{V}^{(h)}}} \rightarrow \mathcal{N}_l(r)_{\overline{\mathcal{V}^{(h)}}}$$

in  $\mathbf{D}_K^+(\overline{X}_{\text{ét}}, \mathbb{Z}_p)$  for every  $h \geq -1$  by Remark B.20 and the process of taking successive homotopy fibers. If  $n - 1 \leq r < p - 1$ , then (B.29) is an equivalence. The desired map (B.23) is simply (B.29) for  $h = -1$ .

To proceed, we need versions of syntomic and crystalline complexes in  $\mathbf{D}^+(X_{\text{ét}}, \mathbb{Z}_p)$  for  $\mathcal{V}^{(h)}$  rather than  $\overline{\mathcal{V}^{(h)}}$ . The construction is similar to  $\mathcal{S}_l(r)_{\overline{\mathcal{V}^{(h)}}}$  and  $\mathcal{C}_{l, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}$  but only taking  $m = 1$  without the colimit (B.25). For  $h \geq -1$ , we have

- $\mathcal{S}_l(r)_{\mathcal{V}^{(h)}}$ , which is obtained in the same way as  $\mathcal{S}_l(r)_{\overline{\mathcal{V}^{(h)}}}$  but only taking  $m = 1$ ,<sup>32</sup>
- $\mathcal{C}_{l, \mathcal{V}^{(h)}/W^{\text{triv}}}$ , which is obtained in the same way as  $\mathcal{C}_{l, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}$  but only taking  $m = 1$ ,<sup>33</sup>
- $\mathcal{C}_{l, \mathcal{V}^{(h)}/W[l]^\circ}$ , which is obtained in the same way as  $\mathcal{C}_{l, \overline{\mathcal{V}^{(h)}}/W[l]^\circ}$  but only taking  $m = 1$ ; it is a  $(\varphi, N)$ -module,
- $\widetilde{\mathcal{C}}_{l, \mathcal{V}^{(h)}/W^\circ}$  and  $\mathcal{C}_{l, \mathcal{V}^{(h)}/W^\circ}$ , which are obtained after we replace  $B_l(0)_{\mathcal{T}, 1}$  (B.24) by the following complexes

$$C_{l, \mathcal{T}/W^\circ} : \mathcal{O}_{\mathcal{D}_l^*} \rightarrow \mathcal{O}_{\mathcal{D}_l^*} \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \Omega_{(\mathcal{Z}^*, N^*)/W^\circ}^1 \rightarrow \mathcal{O}_{\mathcal{D}_l^*} \otimes_{\mathcal{O}_{\mathcal{Z}^*}} \Omega_{(\mathcal{Z}^*, N^*)/W^\circ}^2 \rightarrow \cdots$$

and

$$C_{l, \mathcal{T}/W^\circ} : \mathcal{O}_{\mathcal{D}_l^*} \rightarrow \mathcal{O}_{\mathcal{D}_l^*} \otimes_{\mathcal{O}_{\mathcal{Y}^*}} \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^1 \rightarrow \mathcal{O}_{\mathcal{D}_l^*} \otimes_{\mathcal{O}_{\mathcal{Y}^*}} \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^2 \rightarrow \cdots$$

respectively, where  $(\mathcal{Y}^*, M^*) := (\mathcal{Z}^*, N^*) \times_{W[l]^\circ} W^\circ$  as in §B.2 and  $\mathcal{D}_l^*$  denotes the PD envelop of  $\mathcal{T}$  in  $\mathcal{Y}_l^*$ .<sup>34</sup>

By construction, we have maps

$$\xi_r : \mathcal{S}_l(r)_{\mathcal{V}^{(h)}} \rightarrow \mathcal{C}_{l, \mathcal{V}^{(h)}/W^{\text{triv}}}, \quad \pi_r : \mathcal{S}_l(r)_{\mathcal{V}^{(h)}} \rightarrow \mathcal{N}_l(r)_{\mathcal{V}^{(h)}}$$

$\mathbf{D}^+(X_{\text{ét}}, \mathbb{Z}_p)$  similar to (B.27) and (B.29), and a distinguished triangle

$$(B.30) \quad \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[l]^\circ}^\Delta : \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[l]^\circ}[-1] \rightarrow \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\text{triv}}} \rightarrow \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[l]^\circ} \xrightarrow{N} \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[l]^\circ}$$

in  $\mathbf{D}^+(X_{\text{ét}}, \mathbb{Q}_p)$  similar to (B.28), without *bar*.

In order to prove Lemma B.17 and Lemma B.18, we need to connect the syntomic cohomology to the log rigid cohomology via crystalline complexes we have just constructed. By construction, we have a commutative diagram

$$(B.31) \quad \begin{array}{ccc} \mathcal{S}_{\mathbb{Q}}(r)_{\mathcal{V}^{(h)}} & \xrightarrow{\xi_r} & \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\text{triv}}} \longrightarrow \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[l]^\circ} \\ & & \downarrow \qquad \qquad \downarrow \\ & & \widetilde{\mathcal{C}}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ} \longrightarrow \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ} \end{array}$$

in  $\mathbf{D}^+(X_{\text{ét}}, \mathbb{Q}_p)$  for every  $r \geq 0$  and every  $h \geq -1$ , similar to (B.16). Now we study the cohomology of various crystalline complexes.

We start from  $\mathcal{C}_{l, \mathcal{V}^{(h)}/W^\circ}$ . Note that, as in §B.2, for  $h \geq 0$ , the complex  $\{\mathcal{C}_{l, \mathcal{V}^{(h)}/W^\circ}\}_{l \geq 1}$  is nothing but the modified de Rham–Witt complex  $\{W_l \omega_{\mathcal{V}^{(h)}}^\bullet\}_{l \geq 1}$ . By the construction and (B.5), we have

- a distinguished triangle

$$\mathcal{C}_{l, \mathcal{V}^{(h)}/W^\circ}^\Delta : \mathcal{C}_{l, \mathcal{V}^{(h)}/W^\circ}[-1] \rightarrow \widetilde{\mathcal{C}}_{l, \mathcal{V}^{(h)}/W^\circ} \rightarrow \mathcal{C}_{l, \mathcal{V}^{(h)}/W^\circ} \xrightarrow{N} \mathcal{C}_{l, \mathcal{V}^{(h)}/W^\circ}$$

in  $\mathbf{D}^+(X_{\text{ét}}, \mathbb{Z}_p)$  for every  $h \geq -1$ , similar to (B.30), so that  $\mathcal{C}_{l, \mathcal{V}^{(h)}/W^\circ}$  is a  $(\varphi, N)$ -module,

- a  $K_0$ -linear commutative diagram

$$(B.32) \quad \begin{array}{ccccccc} \omega_{\mathcal{V}^{(-1)}}^\Delta & \longrightarrow & \mathbf{G}_*^{(0)} \omega_{\mathcal{V}^{(0)}}^\Delta & \longrightarrow & \mathbf{G}_*^{(1)} \omega_{\mathcal{V}^{(1)}}^\Delta & \longrightarrow & \cdots \longrightarrow \mathbf{G}_*^{(n-1)} \omega_{\mathcal{V}^{(n-1)}}^\Delta \\ \downarrow & & \downarrow \simeq & & \downarrow \simeq & & \downarrow \simeq \\ \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(-1)}/W^\circ}^\Delta & \longrightarrow & \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(0)}/W^\circ}^\Delta & \longrightarrow & \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(1)}/W^\circ}^\Delta & \longrightarrow & \cdots \longrightarrow \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(n-1)}/W^\circ}^\Delta \end{array}$$

of distinguished triangles in  $\mathbf{D}^+(X_{\text{ét}}, \mathbb{Q}_p)$ , in which all vertical arrows starting from the second are equivalences.

<sup>32</sup>In particular,  $\mathcal{S}_l(d)_{\mathcal{V}^{(0)}}$  is nothing but the complex  $\mathcal{S}_l(d)_X$  used in the previous subsection.

<sup>33</sup>In particular,  $\mathcal{C}_{l, \mathcal{V}^{(0)}/W^{\text{triv}}}$  is nothing but the complex  $\mathcal{C}_{l, X/W^{\text{triv}}}$  used in the previous subsection.

<sup>34</sup>In particular,  $\widetilde{\mathcal{C}}_{l, \mathcal{V}^{(0)}/W^\circ}$  and  $\mathcal{C}_{l, \mathcal{V}^{(0)}/W^\circ}$  coincide with  $\widetilde{\mathcal{C}}_{l, (X, L_X^{\times} | X)/W^\circ}$  and  $\mathcal{C}_{l, (X, L_X^{\times} | X)/W^\circ}$  in §B.2, respectively.

**Lemma B.21.** *We have*

- (1) *The first vertical arrow in (B.32) is also an equivalence.*
- (2) *For every  $h \geq -1$  and every  $q \geq 0$ ,  $\mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{l, \mathcal{V}^{(h)}/W^\circ})$  is finite.*
- (3) *For every  $h \geq -1$  and every  $q \geq 0$ , the  $(\varphi, N)$ -module  $\mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ})$  in  $\mathbf{M}(\mathbb{Q}_p)$  is nilpotent.*

*In particular, for every  $h \geq -1$  and every  $q \geq 0$ , we have canonical isomorphisms*

$$\mathbf{H}_{\text{rig}}^q(\mathcal{V}^{(h)}/W^\circ) \simeq \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \simeq \left( \lim_{\leftarrow l} \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{l, \mathcal{V}^{(h)}/W^\circ}) \right) \otimes \mathbb{Q}$$

*in  $\mathbf{M}(K_0)$  which commute with monodromy operators.*

*Proof.* For (1), it suffices to show that the map  $\omega_{\mathcal{V}^{(-1)}} = \omega_{(U, X)}$  is the successive homotopy fiber of the complex

$$\mathbf{G}_*^{(0)} \omega_{\mathcal{V}^{(0)}} \rightarrow \mathbf{G}_*^{(1)} \omega_{\mathcal{V}^{(1)}} \rightarrow \cdots \rightarrow \mathbf{G}_*^{(n-1)} \omega_{\mathcal{V}^{(n-1)}}.$$

However, this follows from the easy fact that for an embedding system  $\{(X^*, L^*) \hookrightarrow (\mathcal{Y}^*, M^*)\}$  for  $(X, L_X^X|_X)/W^\circ$ , the complex

$$\begin{aligned} 0 \rightarrow \mathbb{F}_{(U^*, X^*)}^1 \left( \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^\bullet \otimes_{\mathcal{O}_{\mathcal{Y}^*}} \mathcal{O}_{|X^*|_{\mathcal{Y}^*}} \right) &\rightarrow \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^\bullet \otimes_{\mathcal{O}_{\mathcal{Y}^*}} \mathcal{O}_{|X^*|_{\mathcal{Y}^*}} \\ &\rightarrow \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^\bullet \otimes_{\mathcal{O}_{\mathcal{Y}^*}} \mathcal{O}_{|\mathcal{V}^{(1)*}|_{\mathcal{Y}^*}} \rightarrow \cdots \rightarrow \Omega_{(\mathcal{Y}^*, M^*)/W^\circ}^\bullet \otimes_{\mathcal{O}_{\mathcal{Y}^*}} \mathcal{O}_{|\mathcal{V}^{(n-1)*}|_{\mathcal{Y}^*}} \rightarrow 0 \end{aligned}$$

in  $\mathbf{C}^+(\mathfrak{Y}_{\eta, \text{qét}}^*, K_0)$  is exact. Here, for  $h \geq 1$ ,

$$\mathcal{O}_{|\mathcal{V}^{(h)*}|_{\mathcal{Y}^*}} := \bigoplus_{\mathcal{T}} \mathcal{O}_{|\Gamma^*|_{\mathcal{Y}^*}}$$

where the direct sum is taken over all irreducible components of  $\mathcal{V}^{(h)}$ .

For (2) and (3), they are already known when  $h \geq 0$ . The case for  $h = -1$  follows from the series of long exact sequences induced by the successive homotopy fiber.  $\square$

Recall that by [Tsu99, (4.5.1)], we have the canonical identification

$$\widehat{\mathbb{B}}_{\text{st}}^+ = \Gamma(\text{Spec } \bar{k}, \mathcal{C}_{\mathbb{Q}, \overline{\text{Spec } \mathcal{O}_K/W[t]^\circ}})$$

of  $(\varphi, N)$ -modules in  $\mathbf{M}_K(\mathbb{Q}_p)$ .

**Lemma B.22.** *The following holds for every  $h \geq -1$ .*

- (1) *There is a canonical isomorphism*

$$\mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[t]^\circ}) \simeq \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{K}$$

*of  $(\varphi, N)$ -modules in  $\mathbf{M}(\mathbb{Q}_p)$  for every  $q \geq 0$ .*

- (2) *The natural map*

$$\mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[t]^\circ}) \otimes_{K_0} \widehat{\mathbb{B}}_{\text{st}}^+ \rightarrow \mathbf{H}^q(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W[t]^\circ}})$$

*of  $(\varphi, N)$ -modules induced by functoriality and cup product descends to an isomorphism*

$$\mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[t]^\circ}) \otimes_{\mathbb{K}} \widehat{\mathbb{B}}_{\text{st}}^+ \xrightarrow{\sim} \mathbf{H}^q(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W[t]^\circ}})$$

*of  $(\varphi, N)$ -modules in  $\mathbf{M}_K(\mathbb{Q}_p)$  for  $q \geq 0$ .*

- (3) *The distinguished triangle  $\mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W[t]^\circ}}^\Delta$  (B.28) induces a Frobenius equivariant isomorphism*

$$\mathbf{H}^q(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W^{\text{triv}}}}) \simeq \left( \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \widehat{\mathbb{B}}_{\text{st}}^+ \right)^{N=0}$$

*in  $\mathbf{M}_K(K_0)$  for every  $q \geq 0$ .*

*Moreover, these isomorphisms are compatible with  $h$  in the obvious sense.*



*Proof.* For (1), we note that [HK94, Lemma 5.2] is applicable to  $\mathcal{V}^{(h)}$  with  $h \geq 0$ , and hence also to the case for  $h = -1$  by the series of long exact sequences induced by the successive homotopy fiber. Thus, there is a canonical isomorphism

$$\left( \varprojlim_I \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{l, \mathcal{V}^{(h)}/W[l]^\circ}) \right) \otimes \mathbb{Q} \simeq \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{K}$$

of  $(\varphi, N)$ -modules for every  $q \geq -1$ . On the other hand, [HK94, Lemma 5.2] together with Lemma B.21(2) imply that

$$\mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[l]^\circ}) \simeq \left( \varprojlim_I \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{l, \mathcal{V}^{(h)}/W[l]^\circ}) \right) \otimes \mathbb{Q}.$$

Thus, (1) follows.

For (2), it is known for  $h \geq 0$  by [Tsu99, Proposition 4.5.4]. For  $h = -1$ , it follows from the series of long exact sequences induced by successive homotopy fiber and Lemma B.14(2).

For (3), by (1), (2), and Lemma B.14(1), it suffices to show that the monodromy map

$$N: \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \widehat{\mathbb{B}}_{\text{st}}^+ \rightarrow \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \widehat{\mathbb{B}}_{\text{st}}^+$$

is surjective for every  $q \geq 0$ . However, this follows from Lemma B.21(3) and Lemma B.14(3).  $\square$

**Lemma B.23.** *Consider integers  $r$  satisfying  $n - 1 \leq r < p - 1$ . For every  $h \geq -1$  and every  $q \geq 0$ , the  $\mathbb{B}_{\text{st}}$ -linear extension of the composite map*

$$\begin{aligned} \text{(B.33)} \quad \mathbf{H}^q(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(0)_{\overline{\mathcal{V}^{(h)}}}) &\xrightarrow{\sim} \mathbf{H}^q(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(r)_{\overline{\mathcal{V}^{(h)}}}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(-r) \\ &\xrightarrow{\sim} \mathbf{H}^q(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{S}_{\mathbb{Q}}(r)_{\overline{\mathcal{V}^{(h)}}}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(-r) \\ &\rightarrow \mathbf{H}^q(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W^{\text{triv}}}}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(-r) \\ &\xrightarrow{\sim} \left( \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+ \right)^{N=0} \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(-r) \\ &\rightarrow \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{B}_{\text{st}} \end{aligned}$$

is independent of  $r$  and induces an isomorphism

$$\mathbf{H}^q(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(0)_{\overline{\mathcal{V}^{(h)}}}) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\text{st}} \simeq \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{B}_{\text{st}}$$

of  $(\varphi, N)$ -modules in  $\mathbf{M}_K(\mathbb{Q}_p)$ . Here in (B.33), the second arrow is induced by the inverse of the period map  $\overline{\pi}_r$  (B.29), the third arrow is induced by the map  $\overline{\xi}_r$  (B.27), the fourth arrow is the isomorphism from Lemma B.22(3), and the last arrow is induced by the canonical map  $\mathbb{Q}_p(-r) \hookrightarrow \mathbb{B}_{\text{st}}$ .

In particular, the above isomorphism induces a Frobenius equivariant isomorphism

$$\mathbf{H}^q(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(0)_{\overline{\mathcal{V}^{(h)}}}) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\text{cris}} \simeq \left( \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{B}_{\text{st}} \right)^{N=0}$$

in  $\mathbf{M}_K(K_0)$ .

*Proof.* For  $h \geq 0$ , the statement follows from [Tsu99, Theorem 4.10.2] (the usual  $C_{\text{st}}$ -comparison theorem for proper strictly semistable schemes) together with the compatibility properties [Tsu99, Corollaries 4.8.8 & 4.9.2] for the independence of  $r$  (which is at least  $\dim V^{(h)}$ ) of the map. The case for  $h = -1$  follows from the series of long exact sequences induced by the successive homotopy fiber.<sup>35</sup>  $\square$

**Lemma B.24.** *Suppose that  $n < p$ . For every  $h \geq -1$ , every  $q \geq 0$ , and every  $0 \leq r < p - 1$ , the following diagram*

$$\begin{array}{ccccc} \mathbf{H}^q(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{S}_{\mathbb{Q}}(r)_{\overline{\mathcal{V}^{(h)}}}) & \xrightarrow[\text{(B.27)}]{\overline{\xi}_r} & \mathbf{H}^q(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W^{\text{triv}}}}) & \xrightarrow{\sim} & \left( \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+ \right)^{N=0} \\ \downarrow \overline{\pi}_r \text{ (B.29)} & & & & \downarrow \\ \mathbf{H}^q(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(r)_{\overline{\mathcal{V}^{(h)}}}) & \xrightarrow{\mathbb{Q}_p(r) \hookrightarrow \mathbb{B}_{\text{cris}}} & \mathbf{H}^q(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(0)_{\overline{\mathcal{V}^{(h)}}}) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\text{cris}} & \xrightarrow{\sim} & \left( \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{B}_{\text{st}} \right)^{N=0} \end{array}$$

<sup>35</sup>Such an isomorphism for  $h = -1$  has already been obtained in [Yam11]. The results there are much stronger than ours and in particular they contain a  $C_{\text{dR}}$ -comparison isomorphism. Thus, the log structure on  $X$  used there is  $L_X^{\mathcal{X} \cup \mathcal{V}}$ , which makes things more complicated.

in  $\mathbf{M}_K(\mathbb{Q}_p)$  commutes, in which the equivalence in the first row is from Lemma B.22, and the equivalence in the second row is from Lemma B.23.

*Proof.* When  $h \geq 0$ , the commutativity follows from the compatibility properties [Tsu99, Corollaries 4.8.8 & 4.9.2]. The case for  $h = -1$  follows from the series of long exact sequences induced by the successive homotopy fiber.  $\square$

**Lemma B.25.** *Suppose that  $n < p$ . The map  $F_!F^*\mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}} \rightarrow \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}} = \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(0)}}$  factors through  $\mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(-1)}}$ . In particular,*

(1) *the map (B.14) factors as*

$$\mathrm{R}\Gamma(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, F_!F^*\mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}}) \rightarrow \mathrm{R}\Gamma(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(-1)}}) \rightarrow \mathrm{R}\Gamma(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{N}_{\mathbb{Q}}(d)_{\mathcal{V}^{(-1)}}) = \mathrm{R}\Gamma_c(U, \mathbb{Q}_p(d)).$$

when  $d < p - 1$ ;

(2) *the map (B.17) factors as*

$$F_!F^*\mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}} \rightarrow \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(-1)}} \rightarrow \widetilde{\mathcal{C}}_{L, \mathcal{V}^{(-1)}/W^\circ} \simeq \widetilde{\omega}_{(U, \mathcal{X})}.$$

*Proof.* Since the complex  $\mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(h)}}$  is supported on  $\mathcal{V}$  for every  $h \geq 1$ , the lemma follows by the construction (and Lemma B.21(1) for the equivalence in (2)).  $\square$

For every  $h \geq -1$ , every  $q \geq 0$ , and every  $0 \leq r < p - 1$ , put

$$(B.34) \quad \mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{S}_{\mathbb{Q}}(r)_{\mathcal{V}^{(h)}})^\heartsuit := \mathrm{Ker}\left(\mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{S}_{\mathbb{Q}}(r)_{\mathcal{V}^{(h)}}) \rightarrow \mathbf{H}^q(\overline{\mathcal{X}}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{N}_{\mathbb{Q}}(r)_{\overline{\mathcal{V}^{(h)}}})\right).$$

Now we can give a proof of Lemma B.17.

*Proof of Lemma B.17.* By Lemma B.25, the proper base change, (B.31), and Lemma B.21(1), it suffices to show that the map

$$\mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(-1)}}) \rightarrow \mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(-1)}/W[t]^\circ})$$

induced by  $\xi_d$  vanishes on  $\mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(-1)}})^\heartsuit$ , which follows from Lemma B.24 with  $h = -1$  and  $r = d$ , and the injectivity of the map

$$\mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(-1)}/W[t]^\circ}) \rightarrow \mathbf{H}^q(\overline{\mathcal{X}}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(-1)}/W[t]^\circ}})$$

by Lemma B.22.  $\square$

In order to prove Lemma B.18, we need to compare edge maps in both étale and crystalline settings. For every  $h \geq -1$ , every  $q \geq 0$ , and every  $r \geq 0$ , put

$$\begin{aligned} \mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{N}_{\mathbb{Q}}(r)_{\mathcal{V}^{(h)}})^0 &:= \mathrm{Ker}\left(\mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{N}_{\mathbb{Q}}(r)_{\mathcal{V}^{(h)}}) \rightarrow \mathbf{H}^q(\overline{\mathcal{X}}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{N}_{\mathbb{Q}}(r)_{\overline{\mathcal{V}^{(h)}}})\right), \\ \mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\mathrm{triv}}})^0 &:= \mathrm{Ker}\left(\mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\mathrm{triv}}}) \rightarrow \mathbf{H}^q(\overline{\mathcal{X}}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W^{\mathrm{triv}}}})\right). \end{aligned}$$

Suppose that  $n < p$ . Then by definition,  $\pi_d$  induces a map

$$\mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(h)}})^\heartsuit \rightarrow \mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{N}_{\mathbb{Q}}(d)_{\mathcal{V}^{(h)}})^0.$$

By Lemma B.24 and the definition of the syntomic complex,  $\xi_d$  induces a map

$$\mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(h)}})^\heartsuit \rightarrow \mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\mathrm{triv}}})^{0, \varphi=p^d}.$$

Composing with the edge maps induced by the corresponding Hochschild–Serre spectral sequences, we obtain

$$\begin{aligned} \beta_q &: \mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(h)}})^\heartsuit \rightarrow \mathrm{H}^1(K, \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{N}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}})), \\ \gamma_q &: \mathbf{H}^q(\mathcal{X}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(h)}})^\heartsuit \rightarrow \mathrm{H}^1(K, \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{\mathrm{e}}\mathrm{t}}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W^{\mathrm{triv}}}})). \end{aligned}$$

The following lemma is an ‘‘Abel–Jacobi’’ version of Lemma B.24.

**Lemma B.26.** *Suppose that  $n < p$ . For every  $h \geq -1$  and every  $q \geq 0$ , the following diagram*

$$\begin{array}{ccc} \mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{S}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}})^{\heartsuit} & \xrightarrow{\gamma_q} & \mathbf{H}^1(K, \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}})) \xrightarrow{\sim} \mathbf{H}^1\left(K, (\mathbf{H}^{q-1}(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W^{\circ}}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0}\right) \\ \downarrow \beta_q & \dashrightarrow & \downarrow \\ \mathbf{H}^1(K, \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}})) & \xrightarrow{\mathbb{Q}_p(d) \hookrightarrow \mathbb{B}_{\text{cris}}} & \mathbf{H}^1(K, \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(0)_{\overline{\mathcal{V}^{(h)}}}) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\text{cris}}) \xrightarrow{\sim} \mathbf{H}^1\left(K, (\mathbf{H}^{q-1}(\mathcal{X}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W^{\circ}}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0}\right) \end{array}$$

in  $\mathbf{M}(\mathbb{Q}_p)$  commutes, in which the equivalence in the first row is from Lemma B.22, and the equivalence in the second row is from Lemma B.23.

This lemma does not follow immediately from Lemma B.24 since  $\mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{S}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}})^{\heartsuit}$ , by our definition (B.34), is in general larger than

$$\text{Ker}\left(\mathbf{H}^q(\mathcal{X}_{\text{ét}}, \mathcal{S}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}}) \rightarrow \mathbf{H}^q(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{S}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}})\right).$$

*Proof.* Take an integer  $r$  satisfying  $(1 \leq d \leq)n-1 \leq r < p-1$  (which exists as  $n < p$ ). We identify  $\mathbf{M}_K((\text{Spec } \bar{k})_{\text{ét}}, L)$  with  $\mathbf{M}_K(L)$ . For example, we have

- the map  $\bar{\pi}_{r-d}: \mathcal{S}_{\mathbb{Q}}(r-d)_{\overline{\text{Spec } O_K}} \xrightarrow{\sim} \mathcal{N}_{\mathbb{Q}}(r-d)_{\overline{\text{Spec } O_K}}$  in  $\mathbf{D}_K^+(\mathbb{Q}_p)$ , both equivalent to  $\mathbb{Q}_p(r-d)$ ,
- the object  $\mathcal{C}_{\mathbb{Q}, \overline{\text{Spec } O_K}/W^{\text{triv}}}$  in  $\mathbf{D}_K^+(\mathbb{Q}_p)$ , which is equivalent to  $(\mathbb{B}_{\text{st}}^+)^{N=0} = \mathbb{B}_{\text{cris}}^+$ ,
- the map  $\bar{\xi}_{r-d}: \mathcal{S}_{\mathbb{Q}}(r-d)_{\overline{\text{Spec } O_K}} \rightarrow \mathcal{C}_{\mathbb{Q}, \overline{\text{Spec } O_K}/W^{\text{triv}}}$  in  $\mathbf{D}_K^+(\mathbb{Q}_p)$ , which is equivalent to the natural map  $\mathbb{Q}_p(r-d) \hookrightarrow \mathbb{B}_{\text{cris}}^+$ .

For every  $h \geq -1$ , consider the following diagram

$$(B.35) \quad \begin{array}{ccc} \text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}) \otimes_{\mathbb{Q}_p} \mathcal{C}_{\mathbb{Q}, \overline{\text{Spec } O_K}/W^{\text{triv}}} & \longrightarrow & \text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}) \\ \uparrow \bar{\xi}_d \otimes \bar{\xi}_{r-d} & & \uparrow \bar{\xi}_r \\ \text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{S}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}}) \otimes_{\mathbb{Q}_p} \mathcal{S}_{\mathbb{Q}}(r-d)_{\overline{\text{Spec } O_K}} & \longrightarrow & \text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{S}_{\mathbb{Q}}(r)_{\overline{\mathcal{V}^{(h)}}}) \\ \downarrow \bar{\pi}_d \otimes \bar{\pi}_{r-d} & & \downarrow \bar{\pi}_r \\ \text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}}) \otimes_{\mathbb{Q}_p} \mathcal{N}_{\mathbb{Q}}(r-d)_{\overline{\text{Spec } O_K}} & \longrightarrow & \text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(r)_{\overline{\mathcal{V}^{(h)}}}) \end{array}$$

in  $\mathbf{D}_K^+(\mathbb{Q}_p)$ , in which all horizontal maps are induced by cup products. We claim that (B.35) commutes. For  $h \geq 0$ , the upper square commutes by definition, and the lower square commutes by the compatibility of period maps with cup products (see [Tsu99, §3.1] in a more general context). The case for  $h = -1$  follows from the process of taking successive homotopy fiber.

Using the equivalences

$$\text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{S}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}}) \simeq \text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{S}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}}) \otimes_{\mathbb{Q}_p} \mathcal{S}_{\mathbb{Q}}(r-d)_{\overline{\text{Spec } O_K}} \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(d-r),$$

$$\text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}}) \simeq \text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}}) \otimes_{\mathbb{Q}_p} \mathcal{N}_{\mathbb{Q}}(r-d)_{\overline{\text{Spec } O_K}} \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(d-r),$$

and the fact that  $\bar{\pi}_r$  is an equivalence, we obtain the following commutative diagram

$$\begin{array}{ccc} \text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{S}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}}) & & \\ \downarrow & \searrow & \\ \text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{N}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}}) & \xrightarrow{(\bar{\xi}_r \circ \bar{\pi}_r^{-1}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(d-r)} & \text{R}\Gamma(\overline{\mathcal{X}}_{\text{ét}}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(d-r) \end{array}$$

in  $\mathbf{D}_K^+(\mathbb{Q}_p)$  from (B.35).<sup>36</sup> Composing with  $R\Phi_K$  and taking edge maps, we obtain a commutative diagram

$$\begin{array}{ccc} \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(h)}})^{\heartsuit} & & \\ \downarrow & \searrow & \\ \mathbf{H}^1(K, \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{N}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}})) & \xrightarrow{(\bar{\xi}_r \circ \bar{\pi}_r^{-1}) \otimes_{\mathbb{Q}_p}(d-r)} & \mathbf{H}^1(K, \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(d-r)). \end{array}$$

The lemma follows since we have the commutative diagrams

$$\begin{array}{ccc} \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{N}_{\mathbb{Q}}(d)_{\overline{\mathcal{V}^{(h)}}}) & \xrightarrow{(\bar{\xi}_r \circ \bar{\pi}_r^{-1}) \otimes_{\mathbb{Q}_p}(d-r)} & \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(d-r) \\ \downarrow \mathbb{Q}_p(d) \hookrightarrow \mathbb{B}_{\text{cris}} & & \simeq \downarrow \text{Lemma B.22} \\ & & (\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\circ}}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0} \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(d-r) \\ & & \downarrow \mathbb{Q}_p(d-r) \hookrightarrow \mathbb{B}_{\text{st}} \\ \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{N}_{\mathbb{Q}}(0)_{\overline{\mathcal{V}^{(h)}}}) \otimes_{\mathbb{Q}_p} \mathbb{B}_{\text{cris}} & \xrightarrow[\text{Lemma B.23}]{\sim} & (\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\circ}}) \otimes_{K_0} \mathbb{B}_{\text{st}})^{N=0} \end{array}$$

and

$$\begin{array}{ccc} \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}) & \longrightarrow & \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}}/W^{\text{triv}}}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(d-r) \\ \simeq \downarrow \text{Lemma B.22} & & \downarrow \\ (\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\circ}}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0} & \longrightarrow & (\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\circ}}) \otimes_{K_0} \mathbb{B}_{\text{st}})^{N=0} \end{array}$$

in which the upper horizontal arrow is induced by the canonical map  $\mathbb{Q}_p(r-d) \hookrightarrow (\mathbb{B}_{\text{st}}^+)^{N=0} \simeq \mathcal{C}_{\mathbb{Q}, \overline{\text{Spec } \mathcal{O}_K}/W^{\text{triv}}}$  and the cup product, and the right vertical arrow is the composition of two right vertical arrows in the previous diagram.  $\square$

*Proof of Lemma B.18.* By Lemma B.25, we may replace the source of both  $\alpha_q$  and  $\rho_q$ , which is originally  $\mathbf{H}^q(\mathcal{X}_{\acute{e}t}, F_! \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{X}})^{\heartsuit}$ , by  $\mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{S}_{\mathbb{Q}}(r)_{\mathcal{V}^{(h-1)}})^{\heartsuit}$ . Then  $\alpha_q$  (B.15) coincides with the dashed arrow in Lemma B.26 (with  $h = -1$ ). By Lemma B.26, we have

$$\text{Ker} \left( \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(h)}})^{\heartsuit} \rightarrow \mathbf{H}^1 \left( K, (\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\circ}}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0} \right) \right) \subseteq \text{Ker}(\alpha_q).$$

Thus, the lemma will follow if we can show

$$(B.36) \quad \text{Ker}(\rho_q) = \text{Ker} \left( \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(h)}})^{\heartsuit} \rightarrow \mathbf{H}^1 \left( K, (\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\circ}}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+)^{N=0} \right) \right).$$

Lemma B.22 implies that

$$\mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\text{triv}}})^0 = \text{Ker} \left( \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\text{triv}}}) \rightarrow \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[l]^{\circ}}) \right).$$

It induces an isomorphism

$$\frac{\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[l]^{\circ}})}{N\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[l]^{\circ}})} \xrightarrow{\sim} \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\text{triv}}})^0$$

by the distinguished triangle  $\mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[l]^{\circ}}^{\Delta}$  (B.30), under which the Frobenius operator on the right side corresponds to  $p$  times the one on the left side.

<sup>36</sup>The above commutative diagram is parallel to [Sat13, (A.6.4)]. However, somehow unfortunately, the roles of  $d$  and  $r$  here are switched from those there.

Consider the following diagram

$$(B.37) \quad \begin{array}{ccc} \left( \frac{\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[t]^\circ})}{N\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[t]^\circ})} \right)^{\varphi=p^{d-1}} & \hookrightarrow & \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\text{triv}}})^0 \\ \downarrow & & \downarrow \\ \frac{\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ})}{N\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ})} & \xrightarrow{-\delta} & \mathbf{H}^1\left(K, \left(\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+\right)^{N=0}\right) \end{array}$$

in which

- the map  $\delta$  is the edge map induced from the short exact sequence

$$0 \longrightarrow \left(\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+\right)^{N=0} \longrightarrow \mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+ \xrightarrow{N} \mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+ \longrightarrow 0$$

in  $\mathbf{M}_K(\mathbb{Q}_p)$ ,

- the left vertical arrow is the specialization map at  $t = 0$ , and
- the right vertical arrow is the composite map

$$\mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\text{triv}}})^0 \rightarrow \mathbf{H}^1\left(K, \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W^{\text{triv}}}})\right) \xrightarrow{\sim} \mathbf{H}^1\left(K, \left(\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ}) \otimes_{K_0} \mathbb{B}_{\text{st}}^+\right)^{N=0}\right)$$

in which the isomorphism is from Lemma B.22(3).

We show that (B.37) commutes. Applying Lemma A.3 to  $S = \overline{\mathcal{X}}_{\acute{e}t}$ , the distinguished triangle

$$\mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W[t]^\circ}[-1]} \xrightarrow{-N[-1]} \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W[t]^\circ}[-1]} \rightarrow \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W^{\text{triv}}}} \xrightarrow{+1} \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W[t]^\circ}}$$

(which is a shift of the distinguished triangle  $\mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W[t]^\circ}^\Delta$  (B.28)), we know that the image of an element  $\gamma_1 \in \mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[t]^\circ})$  under the composite map

$$\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[t]^\circ}) \rightarrow \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\text{triv}}})^0 \rightarrow \mathbf{H}_K^q(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W^{\text{triv}}}})^0 \rightarrow \mathbf{H}^1\left(K, \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W^{\text{triv}}}})\right)$$

can be represented by the 1-cocycle  $g \mapsto g\tilde{\gamma}_0 - \tilde{\gamma}_0$  for  $g \in G_K$ , where  $\tilde{\gamma}_0$  is an arbitrary element in  $\mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W[t]^\circ})}$  satisfying that  $-N(\tilde{\gamma}_0)$  coincides with

$$\gamma_1 \in \mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[t]^\circ}) \subseteq \mathbf{H}^{q-1}(\overline{\mathcal{X}}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \overline{\mathcal{V}^{(h)}/W[t]^\circ}).$$

Then the commutativity of the right square of (B.37) follows from Lemma B.22 and Lemma B.15 (with  $D = \mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ})$ ).

Now (B.36) follows since the composite map

$$\mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{S}_{\mathbb{Q}}(d)_{\mathcal{V}^{(h)}})^\heartsuit \rightarrow \mathbf{H}^q(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^{\text{triv}}})^0 \xrightarrow{\sim} \frac{\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[t]^\circ})}{N\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W[t]^\circ})} \rightarrow \frac{\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ})}{N\mathbf{H}^{q-1}(\mathcal{X}_{\acute{e}t}, \mathcal{C}_{\mathbb{Q}, \mathcal{V}^{(h)}/W^\circ})}$$

is nothing but  $\rho_q$  (B.22) composed with the isomorphism from Lemma B.21.

Lemma B.18 is proved.  $\square$

## REFERENCES

- [Beĭ87] A. Beĭlinson, *Height pairing between algebraic cycles*, Current trends in arithmetical algebraic geometry (Arcata, Calif., 1985), Contemp. Math., vol. 67, Amer. Math. Soc., Providence, RI, 1987, pp. 1–24, DOI 10.1090/conm/067/902590, (to appear in print). MR902590  $\uparrow$ 59
- [Ber94] V. G. Berkovich, *Vanishing cycles for formal schemes*, Invent. Math. **115** (1994), no. 3, 539–571, DOI 10.1007/BF01231772. MR1262943  $\uparrow$ 63
- [Ber96] ———, *Vanishing cycles for formal schemes. II*, Invent. Math. **125** (1996), no. 2, 367–390, DOI 10.1007/s002220050078. MR1395723  $\uparrow$ 63
- [Ber97] P. Berthelot, *Finitude et pureté cohomologique en cohomologie rigide*, Invent. Math. **128** (1997), no. 2, 329–377, DOI 10.1007/s002220050143 (French). With an appendix in English by Aise Johan de Jong. MR1440308  $\uparrow$ 65
- [BK90] S. Bloch and K. Kato, *L-functions and Tamagawa numbers of motives*, The Grothendieck Festschrift, Vol. I, Progr. Math., vol. 86, Birkhäuser Boston, Boston, MA, 1990, pp. 333–400. MR1086888 (92g:11063)  $\uparrow$ 5

- [Bre97] C. Breuil, *Représentations  $p$ -adiques semi-stables et transversalité de Griffiths*, Math. Ann. **307** (1997), no. 2, 191–224, DOI 10.1007/s002080050031 (French). MR1428871 ↑74, 75
- [Car12] A. Caraiani, *Local-global compatibility and the action of monodromy on nearby cycles*, Duke Math. J. **161** (2012), no. 12, 2311–2413, DOI 10.1215/00127094-1723706. MR2972460 ↑37
- [Car14] ———, *Monodromy and local-global compatibility for  $l = p$* , Algebra Number Theory **8** (2014), no. 7, 1597–1646, DOI 10.2140/ant.2014.8.1597. MR3272276 ↑37
- [CH13] G. Chenevier and M. Harris, *Construction of automorphic Galois representations, II*, Camb. J. Math. **1** (2013), no. 1, 53–73, DOI 10.4310/CJM.2013.v1.n1.a2. MR3272052 ↑36
- [dJ96] A. J. de Jong, *Smoothness, semi-stability and alterations*, Inst. Hautes Études Sci. Publ. Math. **83** (1996), 51–93. MR1423020 ↑75, 76
- [Dis17] D. Disegni, *The  $p$ -adic Gross-Zagier formula on Shimura curves*, Compos. Math. **153** (2017), no. 10, 1987–2074, DOI 10.1112/S0010437X17007308. MR3692745 ↑5
- [Disa] ———, *The  $p$ -adic Gross-Zagier formula on Shimura curves, II: nonsplit primes*, J. Inst. Math. Jussieu. to appear. ↑5
- [Disb] ———, *The universal  $p$ -adic Gross-Zagier formula*. arXiv:2001.00045. ↑5
- [EL] E. Eischen and Z. Liu, *Archimedean zeta integrals for unitary groups*. arXiv:2006.04302. ↑9, 32
- [EHL20] E. Eischen, M. Harris, J. Li, and C. Skinner,  *$p$ -adic  $L$ -functions for unitary groups*, Forum Math. Pi **8** (2020), e9, 160, DOI 10.1017/fmp.2020.4. MR4096618 ↑4, 7
- [GG11] Z. Gong and L. Grenié, *An inequality for local unitary theta correspondence*, Ann. Fac. Sci. Toulouse Math. (6) **20** (2011), no. 1, 167–202 (English, with English and French summaries). MR2830396 ↑34
- [GZ86] B. H. Gross and D. B. Zagier, *Heegner points and derivatives of  $L$ -series*, Invent. Math. **84** (1986), no. 2, 225–320, DOI 10.1007/BF01388809. MR833192 ↑2, 5
- [GK05] E. Grosse-Klönne, *Frobenius and monodromy operators in rigid analysis, and Drinfeld’s symmetric space*, J. Algebraic Geom. **14** (2005), no. 3, 391–437, DOI 10.1090/S1056-3911-05-00402-9. MR2129006 ↑64, 67
- [GD77] A. Grothendieck and P. Deligne, *La classe de cohomologie associée à un cycle*, Cohomologie étale, Lecture Notes in Math., vol. 569, Springer, Berlin, 1977, pp. 129–153 (French). MR3727436 ↑59
- [Hid98] H. Hida, *Automorphic induction and Leopoldt type conjectures for  $GL(n)$* , Asian J. Math. **2** (1998), no. 4, 667–710, DOI 10.4310/AJM.1998.v2.n4.a5. Mikio Sato: a great Japanese mathematician of the twentieth century. MR1734126 ↑3, 24
- [Hyo91] O. Hyodo, *On the de Rham-Witt complex attached to a semi-stable family*, Compositio Math. **78** (1991), no. 3, 241–260. MR1106296 ↑65
- [HK94] O. Hyodo and K. Kato, *Semi-stable reduction and crystalline cohomology with logarithmic poles*, Astérisque **223** (1994), 221–268. Périodes  $p$ -adiques (Bures-sur-Yvette, 1988). MR1293974 ↑65, 81
- [Jac79] H. Jacquet, *Principal  $L$ -functions of the linear group*, Automorphic forms, representations and  $L$ -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 63–86. MR546609 ↑4, 29
- [Jan88] U. Jannsen, *Continuous étale cohomology*, Math. Ann. **280** (1988), no. 2, 207–245, DOI 10.1007/BF01456052. MR929536 ↑56
- [KMSW] T. Kaletha, A. Minguez, S. W. Shin, and P.-J. White, *Endoscopic Classification of Representations: Inner Forms of Unitary Groups*. arXiv:1409.3731. ↑36, 37
- [Kar79] M. L. Karel, *Functional equations of Whittaker functions on  $p$ -adic groups*, Amer. J. Math. **101** (1979), no. 6, 1303–1325, DOI 10.2307/2374142. MR548883 ↑16, 19
- [Kat94] K. Kato, *Semi-stable reduction and  $p$ -adic étale cohomology*, Astérisque **223** (1994), 269–293. Périodes  $p$ -adiques (Bures-sur-Yvette, 1988). MR1293975 ↑69, 74
- [KM74] N. M. Katz and W. Messing, *Some consequences of the Riemann hypothesis for varieties over finite fields*, Invent. Math. **23** (1974), 73–77, DOI 10.1007/BF01405203. MR332791 ↑43
- [KSZ] M. Kisin, S. W. Shin, and Y. Zhu, *The stable trace formula for Shimura varieties of abelian type*. arXiv:2110.05381. ↑5, 36
- [Kob13] S. Kobayashi, *The  $p$ -adic Gross-Zagier formula for elliptic curves at supersingular primes*, Invent. Math. **191** (2013), no. 3, 527–629, DOI 10.1007/s00222-012-0400-9. MR3020170 ↑5
- [Kud02] S. S. Kudla, *Derivatives of Eisenstein series and generating functions for arithmetic cycles*, Astérisque **276** (2002), 341–368. Séminaire Bourbaki, Vol. 1999/2000. MR1886765 ↑2
- [Kud03] ———, *Modular forms and arithmetic geometry*, Current developments in mathematics, 2002, Int. Press, Somerville, MA, 2003, pp. 135–179. MR2062318 ↑2
- [Kud04] ———, *Special cycles and derivatives of Eisenstein series*, Heegner points and Rankin  $L$ -series, Math. Sci. Res. Inst. Publ., vol. 49, Cambridge Univ. Press, Cambridge, 2004, pp. 243–270, DOI 10.1017/CBO9780511756375.009, (to appear in print). MR2083214 ↑2
- [KS97] S. S. Kudla and W. J. Sweet Jr., *Degenerate principal series representations for  $U(n, n)$* , Israel J. Math. **98** (1997), 253–306, DOI 10.1007/BF02937337. MR1459856 ↑19
- [Lan12] K.-W. Lan, *Comparison between analytic and algebraic constructions of toroidal compactifications of PEL-type Shimura varieties*, J. Reine Angew. Math. **664** (2012), 163–228, DOI 10.1515/crelle.2011.099. MR2980135 ↑15, 21
- [Lan99] A. Langer, *Local points of motives in semistable reduction*, Compositio Math. **116** (1999), no. 2, 189–217, DOI 10.1023/A:1000829923416. MR1686773 ↑75
- [LL21] C. Li and Y. Liu, *Chow groups and  $L$ -derivatives of automorphic motives for unitary groups*, Ann. of Math. (2) **194** (2021), no. 3, 817–901, DOI 10.4007/annals.2021.194.3.6. ↑2, 4, 6, 7, 27, 32, 34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 54, 59

- [LL22] ———, *Chow groups and  $L$ -derivatives of automorphic motives for unitary groups, II*, Forum Math. Pi **10** (2022), Paper No. e5, DOI 10.1017/fmp.2022.2. MR4390300 ↑2, 7, 34, 36, 38, 41, 42, 49
- [LZ] C. Li and W. Zhang, *Kudla–Rapoport cycles and derivatives of local densities*, J. Amer. Math. Soc. to appear. ↑19, 41
- [Li92] J.-S. Li, *Nonvanishing theorems for the cohomology of certain arithmetic quotients*, J. Reine Angew. Math. **428** (1992), 177–217, DOI 10.1515/crll.1992.428.177. MR1166512 ↑28
- [Liu11a] Y. Liu, *Arithmetic theta lifting and  $L$ -derivatives for unitary groups, I*, Algebra Number Theory **5** (2011), no. 7, 849–921, DOI 10.2140/ant.2011.5.849. MR2928563 ↑6, 17, 32, 48
- [Liu11b] ———, *Arithmetic theta lifting and  $L$ -derivatives for unitary groups, II*, Algebra Number Theory **5** (2011), no. 7, 923–1000, DOI 10.2140/ant.2011.5.923. MR2928564 ↑21, 50
- [Liu19] ———, *Bounding cubic-triple product Selmer groups of elliptic curves*, J. Eur. Math. Soc. (JEMS) **21** (2019), no. 5, 1411–1508, DOI 10.4171/JEMS/865. MR3941496 ↑55
- [Liu22] ———, *Theta correspondence for almost unramified representations of unitary groups*, J. Number Theory **230** (2022), 196–224, DOI 10.1016/j.jnt.2021.03.027. MR4327954 ↑39, 49
- [LTX<sup>+</sup>22] Y. Liu, Y. Tian, L. Xiao, W. Zhang, and X. Zhu, *On the Beilinson–Bloch–Kato conjecture for Rankin–Selberg motives*, Invent. Math. **228** (2022), no. 1, 107–375, DOI 10.1007/s00222-021-01088-4. MR4392458 ↑5, 10, 54
- [Mok15] C. P. Mok, *Endoscopic classification of representations of quasi-split unitary groups*, Mem. Amer. Math. Soc. **235** (2015), no. 1108, vi+248, DOI 10.1090/memo/1108. MR3338302 ↑5, 23, 36
- [Mok93] A. Mokrane, *La suite spectrale des poids en cohomologie de Hyodo–Kato*, Duke Math. J. **72** (1993), no. 2, 301–337, DOI 10.1215/S0012-7094-93-07211-0 (French). MR1248675 ↑67, 72
- [Nek93] J. Nekovář, *On  $p$ -adic height pairings*, Séminaire de Théorie des Nombres, Paris, 1990–91, Progr. Math., vol. 108, Birkhäuser Boston, Boston, MA, 1993, pp. 127–202, DOI 10.1007/s10107-005-0696-y, (to appear in print). MR1263527 ↑6, 38, 46, 54, 59, 62, 63, 72
- [Nek95] ———, *On the  $p$ -adic height of Heegner cycles*, Math. Ann. **302** (1995), no. 4, 609–686, DOI 10.1007/BF01444511. MR1343644 ↑5, 46, 47, 59
- [Nek00] ———,  *$p$ -adic Abel–Jacobi maps and  $p$ -adic heights*, The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), CRM Proc. Lecture Notes, vol. 24, Amer. Math. Soc., Providence, RI, 2000, pp. 367–379, DOI 10.1090/crmp/024/18, (to appear in print). MR1738867 ↑37, 40, 62
- [NT21a] J. Newton and J. A. Thorne, *Symmetric power functoriality for holomorphic modular forms*, Publ. Math. Inst. Hautes Études Sci. **134** (2021), 1–116, DOI 10.1007/s10240-021-00127-3. MR4349240 ↑5
- [NT21b] ———, *Symmetric power functoriality for holomorphic modular forms, II*, Publ. Math. Inst. Hautes Études Sci. **134** (2021), 117–152, DOI 10.1007/s10240-021-00126-4. MR4349241 ↑5
- [PR87] B. Perrin-Riou, *Points de Heegner et dérivées de fonctions  $L$   $p$ -adiques*, Invent. Math. **89** (1987), no. 3, 455–510, DOI 10.1007/BF01388982 (French). MR903381 ↑2, 5, 51
- [Ral82] S. Rallis, *Langlands’ functoriality and the Weil representation*, Amer. J. Math. **104** (1982), no. 3, 469–515, DOI 10.2307/2374151. MR658543 ↑2, 32
- [Ram] D. Ramakrishnan, *A theorem on  $GL(n)$  a la Tchebotarev*. arXiv:1806.08429. ↑36
- [RSZ20] M. Rapoport, B. Smithling, and W. Zhang, *Arithmetic diagonal cycles on unitary Shimura varieties*, Compos. Math. **156** (2020), no. 9, 1745–1824, DOI 10.1112/s0010437x20007289. MR4167594 ↑37, 44
- [Sat07] K. Sato,  *$p$ -adic étale Tate twists and arithmetic duality*, Ann. Sci. École Norm. Sup. (4) **40** (2007), no. 4, 519–588, DOI 10.1016/j.ansens.2007.04.002 (English, with English and French summaries). With an appendix by Kei Hagihara. MR2191526 ↑71, 72
- [Sat13] ———, *Cycle classes for  $p$ -adic étale Tate twists and the image of  $p$ -adic regulators*, Doc. Math. **18** (2013), 177–247. MR3064983 ↑61, 66, 70, 72, 84
- [Shn16] A. Shnidman,  *$p$ -adic heights of generalized Heegner cycles*, Ann. Inst. Fourier (Grenoble) **66** (2016), no. 3, 1117–1174 (English, with English and French summaries). MR3494168 ↑5, 46, 61
- [Tan99] V. Tan, *Poles of Siegel Eisenstein series on  $U(n, n)$* , Canad. J. Math. **51** (1999), no. 1, 164–175, DOI 10.4153/CJM-1999-010-4. MR1692899 ↑21
- [TY07] R. Taylor and T. Yoshida, *Compatibility of local and global Langlands correspondences*, J. Amer. Math. Soc. **20** (2007), no. 2, 467–493, DOI 10.1090/S0894-0347-06-00542-X. MR2276777 ↑37, 43, 62
- [Tsu99] T. Tsuji,  *$p$ -adic étale cohomology and crystalline cohomology in the semi-stable reduction case*, Invent. Math. **137** (1999), no. 2, 233–411, DOI 10.1007/s002220050330. MR1705837 ↑74, 80, 81, 82, 83
- [Tsu00] ———, *On  $p$ -adic nearby cycles of log smooth families*, Bull. Soc. Math. France **128** (2000), no. 4, 529–575 (English, with English and French summaries). MR1815397 ↑69, 78
- [Wal88] N. R. Wallach, *Lie algebra cohomology and holomorphic continuation of generalized Jacquet integrals*, Representations of Lie groups, Kyoto, Hiroshima, 1986, Adv. Stud. Pure Math., vol. 14, Academic Press, Boston, MA, 1988, pp. 123–151, DOI 10.2969/aspm/01410123, (to appear in print). MR1039836 ↑16
- [Yam14] S. Yamana,  *$L$ -functions and theta correspondence for classical groups*, Invent. Math. **196** (2014), no. 3, 651–732, DOI 10.1007/s00222-013-0476-x. MR3211043 ↑27
- [Yam11] G. Yamashita,  *$p$ -adic Hodge theory for open varieties*, C. R. Math. Acad. Sci. Paris **349** (2011), no. 21–22, 1127–1130, DOI 10.1016/j.crma.2011.10.016 (English, with English and French summaries). MR2855488 ↑81

[YZZ13] X. Yuan, S.-W. Zhang, and W. Zhang, *The Gross-Zagier formula on Shimura curves*, Annals of Mathematics Studies, vol. 184, Princeton University Press, Princeton, NJ, 2013. MR3237437 ↑5

DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BE'ER SHEVA 84105, ISREAL  
*E-mail address*: `disegni@bgu.ac.il`

INSTITUTE FOR ADVANCED STUDY IN MATHEMATICS, ZHEJIANG UNIVERSITY, HANGZHOU 310058, CHINA  
*E-mail address*: `liuyf0719@zju.edu.cn`