
p -ADIC HEIGHTS OF HEEGNER POINTS ON SHIMURA CURVES

by

Daniel Disegni

Abstract. — Let f be a primitive Hilbert modular form of parallel weight 2 and level N for the totally real field F , and let p be a rational prime coprime to $2N$. If f is ordinary at p and E is a CM extension of F of relative discriminant Δ prime to Np , we give an explicit construction of the p -adic Rankin–Selberg L -function $L_{p,\alpha}(f_E, \cdot)$. When the sign of its functional equation is -1 , we show, under the assumption that all primes $\wp|p$ are principal ideals of \mathcal{O}_F which split in \mathcal{O}_E , that its central derivative is given by the p -adic height of a Heegner point on the abelian variety A associated to f .

This p -adic Gross–Zagier formula generalises the result obtained by Perrin-Riou when $F = \mathbf{Q}$ and (N, E) satisfies the so-called Heegner condition. We deduce applications to both the p -adic and the classical Birch and Swinnerton-Dyer conjectures for A .

Contents

Introduction	2
The p -adic Rankin–Selberg L -function	3
Heegner points on Shimura curves and the main theorem	4
Applications to the conjecture of Birch and Swinnerton-Dyer	5
Plan of the proof	9
Perspective	9
Acknowledgments	10
Notation	10
Part I. p-adic L-function and measures	11
1. p -adic modular forms	11
1.1. Hilbert modular forms	11
1.2. Fourier expansion	12
1.3. p -adic modular forms	13
1.4. Operators acting on modular forms	13
1.5. Fourier coefficients of old forms	16
1.6. The functional l_{f_α}	17
2. Theta measure	18
2.1. Weil representation	19
2.2. Theta series	19
2.3. Theta measure	21
2.4. Fourier expansion of the theta measure / I	22

2.5. Fourier expansion of the theta measure / II	25
3. Eisenstein measure	26
3.1. Eisenstein series	26
3.2. Fourier expansion of the Eisenstein measure	26
4. The p -adic L -function	31
4.1. Rankin–Selberg convolution	31
4.2. Convoluted measure and the p -adic L -function in the ordinary case	31
4.3. Interpolation property	33
4.4. Factorisation	35
4.5. Fourier expansion of the analytic kernel	36
Part II. Heights	39
5. p -adic heights and Arakelov theory	39
5.1. Local symbols	39
5.2. The p -adic height pairing	41
5.3. p -adic Arakelov theory – local aspects	42
5.4. p -adic Arakelov theory – global aspects	44
6. Heegner points on Shimura curves	45
6.1. Shimura curves	45
6.2. Heegner points	48
6.3. Hecke action on Heegner points	49
7. Heights of Heegner points	50
7.1. Local heights at places not dividing p	51
7.2. Local heights at p / I	52
7.3. Local heights at p / II	56
Part III. Main theorem and consequences	57
8. Proof of the main theorem	57
8.1. Basic case	57
8.2. Reduction to the basic case	59
9. Periods and the Birch and Swinnerton-Dyer conjecture	60
9.1. Real periods	60
9.2. Quadratic periods	62
References	62

Introduction

In this work we generalise the p -adic analogue of the Gross–Zagier formula of Perrin-Riou [41] to totally real fields, in a generality similar to the work of Zhang [57–59]. We describe here the main result and its applications.

The p -adic Rankin–Selberg L -function. — Let f be a primitive (that is, a normalised new eigenform) Hilbert modular form of parallel weight 2, level N and trivial character for the totally real field F of degree g and discriminant D_F . Let p be a rational prime coprime to $2N$. Fix embeddings ι_∞ and ι_p of the algebraic closure $\overline{\mathbf{Q}}$ of F into \mathbf{C} and $\overline{\mathbf{Q}}_p$ respectively; we let v denote the valuation on $\overline{\mathbf{Q}}_p$, normalised by $v(p) = 1$.

Let $E \subset \overline{\mathbf{Q}}$ be a CM (that is, quadratic and purely imaginary) extension of F of relative discriminant Δ coprime to $D_F N p$, let

$$\varepsilon = \varepsilon_{E/F} : F_{\mathbf{A}}^\times / F^\times \rightarrow \{\pm 1\}$$

be the associated Hecke character and $\mathfrak{N} = N_{E/F}$ be the relative norm. If

$$\mathcal{W} : E_{\mathbf{A}}^\times / E^\times \rightarrow \overline{\mathbf{Q}}^\times$$

is a finite order Hecke character⁽¹⁾ of conductor \mathfrak{f} prime to $N\Delta$, the Rankin–Selberg L -function $L(f_E, \mathcal{W}, s)$ is the entire function defined for $\Re(s) > 3/2$ by

$$L(f_E, \mathcal{W}, s) = L^{N\Delta(\mathcal{W})}(\varepsilon \mathcal{W}|_{F_{\mathbf{A}}^\times}, 2s - 1) \sum_m \frac{a(f, m) r_{\mathcal{W}}(m)}{\mathbf{N}m^s},$$

where $\Delta(\mathcal{W}) = \Delta \mathfrak{N}(\mathfrak{f})$, $r_{\mathcal{W}}(m) = \sum_{\mathfrak{N}(\mathfrak{a})=m} \mathcal{W}(\mathfrak{a})$ (the sum running over all nonzero ideals of \mathcal{O}_E) and

$$L^{N\Delta(\mathcal{W})}(\varepsilon \mathcal{W}|_{\mathcal{O}_F}, s) = \sum_{(m, N\Delta(\mathcal{W}))=1} \varepsilon(m) \mathcal{W}(m) \mathbf{N}m^{-s}.$$

This L -function admits a p -adic analogue (§4). Let E'_∞ be the maximal abelian extension of E unramified outside p , and E_∞ the maximal \mathbf{Z}_p -subextension of E'_∞ . Then $\mathcal{G} = \text{Gal}(E_\infty/E)$ is a direct factor of finite, prime to p index in $\mathcal{G}' = \text{Gal}(E'_\infty/E)$. (It has rank $1 + \delta + g$ over \mathbf{Z}_p , where δ is the Leopoldt defect of F .) For each prime \wp of \mathcal{O}_F dividing p , let

$$P_{\wp, f}(X) = X^2 - a(f, \wp)X + \mathbf{N}\wp$$

be the \wp^{th} Hecke polynomial of f , and assume that $v(\iota_p(a(f, \wp))) = 0$; in this case f is said to be *ordinary*, and there is a unique root α_\wp of $P_{\wp, f}(X)$ which is a p -adic unit. Let $L \subset \overline{\mathbf{Q}}_p$ be the finite extension of \mathbf{Q}_p generated by the Fourier coefficients $a(f, m)$ of f and by the α_\wp for $\wp|p$.

Theorem A. — *There exists a unique element $L_{p, \alpha}(f_E)$ of $\mathcal{O}_L[[\mathcal{G}']] \otimes_{\mathcal{O}_L} L$ satisfying the interpolation property*

$$L_{p, \alpha}(f_E)(\mathcal{W}) = \frac{\tau(\overline{\mathcal{W}}) \mathbf{N}(\Delta(\mathcal{W}))^{1/2} V_p(f, \mathcal{W}) \overline{\mathcal{W}}(\Delta)}{\alpha_{\mathfrak{N}(\mathfrak{f})} \Omega_f} L(f_E, \overline{\mathcal{W}}, 1),$$

for all finite order characters \mathcal{W} of \mathcal{G}' of conductor $\mathfrak{f}(\mathcal{W})$. Here both sides are algebraic numbers,⁽²⁾ $\overline{\mathcal{W}} = \mathcal{W}^{-1}$ and

$$\Omega_f = (8\pi^2)^g \langle f, f \rangle_N$$

⁽¹⁾We will throughout use the same notation for a Hecke character, the associated ideal character, and the associated Galois character.

⁽²⁾By a well-known theorem of Shimura [47]. They are compared via ι_p^{-1} and ι_∞^{-1} .

with $\langle \cdot, \cdot \rangle_N$ the Petersson inner product (1.1.2); $\tau(\overline{\mathcal{W}})$ is a normalized Gauß sum; $V_p(f, \mathcal{W})$ is a product of partial Euler factors at p ; and finally $\alpha_{\mathfrak{N}(f)} = \prod_{\mathfrak{p}|p} \alpha_{\mathfrak{p}}^{v_{\mathfrak{p}}(\mathfrak{N}(f))}$.

This is essentially a special case of results of Panchishkin [38] and Hida [23]; we reprove it entirely here (see §4, especially Theorem 4.3.4) because the precise construction of $L_{p,\alpha}(f_E)$ will be crucial for us. It is obtained, using a technique of Hida and Perrin-Riou, by applying a p -adic analogue of the functional ‘‘Petersson product with f ’’ to a convolution Φ of Eisenstein and theta measures on \mathcal{G}' valued in p -adic modular forms (so that $\Phi = \Phi(\mathcal{W})$ is an analogue of the kernel of the classical Rankin–Selberg convolution). The approach we follow is adelic; one novelty introduced here is that the theta measure is constructed via the Weil representation, which seems very natural and would generalise well to higher rank cases.

On the other hand, Manin [35], Dimitrov [15] and others have constructed a p -adic L -function $L_{p,\alpha}(f, \cdot) \in \mathcal{O}_L[[\mathcal{G}_F]]$ as an analogue of the standard L -function $L(f, s)$, where \mathcal{G}_F is the Galois group of the maximal abelian extension of F unramified outside p ; it is characterised by the interpolation property

$$L_{p,\alpha}(f, \chi) = \frac{\tau(\overline{\chi})\mathbf{N}(\mathfrak{f}(\chi))^{1/2}}{\alpha_{\mathfrak{f}(\chi)}} \frac{L(f, \overline{\chi}, 1)}{\Omega_f^+}$$

for all finite order characters χ of conductor $\mathfrak{f}(\chi)$ which are trivial at infinity and ramified at all primes $\mathfrak{v}|p$. (Here Ω_f^+ is a suitable period, cf. §9.1, and $\tau(\chi)$ is again a normalised Gauß sum.)

The corresponding formula for complex L -functions implies a factorisation (Proposition ??)

$$(0.0.1) \quad L_{p,\alpha}(f_E, \chi \circ \mathfrak{N}) = \frac{\Omega_f^+ \Omega_{f_\varepsilon}^+}{D_E^{-1/2} \Omega_f} L_{p,\alpha}(f, \chi) L_{p,\alpha}(f_\varepsilon, \chi),$$

where f_ε is the form with coefficients $a(f_\varepsilon, m) = \varepsilon(m)a(f, m)$ and $D_E = \mathbf{N}(\Delta)$.

Heegner points on Shimura curves and the main theorem. — Suppose that $\varepsilon(N) = (-1)^{g-1}$, where $g = [F : \mathbf{Q}]$. Then for each embedding $\tau : F \rightarrow \mathbf{C}$, there is a quaternion algebra $B(\tau)$ over F ramified exactly at the finite places $\mathfrak{v}|N$ for which $\varepsilon(\mathfrak{v}) = -1$ and the infinite places different from τ ; it admits an embedding $\rho : E \hookrightarrow B(\tau)$, and we can consider an order R of $B(\tau)$ of discriminant N and containing $\rho(\mathcal{O}_E)$. These data define a *Shimura curve* X . It is an algebraic curve over F , whose complex points for any embedding $\tau : F \rightarrow \mathbf{C}$ are described by

$$X(\mathbf{C}_\tau) = B(\tau)^\times \backslash \mathfrak{H}^\pm \times \widehat{B}(\tau)^\times / \widehat{F}^\times \widehat{R}^\times \cup \{\text{cusps}\}.$$

It plays the role of the modular curve $X_0(N)$ in the works of Gross–Zagier [20] and Perrin-Riou [41] who consider the case $F = \mathbf{Q}$ and $\varepsilon(\mathfrak{v}) = 1$ for all $\mathfrak{v}|N$ (it is only in this case that the set of cusps is not empty).

The curve X is connected but not geometrically connected. Let $J(X)$ be its Albanese (\cong Jacobian) variety; it is an abelian variety defined over F , geometrically isomorphic to the product of the Albanese varieties of the geometrically connected components of X . There is a natural map $\iota : X \rightarrow J(X) \otimes \mathbf{Q}$ given by $\iota(x) = [x] - [\xi]$, where $[\xi] \in \text{Cl}(X) \otimes \mathbf{Q}$ is a canonical divisor class constructed in [57] having degree 1 in every geometrically connected component of X ; an integer multiple of ι gives a morphism $X \rightarrow J(X)$ defined over F .

As in the modular curve case, the curve X admits a finite collection of *Heegner points* defined over the Hilbert class field H of E and permuted simply transitively by $\text{Gal}(H/E)$. They are the points represented by (x_0, t) for $t \in \widehat{E}^\times / E^\times \widehat{F}^\times \widehat{\mathcal{O}}_E^\times$ when we use the complex description above

and view $E \subset B$ via ρ . We let γ be any such Heegner point, and let $[z]$ denote the class

$$[z] = u^{-1} \iota \left(\text{Tr}_{H/E} \gamma \right) \in J(X)(E) \otimes \mathbf{Q},$$

where $u = [\mathcal{O}_E^\times : \mathcal{O}_F^\times]$.

As a consequence of Jacquet–Langlands theory, the Hecke algebra on Hilbert modular forms of level N acts through its quaternionic quotient on $J(X)$. Let $z_f \in J(X)(E) \otimes \overline{\mathbf{Q}}$ be the f -component of $[z]$.

Heights and the formula. — On any curve X over a number field E , there is a notion (§5.2) of p -adic height $\langle \cdot, \cdot \rangle_\ell$ attached to the auxiliary choices of splittings of the Hodge filtrations on $H_{\text{dR}}^1(X/E_w)$ for $w|p$ and of a p -adic logarithm $\ell: E_A^\times/E^\times \rightarrow \mathbf{Q}_p$. It is a symmetric bilinear pairing on the group of degree zero divisors on X modulo rational equivalence, which we can view as a pairing on $J(X)(E)$. More generally, for any abelian variety A/E there is defined a p -adic height pairing on $A(E) \times A^\vee(E)$. In our case, there is a canonical choice for the Hodge splittings on the f -components of the Albanese variety $J(X)$, given by the unit root subspaces, and we choose our height pairing on $J(X)$ to be compatible with this choice.

Let \mathcal{W} be a Hecke character of E taking values in $1 + p\mathbf{Z}_p \subset \mathbf{Z}_p^\times$, ramified at all places dividing p . Under the assumption $\varepsilon(N) = (-1)^{g-1}$, the value $L_{p,\alpha}(f_E, \mathbf{1})$ is zero by the complex functional equation and the interpolation property. (In fact, the restriction of $L_{p,\alpha}(f_E)$ to \mathcal{G} also satisfies a functional equation with sign $(-1)^g \varepsilon(N)$ relating its values at \mathcal{W} to its values at \mathcal{W}^{-c} , where c is the nontrivial automorphism of E/F and $\mathcal{W}^c(\sigma) = \mathcal{W}(c\sigma c)$; see Proposition 4.2.2 for a proof in a special case.) Then we can consider the derivative of $L_{p,\alpha}(f_E)$ in the \mathcal{W} -direction

$$L'_{p,\alpha,\mathcal{W}}(f_E, \mathbf{1}) = \left. \frac{d}{ds} \right|_{s=0} L_{p,\alpha}(f_E)(\mathcal{W}^s).$$

Theorem B. — Assume that $\Delta_{E/F}$ is totally odd and that every prime $\wp|p$ is a principal ideal in \mathcal{O}_F and splits in \mathcal{O}_E . Suppose that $\varepsilon_{E/F}(N) = (-1)^{g-1}$. Then $L_{p,\alpha}(f_E, \mathbf{1}) = 0$ and

$$L'_{p,\mathcal{W}}(f_E, \mathbf{1}) = D_F^{-2} \prod_{\wp|p} \left(1 - \frac{1}{\alpha_\wp} \right)^2 \left(1 - \frac{1}{\varepsilon(\wp)\alpha_\wp} \right)^2 \langle z_f, z_f \rangle_{\mathcal{W},\alpha}$$

where $\langle \cdot, \cdot \rangle_{\mathcal{W},\alpha}$ is the height pairing on $J(X)(E)$ associated to the logarithm $\ell = \left. \frac{d}{ds} \right|_{s=0} \mathcal{W}^s$.

The hypothesis that the primes $\wp|p$ are principal is a technical assumption which intervenes only in Proposition 8.1.1. The assumption that they split in E is essential to the argument but, like the assumption on $\Delta_{E/F}$, it can be removed *a posteriori* if the left-hand side of the formula below is nonzero – see §8.2.

Applications to the conjecture of Birch and Swinnerton-Dyer. — It is conjectured that to any Hilbert modular newform f one can attach a simple abelian variety $A = A_f$ over F , characterised uniquely up to isogeny⁽³⁾ by the equality of L -functions

$$L(A, s) = \prod_{\sigma: M_f \rightarrow \mathbf{C}} L(f^\sigma, s).$$

⁽³⁾Thanks to Faltings’s isogeny theorem [17].

Here $M = M_f$ is the field generated by the Fourier coefficients of f ; A has dimension $[M : \mathbf{Q}]$ and its endomorphism algebra contains M (we say that A is of $GL_2(M)$ -type; in fact since F is totally real, A is of *strict* GL_2 -type, that is, its endomorphism algebra equals M – see e.g. [55, Lemma 3.3]). The conjecture is known to be true [57, Theorem B] when either $[F : \mathbf{Q}]$ is odd or $v(N)$ is odd for some finite place v (the assumptions of Theorem B above imply that one of these conditions holds); in this case A is a quotient ϕ of $J(X)$ for a suitable Shimura curve X of the type described above. Viceversa any abelian variety of GL_2 -type (for some field M) over a totally real field F is conjectured to be associated to a Hilbert modular form f as above.

In view of known $\text{Aut}(\mathbf{C}/\mathbf{Q})$ -equivariance properties of automorphic L -functions and the above equality, the order of vanishing of $L(A, s)$ at $s = 1$ will be an integer multiple $r[M : \mathbf{Q}]$ of the dimension of A . We call r the M -order of vanishing of $L(A, s)$ or the *analytic M -rank* of A .

Conjecture (Birch and Swinnerton-Dyer). — *Let A be an abelian variety of $GL_2(M)$ -type over a totally real field F of degree g .*

1. *The M -order of vanishing of $L(A, s)$ at $s = 1$ is equal to the dimension of $A(F)_{\mathbf{Q}}$ as M -vector space.*
2. *The Tate-Shafarevich group $\text{III}(A/F)$ is finite, and the leading term of $L(A, s)$ at $s = 1$ is given by*

$$\frac{L^*(A, 1)}{\Omega_A} = D_F^{-d/2} |\text{III}(A/F)|_{R_A} \prod_{v \neq \infty} c_v = \text{BSD}(A),$$

where $d = \dim A = [M : \mathbf{Q}]$, the c_v are the Tamagawa numbers of A at finite places (almost all equal to 1),

$$\Omega_A = \prod_{\tau: F \rightarrow \mathbf{R}} \int_{A(\mathbf{R}_{\tau})} |\omega_A|_{\tau}$$

for a Néron differential⁽⁴⁾ ω_A , and

$$R_A = \frac{\det(\langle x_i, y_j \rangle)}{[A(F) : \sum \mathbf{Z}x_i][A^{\vee}(F) : \sum \mathbf{Z}y_j]}$$

is the regulator of the Néron-Tate height pairing on $A(F) \times A^{\vee}(F)$, defined using any bases $\{x_i\}$, $\{y_j\}$ of $A(F)_{\mathbf{Q}}$ and $A^{\vee}(F)_{\mathbf{Q}}$.

By the automorphic description of $L(A, s)$ and results of Shimura [47], we know that $L(A, s) / \prod_{\sigma: M_f \rightarrow \mathbf{C}} \Omega_{f\sigma}^+$ is an algebraic number. Comparison with the Birch and Swinnerton-Dyer conjecture suggests the following conjecture.

Conjecture. — *We have*

$$\Omega_A \sim \prod_{\sigma: M_f \rightarrow \mathbf{C}} \Omega_{f\sigma}^+ \quad \text{in } \mathbf{C}^{\times} / \overline{\mathbf{Q}}^{\times}.$$

The conjecture is known for $F = \mathbf{Q}$ [48] or when A has complex multiplication (over $\overline{\mathbf{Q}}$) [6]; see §9 below for a more precise conjecture and some further evidence and motivation.

⁽⁴⁾When it exists, which is only guaranteed if $F = \mathbf{Q}$. Otherwise, we take for ω_A any generator of $H^0(A, \Omega_{A/F}^d)$ and to define Ω_A we divide by the product of the indices $[H^0(\mathcal{A}_v, \Omega_{\mathcal{A}_v}^d) : \mathcal{O}_{F,v} \widetilde{\omega}_A] : \mathcal{O}_{F,v} \widetilde{\omega}_A$ of (the extension of) ω_A in the space of top differentials on the local Néron models $\mathcal{A}_v / \mathcal{O}_{F,v}$ of A .

Assuming the conjecture, given a choice of roots α which we omit from the notation, and automorphic p -adic L -functions $L_p(f^\sigma)$, we can define a p -adic L -function $L_p(A)$ for A by

$$L_p(A) = \frac{\prod_{\sigma} \Omega_{f^\sigma}^+}{\Omega_A} \prod_{\sigma: M_f \rightarrow \mathbb{C}} L_p(f^\sigma)$$

for any prime p of good reduction.

Then, fixing a ramified Hecke character $\nu: \mathcal{G}'_F \rightarrow 1 + p\mathbb{Z}_p \subset \mathbb{Z}_p^\times$ which we omit from the notation, one can formulate a p -adic version of the Birch and Swinnerton-Dyer conjecture similarly as above for $L_p(A, \nu^s)$:⁽⁵⁾ the formula reads

$$\prod_{\wp|p} (1 - \alpha_{\wp}^{-1})^{-2} L_p^*(A, 1) = \text{BSD}_p(A)$$

where $\text{BSD}_p(A)$ differs from $\text{BSD}(A)$ only in the regulator term, which is now the regulator of the p -adic height pairing on $A(F) \times A^\vee(F)$ associated to the p -adic logarithm ℓ deduced from ν as in Theorem B.

Similarly, one can formulate a main conjecture of Iwasawa theory for $L_p(A)$.

Then, just as in [41], we can deduce the following arithmetic application of Theorem B.

Theorem C. — *For the abelian variety $A = A_f$ and an ordinary prime $p > 2$ decomposing into principal prime ideals in \mathcal{O}_F , we have:*

1. *The following are equivalent:*
 - (a) *The p -adic L -function $L_{p,\alpha}(A, \nu^s)$ has M_f -order of vanishing $r \leq 1$ at the central point.*
 - (b) *The complex L -function $L(A, s)$ has M_f -order of vanishing $r \leq 1$ at the central point and the p -adic height pairing associated to ν is non-vanishing on $A(F)$.*
2. *If either of the above assumptions holds, the rank parts of the classical and the p -adic Birch and Swinnerton-Dyer conjecture are true for A and the Tate-Shafarevich group of A is finite.*
3. *If moreover the cyclotomic Iwasawa main conjecture is true for A , then the classical and the p -adic Birch and Swinnerton-Dyer formulas for A are true up to a p -adic unit.*

Proof. — In part 1, the statement follows trivially from the construction of $L_p(A)$ if $r = 0$; if $r = 1$, both conditions are equivalent to the assertion that for a suitable CM extension E , the Heegner point $z_f = z_{f,E}$ is nontorsion: this is obvious from our main theorem in case 1a; in case 1b, by the work of Zhang [57, 58] (generalising Gross–Zagier [20] and Kolyvagin [32, 34]), the Heegner point

$$P = \sum_{\sigma} \text{Tr}_{E/F} \phi(z_{f^\sigma, E}) \in A(F) \otimes \mathbb{Q}$$

(with $\phi: J(X) \rightarrow A$) generates $A(F) \otimes \mathbb{Q}$ as M_f -vector space, so that the p -adic height pairing on $A(F)$ is non-vanishing if and only if it is nonzero at z_f . Part 2 then follows from 1 and the results of Zhang [57, 58].

Schneider [45] proves an “arithmetic” version of the p -adic Birch and Swinnerton-Dyer formula for (the Iwasawa L -function associated to) A ,⁽⁶⁾ which under the assumption of 3 can be compared to the analytic p -adic formula as explained in [41] to deduce the p -adic Birch and Swinnerton-Dyer formula up to a p -adic unit. In the analytic rank 0 case the classical Birch and

⁽⁵⁾Here $s \in \mathbb{Z}_p$ and the central point is $s = 0$, corresponding to $\nu^0 = 1$.

⁽⁶⁾In the general non-ordinary case, the analogue of Schneider’s result is proved by Perrin-Riou [PRbook].

Swinnerton-Dyer formula follows immediately. In the case of analytic rank 1, recall that the main result of [57, 59] is, in our normalisation, the formula

$$\frac{L'(f_E, 1)}{\Omega_f} = \frac{1}{D_F^2 D_E^{1/2}} \langle z_f, z_f \rangle =: D_E^{-1/2} \text{GZ}(f_E)$$

(where $\langle \cdot, \cdot \rangle$ denotes the Néron–Tate height); whereas we introduce the notation $\text{GZ}_p(f_E)$ to write our formula (for any fixed ramified cyclotomic character $\mathscr{W} = \nu \circ \mathfrak{N}$) as

$$L'_p(f_E, 1) = \prod_{\wp | p} \left(1 - \frac{1}{\alpha_\wp} \right)^2 \left(1 - \frac{1}{\varepsilon(\wp) \alpha_\wp} \right)^2 \text{GZ}_p(f_E).$$

Then, after choosing E suitably so that $L(f_\varepsilon, 1) \neq 0$ (which can be done by [5], [52]), we can argue as in [41] to compare the p -adic and the complex Birch and Swinnerton-Dyer formulas via the corresponding Gross–Zagier formulas to get the result. Namely, we have

$$\begin{aligned} \frac{L^*(A, 1)}{\Omega_A \text{BSD}(A)} &= \frac{\prod_\sigma \Omega_{f^\sigma}^+}{\Omega_A} \frac{1}{\text{BSD}(A)} \prod_\sigma \frac{L'(f_E^\sigma, 1)}{\Omega_{f^\sigma}} \frac{\Omega_{f^\sigma}}{\Omega_{f^\sigma}^+ \Omega_{f_\varepsilon}^+} \frac{\Omega_{f_\varepsilon}^+}{L(f_\varepsilon^\sigma, 1)} \\ &= \frac{\prod_\sigma \Omega_{f^\sigma}^+}{\Omega_A} \frac{\prod_\sigma \text{GZ}(f_E^\sigma)}{\text{BSD}(A)} \prod_\sigma \frac{D_E^{-1/2} \Omega_{f^\sigma}}{\Omega_{f^\sigma}^+ \Omega_{f_\varepsilon}^+} \frac{\Omega_{f_\varepsilon}^+}{L(f_\varepsilon^\sigma, 1)} \end{aligned}$$

by the complex Gross–Zagier formula and the factorisation of $L(f_E, s)$. Similarly,

$$\prod_{\wp | p} (1 - \alpha_\wp^{-1})^{-2} \frac{L_p^*(A, 1)}{\text{BSD}_p(A)} = \frac{\prod_\sigma \Omega_{f^\sigma}^+}{\Omega_A} \frac{\prod_\sigma \text{GZ}_p(f_E^\sigma)}{\text{BSD}_p(A)} \prod_\sigma \frac{D_E^{-1/2} \Omega_{f^\sigma}}{\Omega_{f^\sigma}^+ \Omega_{f_\varepsilon}^+} \frac{\Omega_{f_\varepsilon}^+}{L(f_\varepsilon^\sigma, 1)}$$

by the p -adic Gross–Zagier formula, the factorisation of $L_p(f_E)$ and the interpolation property of $L_p(f_\varepsilon)$. Since we are assuming to know that the left-hand side of the last formula is a p -adic unit, the result follows from observing the equality

$$\frac{\prod_\sigma \text{GZ}(f_E^\sigma)}{\text{BSD}(A)} = \frac{\prod_\sigma \text{GZ}_p(f_E^\sigma)}{\text{BSD}_p(A)}$$

of rational numbers.⁽⁷⁾ □

Alternative approaches to the Birch and Swinnerton-Dyer formula in rank one have recently been proposed by Wei Zhang [60] and Xin Wan.

Discussion of the assumptions. — The conjecture on periods could be dispensed of if one were willing to work with a “wrong” p -adic L -function for A (namely, one without the period ratio appearing in the definition above). Then at least the rank part of the p -adic Birch and Swinnerton-Dyer conjecture makes sense and parts 1 and 2 of the Theorem hold. The nonvanishing of the p -adic height pairing is only known for CM elliptic curves [2]. The Iwasawa main conjecture is known in most cases for ordinary elliptic curves over \mathbf{Q} thanks to the work of Rubin, Kato and Skinner–Urban (see [50]). For Hilbert modular forms, in the ordinary case one divisibility in

⁽⁷⁾The rationality of the ratios follows from the fact that the z_{f^σ} essentially belong to $J(X)(F)$ – that is, they belong to the $+1$ -eigenspace for the action of $\text{Gal}(E/F)$ on $J(X)(E) \otimes \overline{\mathbf{Q}}$ – and that in this sense, their images $\phi(z_{f^\sigma})$ form a $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ -invariant basis of $A(F) \otimes \overline{\mathbf{Q}}$, orthogonal for the height pairing.

the CM case is proved by Hsieh [26] and in the general case there is work in progress towards one divisibility by Xin Wan. (The other divisibility is not known but could be within reach with current methods, cf. [51, remarks on top of p.6].) We can then record the following unconditional result, whose assumptions are inherited from [26].

Theorem D. — *Let A/F be an elliptic curve with complex multiplication by the ring of integers \mathcal{O}_K of an imaginary quadratic field K . Let $K' = FK$ and let $h_{K'}^- = h_{K'}/h_K$ be the relative class number of K'/F . Let $p \nmid 6h_{K'}^- D_F$ be a prime such that for all primes $\wp | p$, \wp is principal and A has good ordinary reduction at \wp . Suppose that $\text{ord}_{s=1} L(A, s) \leq 1$. Then*

$$v_p \left(\frac{L^*(A, 1)}{R_A \Omega_A} \right) \leq v_p \left(|\text{III}(A/F)| \prod_{v \nmid \infty} c_v \right).$$

Results toward the divisibility in the opposite direction can be obtained from the method of Kolyvagin, cf. [33] (for $F = \mathbf{Q}$) and [25] (for general F but excluding the CM case).

Plan of the proof. — The proof of the main formula follows the strategy of Perrin-Riou [41]. It is enough (see §8) to study the case where \mathcal{W} is cyclotomic ($\mathcal{W} = \mathcal{W}^c$), since both sides of the formula are zero when \mathcal{W} is anticyclotomic ($\mathcal{W} \mathcal{W}^c = 1$).

In the first part of this paper, we construct the measure Φ on \mathcal{G} valued in p -adic (in fact, overconvergent) modular forms such that $L_{p,\alpha}(f_E)(\mathcal{W})$ essentially equals $l_{f_\alpha}(\Phi(\mathcal{W}))$, where l_{f_α} is a p -adic analogue of the functional “Petersson product with f ” on p -adic modular forms. This allows us to write

$$L'_{p,\alpha,\mathcal{W}}(f_E, 1) \doteq l_{f_\alpha}(\Phi'_{\mathcal{W}}),$$

where \doteq denotes equality up to suitable nonzero factors, and $\Phi'_{\mathcal{W}} = \frac{d}{ds} \Phi(\mathcal{W}^s)|_{s=0}$ is an overconvergent p -adic Hilbert modular form.

On the other hand, there is a modular form $\Psi = \Psi_\alpha$ with Fourier coefficients given by $\langle z, T(m)z \rangle_{\mathcal{W},\alpha}$, so that $l_{f_\alpha}(\Psi) \doteq \langle z_f, z_f \rangle_{\mathcal{W},\alpha}$. It can be essentially written as a sum of modular forms $\Psi_{\text{fin}} + \Psi_{\alpha,p}$, where Ψ_{fin} encodes the local contributions to the height from places not dividing p and $\Psi_{\alpha,p} = \sum \Psi_{\alpha,\wp}$ the contribution from the places \wp above p . Then we can show by explicit computation that the Fourier coefficients of Φ' are equal to the Fourier coefficients of Ψ_{fin} up to the action of suitable Hecke operators at p . The desired formula then follows once we show that $l_{f_\alpha}(\Psi_{\alpha,p})$ vanishes. To prove this we examine the effect of the operator U_\wp on $\Psi_{\alpha,\wp}$, and find that, in a suitable quotient space, the ordinary projection of $\Psi_{\alpha,\wp}$ is zero. The study of $\Psi_{\alpha,\wp}$ follows the methods of Perrin-Riou.

One difficulty in the approach just outlined is that compared to the case of modular curves there are no cusps available, so that in this case the divisors z and $T(m)z$ have intersecting supports and the decomposition of the height pairing into a sum of local pairings is not available. Our solution to this problem, which is inspired from the work of Zhang [57], is to make use of p -adic Arakelov theory as developed by Besser [4] (see §5.3) and work consistently in a suitable quotiented space of Fourier coefficients.

Perspective. — The original Gross–Zagier formula has undergone an impressive transformation since its first appearance in 1986, culminating in the recent book of Yuan–Zhang–Zhang [55]. Obviously, this work is only a first attempt at catching up on the p -adic side. An extension of

the present results to higher weights, as achieved by Nekovář [37] for the case of modular curves, would also be of interest.

Acknowledgments. — The present paper grew out of my Columbia thesis. I am grateful to my advisor Professor Shou-Wu Zhang for suggesting this area of research and for his support and encouragement; and to Amnon Besser, Shinichi Kobayashi, Luis Garcia Martinez, David Loeffler, Yifeng Liu, Giovanni Rosso, Eric Urban, Jeanine Van Order and Shou-Wu Zhang for useful conversations or correspondence.

This work builds upon the works of Perrin-Riou [41] and Zhang [57–59] – and, of course, Gross–Zagier [20]. My debt to their ideas cannot be underestimated and will be obvious to the reader.

Notation. — Throughout this text we use the following notation and assumptions, unless otherwise noted:

- F is a totally real field of degree g ;
- \mathbf{N}_F is the monoid of nonzero ideals of \mathcal{O}_F ;
- $|\cdot|_v$ is the standard absolute value on F_v ;
- $\mathbf{A} = \mathbf{A}_F$ is the adèle ring of F ; if $*$ is a place or a set of places or an ideal of F , the component at $*$ (respectively away from $*$) of an adelic object x is denoted x_* (respectively x^*). For example if $\phi = \prod_v \phi_v$ is a Hecke character and δ is an ideal of \mathcal{O}_F we write $\phi_\delta(y) = \prod_{v|\delta} \phi_v(y_v)$, and $|y|_\delta = \prod_{v|\delta} |y|_v$. We also use the notation

$$|m|_v = |\pi_m|_v, \quad |m|_\delta = |\pi_m|_\delta, \quad \phi_v(m) = \phi_v(\pi_m), \quad \phi_\delta(m) = \phi_\delta(\pi_m)$$

if m is an ideal of \mathcal{O}_F and ϕ is unramified at δ (here π_m satisfies $\pi_m \mathcal{O}_F = m$).

- “ $>$ ” denotes the partial order on \mathbf{A}_F given by $x > 0$ if and only if x_∞ is totally positive;
- $R_{\mathbf{A}} = R \otimes_F \mathbf{A}$ if R is an F -algebra;
- $\mathbf{N}m$ is the absolute norm of an ideal m in a number field (the index of m in the ring of integers: it is a positive natural number);
- d_F is the different of F ;
- π_N , for N an ideal of \mathcal{O}_F , is the idèle with components $\pi_v^{v(N)}$ for $v \nmid \infty$ and 1 for $v|\infty$.
- $D_F = \mathbf{N}d_F$ is the discriminant of F .
- $m^\times = \{a \in F_{\mathbf{A}}^\times \mid a \mathcal{O}_F = m\}$ if m is any nonzero fractional ideal of F (this notation will be used with $m = d_F^{-1}$).
- E is a quadratic CM (that is, totally imaginary) extension of F ;
- $\mathfrak{D} = \mathfrak{D}_{E/F}$ is the different of E/F .
- $\mathfrak{N} = N_{E/F}$ is the relative norm on E or any E -algebra;
- $\Delta = \Delta_{E/F} = \mathfrak{N}(\mathfrak{D})$ is the relative discriminant of E/F and we assume

$$(\Delta_{E/F}, D_F N p) = 1;$$

in §§2.5, 4.5 and part of §3.2 we further assume that

$$(\Delta, 2) = 1$$

and in §§ 7.2, 7.3, 8.1, that

$$(\Delta, 2) = 1 \text{ and all primes } \wp \text{ dividing } p \text{ are split in } E.$$

- $D_E = \mathbf{N}(\Delta)$ is the absolute discriminant of E .

- $U_F(N)$ is the subgroup of $\widehat{\mathcal{O}}_F^\times = \prod_v \mathcal{O}_{F,v}^\times \subset F_{\mathbf{A}^\infty}^\times$ consisting of elements $x \equiv 1 \pmod{N\widehat{\mathcal{O}}_F}$, if N is any ideal of \mathcal{O}_F ;
- $\mathbf{e}_v(x) = \exp(-2\pi i \{\mathrm{Tr}_{F_v/\mathbf{Q}_p}(x)\}_p)$ for $v|p < \infty$ and $\{y\}_p$ the p -fractional part of $y \in \mathbf{Q}_p$ is the standard additive character of F_v , with conductor $d_{F,v}^{-1}$; for $v|\infty$, $\mathbf{e}_v(x) = \exp(2\pi i \mathrm{Tr}_{F_v/\mathbf{R}}(x))$;
- $\mathbf{e}(x) = \prod_v \mathbf{e}_v(x_v)$ is the standard additive character of \mathbf{A}_F .
- $\mathbf{1}_Y$ is the characteristic function of the set Y ;
- if φ is any logical proposition, we define $\mathbf{1}[\varphi]$ to be 1 when φ is true and 0 when φ is false - e.g. $\mathbf{1}[x \in Y] = \mathbf{1}_Y(x)$.

PART I

p -ADIC L -FUNCTION AND MEASURES

This part is dedicated to the construction of the measure giving the p -adic Rankin–Selberg L -function $L_{p,\alpha}(f_E)$ and to the computation of its Fourier coefficients.

1. p -adic modular forms

1.1. Hilbert modular forms. — Let us define compact subgroups of $\mathrm{GL}_2(\mathbf{A}^\infty)$ as follows:

- $K_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\widehat{\mathcal{O}}_F) \mid c \equiv 0 \pmod{N\widehat{\mathcal{O}}_F} \right\}$ if N is an ideal of \mathcal{O}_F ;
- $K_1(N, n) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K_0(N) \mid a \equiv 1 \pmod{N/n\widehat{\mathcal{O}}_F}, d \equiv 1 \pmod{n\widehat{\mathcal{O}}_F} \right\}$ if $n|N$ are ideals of \mathcal{O}_F .

Let k be an element of $\mathbf{Z}_{\geq 0}^{\mathrm{Hom}(F, \overline{\mathbf{Q}})}$ and ψ be a character of $F_{\mathbf{A}^\infty}^\times / F^\times$ of conductor dividing N satisfying $\psi_v(-1) = (-1)^{k_v}$ for $v|\infty$. A **Hilbert modular form** of weight k , level $K_1(N)$ and character ψ is a smooth function

$$f : \mathrm{GL}_2(F) \backslash \mathrm{GL}_2(\mathbf{A}_F) \rightarrow \mathbf{C}$$

of moderate growth⁽⁸⁾ satisfying⁽⁹⁾

$$f \left(\begin{pmatrix} z & \\ & z \end{pmatrix} g \begin{pmatrix} a & b \\ c & d \end{pmatrix} r(\theta) \right) = \psi(z) \psi_N(a) \mathbf{e}_\infty(k \cdot \theta) f(g)$$

for each $z \in F_{\mathbf{A}^\infty}^\times$, $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K_0(N)$ and $\theta = (\theta_v)_{v|\infty} \in F_\infty$, with $r(\theta) = \prod_{v|\infty} r(\theta_v)$ and $r(\theta_v) = \begin{pmatrix} \cos \theta_v & \sin \theta_v \\ -\sin \theta_v & \cos \theta_v \end{pmatrix} \in \mathrm{SO}_2(F_v)$. If k is constant we say that f has parallel weight; in this work we will be almost exclusively concerned with forms of parallel weight, and we will assume that we are in this situation for the rest of this section.

⁽⁸⁾That is, for every g the function $\mathbf{A}^\times \ni y \mapsto f \left(\begin{pmatrix} y & \\ & 1 \end{pmatrix} g \right)$ grows at most polynomially in $|y|$ as $|y| \rightarrow \infty$.

⁽⁹⁾Recall the notation $\psi_N = \prod_{v|N} \psi_v$.

We call f holomorphic if the function on $\mathfrak{H}^{\text{Hom}(F, \overline{\mathbf{Q}})} = \{x_\infty + iy_\infty \in F \otimes \mathbf{C} \mid y_\infty > 0\}$

$$x_\infty + iy_\infty \mapsto (\psi^n)^{-1}(y)|y|^{-k/2} f \left(\begin{pmatrix} y & x \\ & 1 \end{pmatrix} \right)$$

is holomorphic; in this case such function determines f .

Petersson inner product. — We define a Haar measure dg on $Z(\mathbf{A}_F) \backslash \mathbf{GL}_2(\mathbf{A}_F)$ (where $Z \cong \mathbf{G}_m$ denotes the center of \mathbf{GL}_2) as follows. Recall the Iwasawa decomposition

$$(1.1.1) \quad \mathbf{GL}_2(\mathbf{A}_F) = B(\mathbf{A}_F) K_0(1) K_\infty$$

where $K_\infty = \prod_{v|\infty} \mathbf{SO}_2(F_v)$. Let $dk = \otimes_v dk_v$ be the Haar measure on $K = K_0(1) K_\infty$ with volume 1 on each component. Let $dx = \otimes_v dx_v$ be the Haar measure such that dx_v is the usual Lebesgue measure on \mathbf{R} if $v|\infty$, and $\mathcal{O}_{F,v}$ has volume 1 if $v \nmid \infty$. Finally let $d^\times x = \otimes_v d^\times x_v$ on $F_\mathbf{A}^\times$ be the product of the measures given by $d^\times x_v = |dx_v/x_v|$ if $v|\infty$ and by the condition that $\mathcal{O}_{F,v}^\times$ has

volume 1 if $v|\infty$. Then we can use the Iwasawa decomposition $g = \begin{pmatrix} z & \\ & z \end{pmatrix} \begin{pmatrix} y & x \\ & 1 \end{pmatrix} k$ to

define

$$\int_{Z(\mathbf{A}) \backslash \mathbf{GL}_2(\mathbf{A})} f(g) dg = \int_{F_\mathbf{A}^\times} \int_{\mathbf{A}} \int_K f \left(\begin{pmatrix} y & x \\ & 1 \end{pmatrix} k \right) dk dx \frac{d^\times y}{|y|}.$$

The Petersson inner product of two forms f_1, f_2 on $GL_2(F) \backslash GL_2(\mathbf{A})$ such that $f_1 f_2$ is invariant under $Z(\mathbf{A})$ is defined by

$$\langle f_1, f_2 \rangle_{\text{Pet}} = \int_{Z(\mathbf{A}) \backslash \mathbf{GL}_2(\mathbf{A})} \overline{f_1(g)} f_2(g) dg$$

whenever this converges (this is ensured if either f_1 or f_2 is a cuspform as defined below). It will be convenient to introduce a level-specific inner product on forms f, g of level N :

$$(1.1.2) \quad \langle f, g \rangle_N = \frac{\langle f, g \rangle_{\text{Pet}}}{\mu(N)}$$

where $\mu(N)$ is the measure of $K_0(N)$.

1.2. Fourier expansion. — Let f be a (not necessarily holomorphic) Hilbert modular form. We can expand it as

$$f(g) = C_f(g) + \sum_{\alpha \in F^\times} W_f \left(\begin{pmatrix} \alpha & \\ & 1 \end{pmatrix} g \right)$$

where

$$C_f(g) = D_F^{-1/2} \int_{\mathbf{A}/F} f \left(\begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} g \right) dx,$$

$$W_f(g) = D_F^{-1/2} \int_{\mathbf{A}/F} f \left(\begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} g \right) \mathbf{e}(-x) dx$$

are called the *constant term* and the *Whittaker function* of f respectively. The form f is called *cuspidal* if its constant term C_f is identically zero. The functions of y obtained by restricting the constant term and the Whittaker function to the elements $\begin{pmatrix} y & \\ & 1 \end{pmatrix}$ are called the *Whittaker coefficients*

of f . When f is holomorphic, they vanish unless $y_\infty > 0$ and otherwise have the simple form

$$\begin{aligned} C_f \left(\begin{pmatrix} y & x \\ & 1 \end{pmatrix} \right) &= \tilde{a}^0(f, y) = \psi(y)|y|^{k/2} a(f, 0), \\ W_f \left(\begin{pmatrix} y & x \\ & 1 \end{pmatrix} \right) &= \tilde{a}(f, y) \mathbf{e}_\infty(iy_\infty) \mathbf{e}(x) = \psi(y)|y|^{k/2} a(f, y^\infty d_F) \mathbf{e}_\infty(iy_\infty) \mathbf{e}(x) \end{aligned}$$

for functions $\tilde{a}^0(f, y)$, $\tilde{a}(f, y)$ of $y \in F_{\mathbb{A}}^{\infty, \times}$ which we call the *Whittaker-Fourier coefficients* of f , and a function $a(f, m)$ of the fractional ideals m of F which vanishes on nonintegral ideals whose values are called the **Fourier coefficients** of f .

For any \mathbf{Z} -submodule A of \mathbf{C} , we denote by $M_k(K_1(N), \psi, A)$ the space of holomorphic Hilbert modular forms with Fourier coefficients in A of weight k , level $K_1(N)$, and character ψ ; and by $S_k(K_1(N), \psi, A)$ its subspace of cuspidal forms. When the character ψ is trivial we denote those spaces simply by $M_k(K_0(N), A)$ and $S_k(K_0(N), A)$, whereas linear combinations of forms of level $K_1(N)$ with different characters form the space $M_k(K_1(N), A)$. The notion of Whittaker-Fourier coefficients extends by linearity to the spaces $M_k(K_1(N), \mathbf{C})$.

We can allow more general coefficients: if A is a $\mathbf{Z}[1/N]$ -algebra, we define $S_k(K_0(N), A) = S_k(K_0(N), \mathbf{Z}[1/N]) \otimes A$; this is well-defined thanks to the q -expansion principle [1].

1.3. p -adic modular forms. — Let N, P be coprime ideals of \mathcal{O}_F , ψ a character of conductor dividing N . If f is a holomorphic form of weight k , level $K_1(NP)$ and prime-to- P character ψ (that is, f is a linear combination of forms of level NP and character $\psi\psi'$ with ψ' a character of conductor dividing P), we associate to it the *formal q -expansion coefficients*

$$a_p(f, y^\infty) = \psi^{-1}(y)|y|^{-k/2} \tilde{a}(f, y).$$

If ψ' is trivial we set $a_p(f, m) = a_p(f, y^\infty)$ if m is the ideal $m = y^\infty d_F$.

Let N be an ideal prime to p , ψ a character of level dividing N . Consider the space of classical modular forms $M_k(K_1(Np^\infty, Np^\infty), \overline{\mathbf{Q}})_\psi$ with character whose prime-to- p part is equal to ψ , and endow it with the norm given by the maximum of the p -adic absolute values (for the chosen embedding $\overline{\mathbf{Q}} \hookrightarrow \mathbf{C}_p$) of the Fourier coefficients. Its completion

$$\mathbf{M}_k(K_1(N), \psi, \mathbf{C}_p)$$

of this space is a p -adic Banach space called the space of **p -adic modular forms** of weight k , *tame level* $K_1(N)$ and tame character ψ . We shall view $M_k(K_1(Np^r, \psi, \mathcal{A}))$ as a subset of $\mathbf{M}_k(K_1(N), \psi, \mathcal{A})$ via the q -expansion map.

If \mathcal{A} be a complete \mathbf{Z}_p -submodule of \mathbf{C}_p , we also use the notation $\mathbf{M}_k(K_1(N), \psi, \mathcal{A})$ with obvious meaning, and $\mathbf{S}_k(K_1(N), \psi, \mathcal{A})$ or $\mathbf{S}_k(K_0(N), \mathcal{A})$ (in the case of trivial tame character) for cuspforms; when $k = 2$ we write simply

$$\mathbf{S}_N(\mathcal{A}) = \mathbf{S}_2(K_0(N), \mathcal{A})$$

or just \mathbf{S}_N if $\mathcal{A} = \mathbf{Q}_p$ or $\mathcal{A} = \mathbf{C}_p$ (as understood from context).

1.4. Operators acting on modular forms. — There is a natural action of the group algebra $\mathbf{Q}[\mathrm{GL}_2(\mathbf{A}^\infty)]$ on modular forms induced by right translation. Here we describe several interesting operators arising from this action.

Let m be an ideal of \mathcal{O}_F , $\pi_m \in F_{\mathbf{A}^\infty}^\times$ a generator of $m\widehat{\mathcal{O}}_F$ which is trivial at places not dividing m .

The operator $[m]: M_k(K_1(N), \psi) \rightarrow M_k(K_1(Nm), \psi)$ is defined by

$$(1.4.1) \quad [m]f(g) = \mathbf{N}(m)^{-k/2} f\left(g \begin{pmatrix} 1 & & & \\ & \pi_m & & \\ & & 1 & \\ & & & \ddots \end{pmatrix}\right).$$

It acts on Fourier coefficients by

$$a([m]f, n) = a(f, m^{-1}n).$$

If χ is a Hecke character of F , we denote by $f|\chi$ the form with coefficients

$$a(f|\chi, n) = \chi(n)a(f, n).$$

For any double coset decomposition

$$K_1(N) \begin{pmatrix} \pi_m & & & \\ & 1 & & \\ & & & \\ & & & \ddots \end{pmatrix} K_1(N) = \coprod_i \gamma_i K_1(N),$$

the **Hecke operator** $T(m)$ is defined by the following level-preserving action on forms f in $M_k(K_1(N))$:

$$T(m)f(g) = \mathbf{N}(m)^{k/2-1} \sum_i f(g\gamma_i);$$

For m prime to N , its effect on Fourier coefficients of forms with trivial character is described by

$$a(T(m)f, n) = \sum_{d|(m,n)} \mathbf{N}(d)^{k/2-1} a(f, mn/d^2).$$

When m divides N , we can pick as double coset representatives the matrices $\gamma_i = \begin{pmatrix} \pi_m & c_i \\ & 1 \end{pmatrix}$ for $\{c_i\} \subset \widehat{\mathcal{O}}_F$ a set of representatives for $\mathcal{O}_F/m\mathcal{O}_F$. Then the operator $T(m)$ is more commonly denoted $U(m)$ and we will usually follow this practice. It acts on Fourier coefficients of forms with trivial character by

$$a(U(m)f, n) = \mathbf{N}(m)^{k/2-1} a(f, mn).$$

Let \mathbf{T}_N be the (commutative) subring of $\text{End}_{S_2(K_0(N), \mathbf{Z})}$ generated by the $T(m)$ for m prime to N . A form f which is an eigenfunction of all the operators in \mathbf{T}_N is called a Hecke *eigenform*. It is called a *primitive* form if moreover it is normalised by $a(f, 1) = 1$ and it is a newform (see §1.5 below for the definition) of some level dividing N .

As usual (cf. [41, Lemme 1.10]) we will need the following well-known lemma to ensure the modularity of certain generating functions.

Lemma 1.4.1. — *Let A be a \mathbf{Q} -algebra. For each linear form*

$$a: \mathbf{T}_N \rightarrow A$$

there is a unique modular form in $\oplus_{N'|N} S_k(K_0(N'), A)$ whose Fourier coefficients are given by $a(T(m))$ for m prime to N .

Proof. — In [57, Corollary 3.18], the result is stated and proved when $A = \mathbf{C}$ as a consequence of the existence of a pairing $(T, f) \mapsto a_1(Tf)$ between \mathbf{T}_N and the space of modular forms of interest;

but this pairing is defined over \mathbf{Q} , hence the result is true for $A = \mathbf{Q}$ and by extending scalars for any \mathbf{Q} -algebra A . \square

Atkin-Lehner theory. — For any nonzero ideal M of \mathcal{O}_F , let $W_M \in \mathrm{GL}_2(\mathbf{A}^\infty)$ be a matrix with components

$$(1.4.2) \quad W_{M,v} = \begin{pmatrix} & 1 \\ -\pi_v^{v(M)} & \end{pmatrix} \quad \text{if } v|M, \quad W_{M,v} = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} \quad \text{if } v \nmid M$$

where π_v is a uniformiser at v . We denote by the same name W_M the operator acting on modular forms of level N and trivial character by

$$W_M f(g) = f(g W_M);$$

it is self-adjoint for the Petersson inner product, and when M is prime to N it is proportional to the operator $[M]$ of (1.4.1). On the other hand when M equals N , or more generally M divides N and is coprime to NM^{-1} , the operator W_M is an involution and its action is particularly interesting. In this case, extending the definition to forms of level $K_1(N)$ and character⁽¹⁰⁾ $\psi = \psi_{(M)}\psi_{(NM^{-1})}$ with $\psi_{(C)}$ of conductor dividing C , we have

$$(1.4.3) \quad W_M f(g) = \psi_{(M)}^{-1}(\det g)\psi_{(M)}^{-1}(\pi_M)f(g W_M)$$

where π_M is the idèle with nontrivial components only at $v|M$ and given there by $\pi_v^{v(M)}$. It is easy to check that this definition is independent of the choice of uniformisers. The effect of the W_M -action on newforms is described by Atkin–Lehner theory; we summarise it here (in the case $M = N$), referring to [10] for the details.

Let π be an irreducible infinite-dimensional automorphic representation of $\mathrm{GL}_2(\mathbf{A}_F)$ of central character ψ . Up to scaling, there is a unique *newform* f in the space of π . It is characterised by either of the equivalent properties: (a) it is fixed by a subgroup $K_1(N)$ with N minimal among the N' for which $\pi^{K_1(N')} \neq 0$; (b) its Mellin transform is (a multiple of) the L -function $L(\pi, s)$ of π . In the case of a holomorphic cuspform, this is equivalent to requiring that it belongs to the space of newforms defined in §1.5 below. There is a functional equation relating the L -function $L(s, \pi)$ of π and the L -function $L(1-s, \tilde{\pi})$ of the contragredient representation; as $\tilde{\pi} \cong \psi^{-1} \cdot \pi$, it translates into the following description of the action of W_N on newforms. Suppose that the eigenform $f \in S_k(K_1(N), \psi)$ is a newform in the representation π it generates, then we have

$$(1.4.4) \quad W_N f(g) = (-i)^{[F:\mathbf{Q}]k} \tau(f) f^\rho(g)$$

where f^ρ is the form with coefficients

$$(1.4.5) \quad a(f^\rho, m) = \overline{a(f, m)}$$

and $\tau(f) = \tau(\pi)$ is the central epsilon factor of the functional equation for $L(s, \pi)$.

Trace of a modular form. — The **trace** of a modular form f of level ND and trivial character is the form of level N

$$\mathrm{Tr}_{ND/N}(f)(g) = \sum_{\gamma \in K_0(ND)/K_0(N)} f(g\gamma).$$

⁽¹⁰⁾Notice that a decomposition of ψ as described is only unique up to class group characters (that is, Hecke characters of level one). We will only be using the operator W_M for M a proper divisor of N in a case in which a decomposition is naturally given.

It is the adjoint of inclusion of forms of level N for the rescaled Petersson product:

$$\langle f, \mathrm{Tr}_{ND/N} g \rangle_N = \langle f, g \rangle_{ND}$$

if f has level N and g has level D .

Suppose that D is squarefree and prime to N , in which case we can write $\mathrm{Tr}_D = \mathrm{Tr}_{ND/N}$ without risk of ambiguity. A set of coset representatives for $K_0(ND)/K_0(N)$ is given by elements $\gamma_{j,\delta}$ for $\delta|D$, $j \in \mathcal{O}_{F,v}/\delta \mathcal{O}_{F,v}$, having components

$$\gamma_{j,\delta,v} = \begin{pmatrix} 1 & j \\ & 1 \end{pmatrix} \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} = \frac{1}{\pi_v} \begin{pmatrix} \pi_v & j \\ & 1 \end{pmatrix} \begin{pmatrix} & 1 \\ -\pi_v & \end{pmatrix}$$

at places $v|\delta$, and $\gamma_{j,\delta,v} = 1$ everywhere else. From the second decomposition given just above, if f has weight 2 we obtain

$$(1.4.6) \quad a(\mathrm{Tr}_D(f), m) = \sum_{\delta|D} a(U(\delta)f^{(\delta)}, m) = \sum_{\delta|D} a(f^{(\delta)}, m\delta)$$

where $f^{(\delta)}(g) = f(gW_\delta)$ with W_δ as in (1.4.2).

Remark 1.4.2. — If D is prime to p , the various trace operators Tr_{NDp^r/Np^r} extend to a continuous operator $\mathrm{Tr}_{ND/N}$ on p -adic modular forms of tame level ND . Similarly the operators $[m]$, $T(m)$ and W_m for m prime to Np extend to continuous operators on p -adic modular forms of tame level N .

Ordinary projector. — Let L be a complete subfield of \mathbf{C}_p . Following Hida (see e.g. [23, §3]) we can define for each $\wp|p$ an idempotent

$$e_\wp = \lim_{n \rightarrow \infty} U_\wp^{n!} : S_N(L) \rightarrow S_{N\wp}(L)$$

which is surjective onto $S_{N\wp}^{\wp\text{-ord}}(L)$, the subspace of $S_{N\wp}(L)$ spanned by U_\wp -eigenforms with unit eigenvalue.

Let $P = \prod_{\wp|p} \wp$. Then we similarly have a surjective idempotent

$$e = \prod_{\wp|p} e_\wp : S_N(L) \rightarrow S_{NP}^{\mathrm{ord}}(L),$$

where $S_{NP}^{\mathrm{ord}}(L)$ is the subspace of $S_{NP}(L)$ spanned by simultaneous U_\wp -eigenforms with unit eigenvalue.

1.5. Fourier coefficients of old forms. — As we will study modular forms through their Fourier coefficients, we give here a criterion for recognising the coefficients of certain old forms.⁽¹¹⁾ Let N, P be coprime ideals of \mathcal{O}_F . The space $S_{NP}^{N\text{-old}} \subset S_{NP}$ is the space spanned by forms $f = [d]f'$ for some $1 \neq d|N$ and some cuspform f' of level $N'P$ with $N'|d^{-1}N$. In the case $P = 1$, we define the space of **newforms** of level dividing N to be the orthogonal to the space of old forms for the Petersson inner product. We denote by $\mathbf{S}_N^{\mathrm{old}} \subset \mathbf{S}_N$ the closed subspace generated by the image of $S_{NP}^{N\text{-old}}$ in \mathbf{S}_N .

(The coefficient ring will always be either a finite extension of \mathbf{Q}_p or \mathbf{C}_p , as understood from context when not present explicitly in the notation).

⁽¹¹⁾Cf. [57, §4.4.4].

Let now \mathcal{S} be the space of functions $f: \mathbf{N}_F \rightarrow \mathcal{A}$ modulo those for which there is an ideal M prime to p such that $f(n) = 0$ for all n prime to M . A function $f \in \mathcal{S}$ is called *multiplicative* if it satisfies⁽¹²⁾ $f(mn) = f(m)f(n)$ for all $(m, n) = 1$. For b a multiplicative function, a function f is called an *b -derivative* if it satisfies $f(mn) = b(m)f(n) + b(n)f(m)$ for all $(m, n) = 1$.

Let σ_1 and r be the multiplicative elements of \mathcal{S} defined by

$$\sigma_1(m) = \sum_{d|m} \mathbf{N}(d), \quad r(m) = \sum_{d|m} \varepsilon_{E/F}(d)$$

(where E is a totally imaginary quadratic extension of F of discriminant prime to p).⁽¹³⁾ Let $P = \prod_{\varphi|p} \varphi \subset \mathcal{O}_F$. We define a subspace $\mathcal{D}_N \subset \mathcal{S}$ to be generated by σ_1 , r , σ_1 -derivatives, r -derivatives, and Fourier coefficients of forms in $S_{NP}^{N\text{-old}}$.

Lemma 1.5.1. — *The q -expansion map $S_{NP}^{\text{ord}}/S_{NP}^{N\text{-old}} \rightarrow \mathcal{S}/\mathcal{D}_N$ is injective.*

Proof. — First notice that it is enough to show this when the coefficient ring is a number field L over which S_{NP}^{ord} is defined (it suffices for L to contain all the eigenvalues of the operators T_ℓ ($\ell \nmid NP$) and U_φ on $S_{NP}(L)$). By [57, Proposition 4.5.1], the kernel of $S_{NP}(L)/S_{NP}^{N\text{-old}}(L) \rightarrow \mathcal{S}/\mathcal{D}_N$ is at most generated by $S_{NP}^{p\text{-old}}(L) = \sum_{\varphi|p} S_{NP}^{\varphi\text{-old}}(L)$, the space of forms which are old at some $\varphi|p$. To conclude, it suffices to show that for each $\varphi|p$ we have $I := S_{NP}^{\varphi\text{-old}} \cap S_{NP}^{\varphi\text{-old}} = 0$. The intersection I is stable under the action of \mathbf{T}_{NP} which decomposes it into spaces $I[f_i] \subset S_{NP}[f_i]$ corresponding to eigenforms f_i of level N' or $N'\varphi$ for some $N'|NP\varphi^{-1}$. If f_i has level $N'\varphi$ then $S_{NP}[f_i]$ does not contain φ -oldforms. If f_i has level N' with $\varphi \nmid N'$ then $S_{NP}^{\text{ord}}[f_i]$ is either zero or the line spanned by the ordinary φ -stabilisation of f_i , whereas $S_{NP}^{\varphi\text{-old}}[f_i]$ is the line spanned by $[\varphi]f_i$. We conclude that $I[f_i] = 0$ in all cases. \square

Remark 1.5.2. — The operators U_φ for $\varphi|p$ extend to operators on \mathcal{S} via $U_\varphi f(m) = f(m\varphi)$. The Hecke algebra \mathbf{T}_{NP} acts on the image \mathcal{S}_N of \mathbf{S}_N in $\mathcal{S}/\mathcal{D}_N$.

1.6. The functional l_{f_α} . — Recall from the Introduction that we have fixed an ordinary primitive Hilbert modular newform f of level $K_0(N)$. If α_φ is the unit root of the φ^{th} Hecke polynomial of f , β_φ is the other root, and the operator $[\varphi]$ is as in (1.4.1), then the *p -stabilisation* of f is

$$f_\alpha = \prod_{\varphi|p} (1 - \beta_\varphi [\varphi]) f,$$

a form of level $K_0(N \prod_{\varphi|p} \varphi)$ satisfying $U_\varphi f_\alpha = \alpha_\varphi f_\alpha$ for all $\varphi|p$.

We define a functional, first introduced by Hida, which plays the role of projection onto the f -component. Both sides of our main formula will be images of p -adic modular forms under this operator.

Let P be an ideal of \mathcal{O}_F divisible exactly by the primes $\varphi|p$. For a form $g \in M_2(K_0(NP))$ with $r \geq 1$, let

$$l_{f_\alpha}(g) = \frac{\langle W_{NP} f_\alpha^\rho, g \rangle}{\langle W_{NP} f_\alpha^\rho, f_\alpha \rangle}.$$

⁽¹²⁾This relation and the following are of course to be understood to hold in \mathcal{S} .

⁽¹³⁾We will see below that σ_1 and r are the Fourier coefficients of weight 1 Eisenstein series and theta series.

Let $L \subset \overline{\mathbf{Q}}_p$ be the extension of \mathbf{Q}_p generated by $a(f, m)$ for all ideals m and α_\wp for $\wp|p$.

Lemma 1.6.1 (Hida). — *The above formula defines a linear functional*

$$l_{f_\alpha} : M_2(K_0(Np^\infty), L) \rightarrow L$$

satisfying:

1. On $M_2(K_0(N), L)$ we have

$$l_{f_\alpha} = \prod_{\wp|p} \left(1 - \frac{N_\wp}{\alpha_\wp^2} \right)^{-1} \mathbf{1}_f$$

where $\mathbf{1}_f(g) = \langle f, g \rangle / \langle f, f \rangle$.

2. On $M_2(K_0(N\wp^r))$ we have, for each nonnegative $t \leq r-1$,

$$l_{f_\alpha} \circ U_\wp^t = \alpha_\wp(f)^t l_{f_\alpha}.$$

3. If each $\iota_p(\alpha_\wp)$ is a p -adic unit, l_{f_α} admits a continuous extension to p -adic modular forms still denoted

$$l_{f_\alpha} : \mathbf{M}_N(L) \rightarrow L.$$

Proof. — See [23, Lemma 9.3] where the well-definedness of the functional and its extension to p -adic modular forms are proved more generally for Hida families. For part 1 the computation is the same as in the case of elliptic modular forms: see [40] or [22, §4]. \square

Some quotient spaces. — Let $\overline{\mathcal{F}} = \mathcal{S} / \mathcal{D}_N$. The ordinary projection operator e is not defined on all arithmetic functions; however its kernel $\text{Ker}(e)$ is a well-defined subspace of \mathcal{S} . We define

$$\overline{\mathcal{F}}^{\text{ord}} := \mathcal{S} / \mathcal{D}_N + \text{Ker}(e).$$

The quotient map $\overline{\mathcal{F}} \rightarrow \overline{\mathcal{F}}^{\text{ord}}$ is clearly injective when restricted to the image of S_{NP}^{ord} , where $P = \prod_{\wp|p} \wp$. Then we denote by $\overline{\mathcal{F}}_N^{\text{ord}}$ the image of S_{NP}^{ord} in either $\overline{\mathcal{F}}$ or $\overline{\mathcal{F}}^{\text{ord}}$. It is also identified with the common image of S_{NP} and S_N in $\overline{\mathcal{F}}^{\text{ord}}$. We denote by $\overline{\mathcal{F}}_N^{p\text{-adic}} \subset \overline{\mathcal{F}}$ the image of S_N .

We obtain a commutative diagram (where L is as usual any sufficiently large finite extension of \mathbf{Q}_p):

$$(1.6.1) \quad \begin{array}{ccc} S_N(L) & \longrightarrow & \overline{\mathcal{F}}_N^{p\text{-adic}}(L) \\ \downarrow e & & \downarrow \\ S_{NP}^{\text{ord}}(L) / S_{NP}^{N\text{-old}} & \xrightarrow{\sim} & \overline{\mathcal{F}}_N^{\text{ord}}(L) \xrightarrow{l_{f_\alpha}} L \end{array}$$

where the right-hand vertical map is the restriction of the quotient $\overline{\mathcal{F}} \rightarrow \overline{\mathcal{F}}^{\text{ord}}$, and the bottom horizontal map is an isomorphism by Lemma 1.5.1.

2. Theta measure

We construct a measure on the Galois group of the maximal abelian extension of E unramified outside p with values in p -adic theta series, and compute its Fourier expansion.

2.1. Weil representation. — We first define the Weil representation. See [7, §4.8] for an introduction, and [52] or [55] for our conventions on the representation for similitude groups.

Local setting. — Let $V = (V, q)$ be a quadratic space over a local field F of characteristic not 2, with a quadratic form q ; we choose a nontrivial additive character \mathbf{e} of F . For simplicity we assume V has even dimension. For $u \in F^\times$, we denote by V_u the quadratic space (V, uq) . We let $\mathbf{GL}_2(F) \times \mathbf{GO}(V)$ act on the space $\mathcal{S}(V \times F^\times)$ of Schwartz functions as follows (here $\nu: \mathbf{GO}(V) \rightarrow \mathbf{G}_m$ denotes the similitude character):

- $r(h)\phi(t, u) = \phi(h^{-1}t, \nu(h)u)$ for $h \in \mathbf{GO}(V)$;
- $r(n(x))\phi(t, u) = \mathbf{e}(x u q(t))\phi(t, u)$ for $n(x) = \begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} \in \mathbf{GL}_2$;
- $r\left(\begin{pmatrix} a & \\ & d \end{pmatrix}\right)\phi(t, u) = \chi_V(a) |a/d|^{\frac{\dim V}{4}} \phi(at, d^{-1}a^{-1}u)$;
- $r(w)\phi(x, u) = \gamma(V_u)\hat{\phi}(x, u)$ for $w = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix}$.

Here χ_V is the quadratic character associated to V , $\gamma(V_u)$ is a certain square root of $\chi(-1)$, and $\hat{\phi}$ denotes the Fourier transform in the first variable

$$\hat{\phi}(x, u) = \int_V \phi(y, u) \mathbf{e}(-u\langle x, y \rangle) dy$$

where \langle , \rangle is the bilinear form associated to q and dy is the self-dual Haar measure.

Global setting. — Given a quadratic space (V, q) over a global field F of characteristic not 2 (and a nontrivial additive character $\mathbf{e}: F \setminus \mathbf{A}_F \rightarrow \mathbf{C}^\times$), the Weil representation is the restricted tensor product r of the associated local Weil representations, with spherical functions $\phi_v(t, u) = \mathbf{1}_{\mathcal{Y}_v \times \mathcal{O}_{F,v}^\times}(x, u)$ for some choice of lattices $\mathcal{Y}_v \subset V(F_v)$.

The case of interest to us is the following: F is a totally real number field, $V = (E, \mathfrak{N})$ is given by a quadratic CM extension E/F with the norm form $\mathfrak{N} = N_{E/F}$ and the lattices $\mathcal{O}_{E,v} \subset E_v$, and the additive character \mathbf{e} is the standard one. We denote $G = \mathbf{GL}_2$, $H = \mathbf{GSO}(V)$, two algebraic groups defined over F ; we have $H \cong \text{Res}_{E/F} \mathbf{G}_m$. In this case we have

$$\chi_V = \varepsilon_{E/F} = \varepsilon,$$

where $\varepsilon_{E/F}$ is the quadratic character of $F_\mathbf{A}^\times$ associated to the extension E/F . The self-dual measure on E_v is the one giving $\mathcal{O}_{E,v}$ volume $|\mathcal{O}_{E,v}/\mathfrak{D}_v|^{-1/2}$ where \mathfrak{D}_v is the relative different. Moreover the constant γ can be explicitly described (see [8, §§ 38.6, 30.4, 23.5]): in the case $v|\Delta_{E/F}$, which is the only one we will be using, such description is in terms of a local Gauß sum $\varkappa(v)$:

$$(2.1.1) \quad \gamma(E_v, u\mathfrak{N}) = \varepsilon_v(u)\varkappa(v) = \varepsilon_v(u)|\pi_v|^{1/2} \sum_{x \in (\mathcal{O}_{F,v}/\pi_v \mathcal{O}_{F,v})^\times} \varepsilon(x/\pi_v) \mathbf{e}_v(x/\pi_v).$$

Notice that our $\varkappa(v)$ is the *inverse* of the quantity denoted by the same name in [57, Proposition 3.5.2].

2.2. Theta series. — We define the **theta kernel** to be

$$\theta_\phi(g, h) = \sum_{(t, u) \in V \times F^\times} r(g, h)\phi(t, u)$$

which is an automorphic form for the group $\mathbf{GL}_2(F)\backslash\mathbf{GL}_2(\mathbf{A}_F) \times \mathbf{GO}(V)\backslash\mathbf{GO}(V_{\mathbf{A}_F})$.

If \mathcal{W} is an automorphic function for H which is trivial at infinity (which is the same thing as a linear combination of finite order Hecke characters of E), we define the **theta series**⁽¹⁴⁾

$$\theta_\phi(\mathcal{W})(g) = \int_{H(F)\backslash H(\mathbf{A}_F)} \mathcal{W}(b^{-1})\theta_\phi(g, b) db$$

which is an automorphic form on G . Here the measure db is the product of the measure on $H(\mathbf{A}^\infty)$ which gives volume 1 to the compact $U_0 = \widehat{\mathcal{O}}_E^\times$, and any fixed measure⁽¹⁵⁾ on $H(\mathbf{A}_\infty)$.

Let us explain how to explicitly compute the integral in our situation. For each open compact subgroup $U \subset H(\mathbf{A}_F^\infty) = E_{\mathbf{A}^\infty}$, we have exact sequences

$$1 \rightarrow \mathcal{O}_{E,U}^\times \backslash UE_\infty^\times \rightarrow E^\times \backslash E_{\mathbf{A}}^\times \rightarrow E^\times U \backslash E_{\mathbf{A}^\infty}^\times \rightarrow 1$$

and

$$1 \rightarrow \mu(U) \backslash UE_\infty^1 \rightarrow \mathcal{O}_{E,U}^\times \backslash UE_\infty^\times \xrightarrow{\mathfrak{N}_\infty} N(\mathcal{O}_{E,U}^\times) \backslash F_\infty^+ \rightarrow 1.$$

The notation used is the following: $\mathcal{O}_{E,U}^\times = E^\times \cap U \supset \mu(U) =$ the subset of roots of unity, $\mathfrak{N}_\infty: E_\infty^\times \rightarrow F_\infty^+$ is the norm map at the infinite places and E_∞^1 is its kernel.

We can choose a splitting ι of the first sequence, for example

$$\iota: E^\times U \backslash E_{\mathbf{A}^\infty}^\times \cong E^\times U \backslash (E_{\mathbf{A}}^\times)^{1,\parallel} \hookrightarrow E^\times \backslash E_{\mathbf{A}}^\times,$$

where $(E_{\mathbf{A}}^\times)^{1,\parallel}$ denotes the set of idèles of adelic norm 1 with infinity component $h_\infty = (b, \dots, b)$ for some real number $b > 0$ and the isomorphism is the unique one which gives the identity once composed with projection onto the finite part.

We begin to expand the series, evaluating the integral as explained above and exploiting the fact that the action of $H(F_\infty) = E_\infty^\times$ on $\phi(t, u)$ factors through the norm. We take U to be small enough so that \mathcal{W} and ϕ are invariant under U , and denote

$$\bar{\phi}_v(t, u) = \int_{H(F_v)} r(b)\phi_v(t, u) db \quad \text{if } v|\infty$$

and $\bar{\phi} = \prod_{v|\infty} \phi_v \prod_{v|\infty} \bar{\phi}_v$. A specific choice of $\bar{\phi}_v$ will be made shortly: for the moment we just record, and use in the following computation, that we will take $u \mapsto \bar{\phi}_v(t, u)$ to be supported on \mathbf{R}^+ .

We have

$$\begin{aligned} \theta_\phi(\mathcal{W})(g) &= \int_{E^\times \backslash E_{\mathbf{A}}^\times} \mathcal{W}(b^{-1})\theta_\phi(g, b) db \\ &= \omega_U^{-1} \int_U \int_{E_\infty^1} \int_{\mathfrak{N}(\mathcal{O}_{E,U}^\times) \backslash F_\infty^+} \int_{E^\times U \backslash E_{\mathbf{A}^\infty}^\times} \mathcal{W}(\iota(a)^{-1}) \sum_{(t,u) \in E \times F^\times} r(g, \iota(a)b)\phi(t, u) da db \end{aligned}$$

⁽¹⁴⁾The reason for taking $\mathcal{W}(b^{-1})$ rather than $\mathcal{W}(b)$ is that we want $\theta_\phi(\mathcal{W})$ to be the series classically denoted $\Theta(\mathcal{W})$ for a suitable choice of ϕ - this will be clear from the computations below.

⁽¹⁵⁾There will be no ambiguity since later we will choose ϕ_∞ to be again any fixed Schwartz function, whose integral over $H(\mathbf{A}_\infty)$ with respect to the chosen measure is a specified function $\bar{\phi}_\infty$.

Here $w_U = |\mu(U)|$ and dh denotes the measure on $U \times E_\infty^1 \times F_\infty^+ = U \times H(F_\infty)$. We partially collapse the integral over $\mathfrak{N}(\mathcal{O}_{E,U}^\times) \backslash F_\infty^+$ and the sum over $u \in F^\times$ and use our choice of ϕ_∞ to get

$$\begin{aligned}
 &= w_U^{-1} \text{vol}(U) \int_{E^\times U \backslash E_{A_\infty}^\times} \mathcal{W}(\iota(a)^{-1}) \sum_{u \in \mathfrak{N}(\mathcal{O}_{E,U}^\times) \backslash F^+} \sum_{t \in E} r(g, \iota(a)) \bar{\phi}(t, u) da \\
 (2.2.1) \quad &= w_U^{-1} \frac{h}{h_U} \int_{E^\times U \backslash E_{A_\infty}^\times} \mathcal{W}(\iota(a)^{-1}) \nu_U \sum_{u \in \mathfrak{N}(\mathcal{O}_{E,U}^\times) \backslash F^+} \sum_{t \in E} r(g, \iota(a)) \bar{\phi}(t, u) da
 \end{aligned}$$

Here in the last step we have defined $\nu_U = [\mathfrak{N}(\mathcal{O}_E^\times) : \mathfrak{N}(\mathcal{O}_{E,U}^\times)]$ and computed $\text{vol}(U) = \text{vol}(U_0)(h/h_U)(w_U/w)\nu_U^{-1}$, where $U_0 = \widehat{\mathcal{O}}_E^\times$, $h_U = |E^\times U \backslash E_{A_\infty}^\times|$, $h = h_{U_0}$, $w = w_{U_0}$. Recall that our measure satisfies $\text{vol}(U_0) = 1$. The remaining integral is just a finite sum.

The sum over u is actually finite owing to the integrality constraints imposed by ϕ at finite places.⁽¹⁶⁾

2.3. Theta measure. — We define a measure with values in p -adic modular forms on the group

$$\mathcal{G}' = \text{Gal}(E'_\infty/E) \cong \varprojlim \overline{E^\times U_{p^n}} \backslash E_{A_\infty}^\times$$

where the over line denotes closure and E'_∞ is the maximal abelian extension of E unramified outside p , that is, the union of the ray class fields of E of p -power ray $U_{p^n} = \prod_v \{\text{units} \equiv 1 \pmod{p^n \mathcal{O}_{E,v}}\}$ and the isomorphism is given by class field theory. The topology is the profinite topology.

Recall that a **measure** on a topological space \mathcal{G} with values in a p -adic Banach space \mathbf{M} is a \mathbf{C}_p -linear functional

$$\mu : \mathcal{C}(\mathcal{G}, \mathbf{C}_p) \rightarrow \mathbf{M}$$

on continuous \mathbf{C}_p -valued functions, which is continuous (equivalently, bounded) with respect to the sup norm on $\mathcal{C}(\mathcal{G}, \mathbf{C}_p)$. The linearity property will be called *distributional property* in what follows. The boundedness property will in each case at hand be verified on the set of p -adic characters of \mathcal{G} , which in our cases generates the whole of $\mathcal{C}(\mathcal{G}, \mathbf{C}_p)$ (classically, the continuity of μ goes under the name of *abstract Kummer congruences* for μ).

When $\mathbf{M} = \mathbf{M}_0 \otimes_{\mathbf{Q}_p} \mathbf{C}_p$ for a p -adic Banach space \mathbf{M}_0 over \mathbf{Q}_p , the measure μ is said to be *defined over \mathbf{Q}_p* if $\mu(\mathcal{W}) \in \mathbf{M}_0 \otimes_{\mathbf{Q}_p} \mathbf{C}_p(\mathcal{W})$ whenever the function \mathcal{W} on \mathcal{G} has values in $\mathbf{Q}_p(\mathcal{W}) \subset \overline{\mathbf{Q}_p} \subset \mathbf{C}_p$.

Definition 2.3.1. — The **theta measure** $d\Theta$ on \mathcal{G}' is defined by

$$\Theta(\mathcal{W}) = \int_{\mathcal{G}'} \mathcal{W}(\sigma) d\Theta(\sigma) = \theta_\phi(\mathcal{W}),$$

for any function $\mathcal{W} : \mathcal{G}' \rightarrow \overline{\mathbf{Q}}$ factoring through a finite quotient of \mathcal{G}' , where the function ϕ is chosen as follows:

$$- \text{ for } v \nmid p\infty, \phi_v(t, u) = \mathbf{1}_{\mathcal{O}_{E,v}}(t) \mathbf{1}_{d_{F_v}^{-1,\times}}(u);$$

⁽¹⁶⁾We will see this in more detail shortly. We are also using the definition of $\bar{\phi}_\infty$ in order to freely replace the sum over $u \in F^\times$ with a sum over $u \in F^+$ – in fact a slight variation would be necessary when $\det g_\infty \notin F_\infty^+$, but this is a situation we won't encounter.

– for $v|p$,

$$\phi_v(t, u) = [\mathcal{O}_{E,v}^\times : U'_v] \mathbf{1}_{U'_v}(t) \mathbf{1}_{d_{F_v}^{-1,\times}}(u),$$

where $U'_v \subset \mathcal{O}_{E,v}^\times$ is any small enough compact set – that is, $U'_v \subset U_v$ if \mathcal{W} is invariant under $U = \prod_v U_v$, and the definition does not depend on the choice of U_v . (In practice, we will choose $U'_v = U_v$ if U_v is maximal with respect to the property just mentioned.)

– for $v|\infty$, $\phi_v(t, u)$ is a Schwartz function such that

$$\int_{H(F_v)} r(h) \phi_v(t, u) dh = \bar{\phi}_v(t, u) = \mathbf{1}_{\mathbf{R}^+}(u) \exp(-2\pi u N(t)).$$

(See [55, 4.1] for more details on this choice.)

In Corollary 2.4.2 below we will show that this in fact defines a measure on \mathcal{G}' with values in p -adic Hilbert modular forms of weight one, tame level $\Delta_{E/F}$ and character ε .

2.4. Fourier expansion of the theta measure / I. – We compute the Fourier expansion of the theta measure on \mathcal{G}' , carrying on the calculation started in §2.2.

In the case where $g = \begin{pmatrix} y & x \\ & 1 \end{pmatrix}$ with $y_\infty > 0$, the sum over (u, t) in (2.2.1) evaluates to

$$(2.4.1) \quad \varepsilon(y) |y|^{1/2} \sum_{u,t} \phi^\infty(a^{-1}yt, \mathfrak{N}(a)y^{-1}u) \mathbf{e}_\infty(iy_\infty u N(t)) \mathbf{e}(xu \mathfrak{N}(t)).$$

Then we compute the sum of this expression over the finite quotient \mathcal{G}'_U of \mathcal{G}' , with $\mathcal{G}'_U \cong E^\times U \backslash E_{A^\infty}^\times$.

We assume \mathcal{W} is a character so $\mathcal{W}(a^{-1}) = \overline{\mathcal{W}}(a)$ where $\overline{\mathcal{W}} = \mathcal{W}^{-1}$.

First we pre-compute the product of all the constants appearing in the theta series of (2.2.1), including the one from ϕ – we take

$$\phi_v(t, u) = [\mathcal{O}_{E,v}^\times : U_v] \mathbf{1}_{U_v}(t) \mathbf{1}_{\mathcal{O}_F^\times}(u),$$

so:

$$\begin{aligned} \omega \frac{h}{h_U} \nu_U[\mathcal{O}_{E,v}^\times : U_v] &= \omega[\mathcal{O}_E^\times \backslash \widehat{\mathcal{O}}_{E,v}^\times : \mathcal{O}_{E,U}^\times \backslash U]^{-1} [\mathfrak{N}(\mathcal{O}_E^\times) : \mathfrak{N}(\mathcal{O}_{E,U}^\times)]^{-1} [\widehat{\mathcal{O}}_E^\times : U] \\ &= \omega[\mu(\mathcal{O}_E) : \mu(U)] = \omega_U^{-1}. \end{aligned}$$

This computation together with (2.2.1), (2.4.1) gives

$$\begin{aligned} \Theta(\mathcal{W}) &= \varepsilon(y) |y|^{\frac{1}{2}} \omega_U^{-1} \sum_{a \in E^\times U \backslash E_{A^\infty}^\times} \overline{\mathcal{W}}(a) \sum_{t \in E, u \in \mathfrak{N}(\mathcal{O}_{E,U}^\times) \backslash F^+} \phi^{p^\infty}(a^{-1}yt, \mathfrak{N}(a)y^{-1}u) \\ &\quad \times \mathbf{1}_{\mathcal{O}_{E,U,p}^\times}(a^{-1}yt) \mathbf{1}_{d_{F_p}^{-1,\times}}(\mathfrak{N}(a)y^{-1}u) \mathbf{e}_\infty(iy_\infty u \mathfrak{N}(t)) \mathbf{e}(xu \mathfrak{N}(t)) \\ &= \varepsilon(y) \overline{\mathcal{W}}(y) |y|^{\frac{1}{2}} \omega_U^{-1} \sum_{a \in E^\times U \backslash E_{A^\infty}^\times} \overline{\mathcal{W}}(a) \sum_{t \in E, u \in \mathfrak{N}(\mathcal{O}_{E,U}^\times) \backslash F^+} \mathbf{1}_{\widehat{\mathcal{O}}_{E,U}^\times \cap \mathcal{O}_{E,U,p}^\times}(a^{-1}t) \\ &\quad \times \mathbf{1}[\mathfrak{N}(a)yu \mathcal{O}_F = d_F^{-1}] \mathbf{e}_\infty(iy_\infty u \mathfrak{N}(t)) \mathbf{e}(xu \mathfrak{N}(t)) \end{aligned}$$

where we have made the change of variable $a \rightarrow ay^\infty$.

Now we make the substitution $u\mathfrak{N}(t) = \xi$ and observe that the contribution to the ξ^{th} term is equal to zero if $(\xi \gamma d_F, p) \neq 1$, and otherwise it equals $\overline{\mathcal{W}}(a)$ times the cardinality of the set

$$R_{a^{-1}}(\xi, \gamma) = \left\{ (t, u) \in \mathcal{O}_E \times F^+ \mid t \in U_p, u\mathfrak{N}(t) = \xi, \mathfrak{N}(t/a)\mathcal{O}_F = \xi \gamma d_F \right\} / \mathfrak{N}(\mathcal{O}_{E,U}^\times),$$

which admits a surjection $\pi : (t, u) \mapsto a^{-1}t\mathcal{O}_E$ to the set $\mathfrak{r}_{a^{-1}}(\xi \gamma d_F)$ of ideals $\mathfrak{b} \subset \mathcal{O}_E$ in the U -class a^{-1} , whose norm is $\mathfrak{N}(\mathfrak{b}) = \xi \gamma d_F$. The fibres of π are in bijection with $\mathcal{O}_{E,U}^\times / \mathfrak{N}(\mathcal{O}_{E,U}^\times)$ which has cardinality w_U . We deduce the following description of the Fourier coefficients of $\Theta(\mathcal{W})$.

Proposition 2.4.1. — *The series $\Theta(\mathcal{W})$ belongs to $S_1(K_1(\Delta(\mathcal{W})), \varepsilon \overline{\mathcal{W}}|_{F^\times})$, where $\Delta(\mathcal{W}) = \Delta \mathfrak{N}(\mathfrak{f}(\mathcal{W}))$. Its Fourier coefficients are given by*

$$a(\Theta(\mathcal{W}), m) = \sum_{\substack{\mathfrak{b} \subset \mathcal{O}_E \\ \mathfrak{N}(\mathfrak{b})=m}} \mathcal{W}(\mathfrak{b}) = r_{\mathcal{W}}(m)$$

for $(m, p) = 1$ and vanish for $(m, p) \neq 1$.

Corollary 2.4.2. — *The functional Θ of Definition 2.3.1 is a measure on \mathcal{G}^l with values in $S_1(K_1(\Delta), \varepsilon)$, defined over \mathbf{Q}_p .*

Proof. — The distributional property is obvious from the construction or can be seen from the q -expansion given above, from which boundedness is also clear. Cf. also [24, Theorem 6.2], where a slightly different theta measure is constructed. \square

Lemma 2.4.3. — *The theta series admits a functional equation*

$$W_{\Delta(\mathcal{W})} \Theta(\mathcal{W}) = (-i)^{[F:\mathbf{Q}]} \overline{\mathcal{W}}(\mathfrak{D}_E) \tau(\overline{\mathcal{W}}) \Theta(\overline{\mathcal{W}})$$

where \mathfrak{D}_E is the relative different and $\tau(\overline{\mathcal{W}}) = \prod_{v|p} \tau(\overline{\mathcal{W}}_v)$ with

$$\tau(\overline{\mathcal{W}}_v) = |\pi_v|^{-c/2} \int_{E_v^\times} \overline{\mathcal{W}}_v(h_v) \mathbf{e}_v(-\text{Tr}_{E_v/F_v}(h_v)) dh_v$$

if the relative norm of the conductor of $\overline{\mathcal{W}}_v$ is $\pi_v^c \mathcal{O}_{F,v}$.

Proof. — Let

$$\begin{aligned} \phi_{\mathcal{W}}(g, t, u) &= \int_{H(F) \backslash H(\mathbf{A})} \mathcal{W}(h^{-1}) r(g, h) \phi(t, u) dh \\ \phi'_{\mathcal{W}}(g, t, u) &= \varepsilon \mathcal{W}(\pi_{\Delta(\mathcal{W})}) \int_{H(F) \backslash H(\mathbf{A})} \mathcal{W}(h^{-1}) r(g W_{\Delta(\mathcal{W})}, h) \phi(t, u) dh \end{aligned}$$

for $(t, u) \in E_{\mathbf{A}} \times F_{\mathbf{A}}^\times$. The behaviour in g is through the Weil representation.

Then we have

$$\begin{aligned} W_{\Delta(\mathcal{W})} \Theta(\mathcal{W})(g) &= \varepsilon \mathcal{W}(\det(g)) \sum_{(t,u) \in E \times F^\times} \phi'_{\mathcal{W}}(g, t, u) \\ \Theta(\overline{\mathcal{W}})(g) &= \sum_{(t,u) \in E \times F^\times} \phi_{\overline{\mathcal{W}}}(g, t, u) \end{aligned}$$

so that the lemma follows if we show that for all $(t, u) \in E_{\mathbf{A}} \times F_{\mathbf{A}}^{\times}$

$$(2.4.2) \quad \varepsilon \mathcal{W}(\det(g)) \phi'_{\mathcal{W}}(g, t, u) = (-i)^{[F:\mathbb{Q}]} \overline{\mathcal{W}}(\mathfrak{D}_E) \tau(\overline{\mathcal{W}}) \varepsilon \mathcal{W}(u) \phi_{\overline{\mathcal{W}}}(g, \bar{t}, u)$$

where \bar{t} is the conjugate of t under the nontrivial automorphism of E over F . We write

$$\tilde{\tau}(\overline{\mathcal{W}}) = (-i)^{[F:\mathbb{Q}]} \overline{\mathcal{W}}(\mathfrak{D}_E) \tau(\overline{\mathcal{W}})$$

for short.

We claim that it suffices to prove (2.4.2) for $g = 1$. Indeed it is clear that this implies the same result for all $g \in \mathbf{SL}_2(\mathbf{A})$ by acting via the Weil representation on both sides (viewed as functions of (t, u)). Then it suffices to verify it for the elements of the form $d(y) = \begin{pmatrix} 1 & \\ & y \end{pmatrix}$:

$$\begin{aligned} \varepsilon(y) \mathcal{W}(y) r(d(y)) \phi'_{\mathcal{W}}(1, t, u) &= \tilde{\tau}(\overline{\mathcal{W}}) \varepsilon(y) \mathcal{W}(y) r(d(y)) [\varepsilon \mathcal{W}(u) \phi_{\overline{\mathcal{W}}}(1, \bar{t}, u)] \\ &= \tilde{\tau}(\overline{\mathcal{W}}) \varepsilon \mathcal{W}(y) \varepsilon \mathcal{W}(y^{-1}u) r(d(y)) \phi_{\overline{\mathcal{W}}}(1, \bar{t}, u) = \tilde{\tau}(\overline{\mathcal{W}}) \varepsilon(u) \mathcal{W}(u) \phi_{\overline{\mathcal{W}}}(d(y), \bar{t}, u). \end{aligned}$$

We now prove (2.4.2) for $g = 1$, thus dropping g from the notation. We can write

$$\begin{aligned} \phi'_{\mathcal{W}}(t, u) &= \int_{H(F) \backslash H(\mathbf{A}^{p\Delta})} \mathcal{W}(h_0^{-1}) r(1, h) \phi^{p\Delta}(t, u) dh_0 \\ &\quad \prod_{v|p\Delta} \mathcal{W}(\pi_v^{c_v}) \int_{H(F_v)} \mathcal{W}(h_v^{-1}) r(W_{\pi_v^{c_v}}, 1) \phi(h_v^{-1}t, v(h_v u)) dh_v \end{aligned}$$

where c_v is the appropriate exponent. We can rewrite this as

$$\phi'_{\mathcal{W}}(t, u) = \phi'^{\Delta p}_{\mathcal{W}}(t, u) \prod_{v|p\Delta} \phi'_{\mathcal{W}, p}(t, u)$$

with obvious notation. A similar factorisation holds for $\phi_{\mathcal{W}}(t, u)$.

For $v \nmid p$ we have, by the explicit description of ϕ_v (dropping the subscripts v):

$$\begin{aligned} r(h) \phi(t, u) &= \phi(h^{-1}t, v(h)u) = \phi(\pi_{d_F} u \bar{h}t, v(h)u) \\ &= \phi(\pi_{d_F} u \bar{h} \bar{t}, v(h)^{-1} u^{-1} \pi_{d_F}^{-2}) = r((\pi_{d_F} u \bar{h})^{-1}) \phi(\bar{t}, u). \end{aligned}$$

A change of variable and integration over $H(F) \backslash H(\mathbf{A}^{\Delta p})$ then gives

$$(2.4.3) \quad \phi'^{\Delta p}_{\mathcal{W}}(t, u) = \mathcal{W}^{\Delta p}(u d_F) \varepsilon^{\Delta p}(d_F) \varepsilon^{\Delta p}(u) \phi'_{\mathcal{W}}(\bar{t}, u)$$

where the added factor $\varepsilon^{\Delta p}(d_F) \varepsilon^{\Delta p}(u) = 1$ if either side is nonzero (recall that $\phi(t, u) = 0$ unless $u^{-1} \in d_F^{\Delta p, \times}$).

For $v|p$ we have by (2.5.1) below and the previous argument

$$\varepsilon(\pi) r(W_{\pi}, h) (\pi) \phi(t, u) = \varepsilon(u) \chi(v) \phi(h^{-1}t \pi_{\mathfrak{D}}, \pi^{-1}v(h)u) = \varepsilon(u) \chi(v) r(u^{-1}h^{-1}\pi_{\mathfrak{D}}) \phi(\bar{t}, u)$$

where $\pi_{\mathfrak{D}} \in \mathcal{O}_{E, v}^{\times}$ is a generator of the local relative different of E_v/F_v . After change of variable and integration, we obtain

$$(2.4.4) \quad \phi'_{\mathcal{W}, v}(t, u) = \varepsilon \mathcal{W}(u) \chi(v) \overline{\mathcal{W}}_v(\mathfrak{D}) \phi_{\overline{\mathcal{W}}, v}(\bar{t}, u).$$

For $v|p$ we have

$$\begin{aligned} & \mathcal{W}(\pi^c) \int_{H(F_v)} \mathcal{W}(h^{-1}) r(h, w_{\pi^c}) \phi(t, u) d^\times h \\ &= |\pi|^{-c/2} \int_{E^\times} \int_E \overline{\mathcal{W}}(\pi^{-c} h) \mathbf{e}(-\pi^{-c} u \nu(h) \mathrm{Tr}(h^{-1} t \bar{\xi})) \phi(\xi, \pi^{-c} \nu(h) u) d\xi d^\times h \end{aligned}$$

Using the fact that $\phi(\xi, u) d\xi = \phi(\xi, u) d^\times \xi$, and a change of variables $\zeta = \pi^{-c} u h \xi \bar{t}$, this equals

$$\mathcal{W}(u) \tau(\overline{\mathcal{W}}) \int_{E^\times} \mathcal{W}(\xi \bar{t}) \phi(\xi, \nu(t \xi) u) d^\times \xi$$

after integration, where the new second argument in ϕ gives the condition for the integral in $d\zeta$ to be nonzero. We observe that $\phi(\xi) = \phi(\xi^{-1})$ so that with the the new variable $h' = \xi \bar{t}$, and reintroducing v in the notation, this can be rewritten as

$$\mathcal{W}_v(u) \tau(\overline{\mathcal{W}}_v) \int_{E_v^\times} \overline{\mathcal{W}}_v(h'_v)^{-1} \phi(h'^{-1} \bar{t}, \nu(h'_v) u) d^\times h'_v$$

so that

$$(2.4.5) \quad \phi'_{\mathcal{W}_v}(t, u) = \varepsilon_v(d_F) \varepsilon_v(u) \mathcal{W}_v(u) \tau(\overline{\mathcal{W}}_v) \phi_{\overline{\mathcal{W}}_v}(\bar{t}, u).$$

Putting together (2.4.3), (2.4.4), (2.4.5) and using the formula $\prod_{v|\Delta} \chi(v) = (-i)^{[F:\mathbb{Q}]} \varepsilon(d_F)$ from [57, p. 127],⁽¹⁷⁾ we obtain (2.4.2) as desired. \square

2.5. Fourier expansion of the theta measure / II. — For later use in computing the trace of the convolution of the theta measure with the Eisenstein measure (defined below), we need to consider the expansion of $\Theta(\mathcal{W})^{(\delta)}(g) = \Theta(\mathcal{W})(g W_\delta)$ for $g = \begin{pmatrix} y & x \\ & 1 \end{pmatrix}$; for such a g we have

$$\begin{pmatrix} y & x \\ & 1 \end{pmatrix} W_\delta = \begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} \begin{pmatrix} y & \\ & \pi_\delta \end{pmatrix} w_\delta$$

where π_δ is an idèle with components π_v at $v|\delta$ and 1 everywhere else. Here π_v is a uniformiser chosen to satisfy $\varepsilon(\pi_v) = 1$.

The modular form $\Theta(\mathcal{W})^\delta$ can be expanded in the same way as in §2.4, except that for $v|\delta$ we need to replace $\phi_v(t, u) = \mathbf{1}_{\mathcal{O}_{E,v}}(t) \mathbf{1}_{d_F^{-1,\times}}(u)$ by

$$(2.5.1) \quad \begin{aligned} W_\delta \phi_v(t, u) &= \varepsilon_v(\pi_v) \begin{pmatrix} 1 & \\ & \pi_v \end{pmatrix} \gamma(u) \widehat{\mathbf{1}}_{\mathcal{O}_{E,v}}(t) \mathbf{1}_{d_F^{-1,\times}}(u) \\ &= \varepsilon_v(u) \chi(v) \mathbf{1}_{\mathfrak{D}_v^{-1}}(t) \mathbf{1}_{d_F^{-1,\times}}(\pi_\delta^{-1} u) \end{aligned}$$

Here recall that \mathfrak{D} is the relative different of E/F ; and that w acts as Fourier transform in t with respect to the quadratic form associated to $u\mathfrak{N}$, with the normalising constant $\gamma(u) = \gamma(E_v, u\mathfrak{N})$ as described in (2.1.1).

The computation of the expansion can then be performed exactly as in §2.4. We omit the details but indicate that the relevant substitution is now $a \rightarrow \pi_\mathfrak{d} a y$, where \mathfrak{d} is an ideal of \mathcal{O}_E of norm δ and $\pi_\mathfrak{d} \in \widehat{\mathcal{O}}_E$ is a generator with components equal to 1 away from \mathfrak{d} .

⁽¹⁷⁾Recall that our $\chi(v)$ are the inverses of the $\chi(v)$ of *loc. cit.*

Proposition 2.5.1. — *The series $\Theta(\mathcal{W})^{(\delta)}$ belongs to $S_1(K_1(\Delta(\mathcal{W}), \delta), \varepsilon \mathcal{W}|_{F_A^\times})$. Its Whittaker-Fourier coefficients are given by*

$$\tilde{a}(\Theta(\mathcal{W})^{(\delta)}, y) = \varepsilon \mathcal{W}(y) |y|^{1/2} \chi(\delta) \mathcal{W}(\mathfrak{d}) \varepsilon_\delta(y) r_{\mathcal{W}}(y d_F),$$

where $\chi(\delta) = \prod_{v|\delta} \chi(v)$.

3. Eisenstein measure

In this section we construct a measure (cf. §2.3) valued in Eisenstein series of weight one, and compute its Fourier expansion.

3.1. Eisenstein series. — Let k be a positive integer, M an ideal of \mathcal{O}_F , and $\varphi: F_A^\times/F^\times \rightarrow \mathbf{C}^\times$ a finite order character of conductor dividing M satisfying $\varphi_v(-1) = (-1)^k$ for $v|\infty$. Let

$$(3.1.1) \quad L^M(s, \varphi) = \sum_{(m, M)=1} \varphi(m) \mathbf{N}(m)^{-s}$$

where the sum runs over all nonzero ideals of \mathcal{O}_F .

Let $B \subset \mathbf{GL}_2$ be the Borel subgroup of upper triangular matrices; recall the notation from §1.1, and the Iwasawa decomposition (1.1.1); the decomposition is not unique but the ideal of $\widehat{\mathcal{O}}_F$ generated by the lower left entry of the $K_0(1)$ -component is well-defined.

For $s \in \mathbf{C}$, define a function $H_{k,s}(g, \varphi)$ on $\mathbf{GL}_2(\mathbf{A}_F)$ by

$$H_{k,s}(g = qur(\theta); \varphi) = \begin{cases} \left| \frac{y_1}{y_2} \right|^s \varphi(y_1 a) \mathbf{e}_\infty(k\theta) & \text{if } u = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K_0(M) \\ 0 & \text{if } u \in K_0(1) \setminus K_0(M_0). \end{cases}$$

where we have written $g = qur(\theta)$ with $q = \begin{pmatrix} y_1 & x \\ & y_2 \end{pmatrix} \in B(\mathbf{A}_F)$, $u \in K_0(1)$, $r(\theta) \in K_\infty$.

We define two **Eisenstein series**

$$E_k^M(g, s; \varphi) = L^M(2s, \varphi) \sum_{\gamma \in B(F) \backslash \mathbf{GL}_2(F)} H_{k,s}(\gamma g; \varphi),$$

$$\tilde{E}_k^M(g, s; \varphi) = W_M E_k^M(g, s; \varphi) = \varphi^{-1}(\det g \pi_M) E_k^M(g W_M, s; \varphi)$$

which are absolutely convergent for $\Re s > 1$ and continue analytically for all s to (non-holomorphic) automorphic forms of level M , parallel weight k and character φ (for E) and φ^{-1} (for \tilde{E}). Here W_M is as in (1.4.3). The superscript M will be omitted from the notation when its value is clear from context.

3.2. Fourier expansion of the Eisenstein measure. — We specialise to the case where k is odd, $M = \Delta P$ with $(\Delta, P) = 1$, $\varphi = \varepsilon \phi$ with $\varepsilon = \varepsilon_{E/F}$ and ϕ a character of conductor dividing P , trivial at infinity (in particular we have $\varphi_v(-1) = \varepsilon_v(-1) \phi_v(-1) = -1$ as required). We assume that Δ is squarefree. For $\delta|\Delta$ we compute⁽¹⁸⁾ the Whittaker coefficients (c f. §1.2; we suppress φ , M and k

⁽¹⁸⁾Cf. [57, §§ 3.5, 6.2].

from the notation) of $\tilde{E}^{(\delta)}$;

$$c_s^\delta(\alpha, \gamma) = D_F^{-1/2} \int_{\mathbf{A}_F/F} \tilde{E} \left(\begin{pmatrix} \gamma & x \\ & 1 \end{pmatrix} W_{\delta, s} \right) \mathbf{e}(-\alpha x) dx$$

for $\alpha \in F$ and δ dividing Δ ; since $c_s(\alpha, \gamma) = c_s(1, \alpha\gamma)$ for $\alpha \neq 0$, we can restrict to $\alpha = 0$ or 1. The choice of uniformisers π_v at $v|\delta$ implicit in the above formula is made so that $\varepsilon(\pi_v) = 1$ to save some notation.

Proposition 3.2.1. — *In the case just described, the Whittaker coefficients $c_s^\delta(\alpha, \gamma)$ of the Eisenstein series $\tilde{E}_k^{(\delta)}(g, s; \varphi)$ are given by*

$$c_s^\delta(0, \gamma) = \begin{cases} \frac{1}{D_F^{1/2} \mathbf{N}(\Delta P)^s} \varepsilon \phi(\gamma) |\gamma|^{1-s} V_{k, s}(0)^{[F:\mathbf{Q}]} L^{(P)}(2s-1, \varepsilon \phi) & \text{if } \delta = 1 \\ 0 & \text{if } \delta \neq 1, \end{cases}$$

$$c_s^\delta(1, \gamma) = \frac{\mathbf{N}(\delta)^{s-1/2}}{D_F^{1/2} \mathbf{N}(\Delta P)^s} \varepsilon \phi(\gamma) |\gamma|^{1-s} \chi(\delta) \phi(\delta) \varepsilon_\delta(\gamma) \phi_\delta(\gamma^\infty d_F) |\gamma \pi_\delta d_F|_\delta^{2s-1} \sigma_{k, s, \varepsilon \phi}(\gamma)$$

if γd_F is integral, and $c_s^\delta(1, \gamma) = 0$ otherwise,

where $\chi(\delta) = \prod_{v|\delta} \chi(v)$ with $\chi(v)$ as in (2.1.1) and

$$\sigma_{k, s, \varphi}(\gamma) = \prod_{v \nmid \Delta M \infty} \sum_{n=0}^{v(\gamma d_F)} \varphi_v(\pi_v)^n |\pi_v|^{n(2s-1)} \prod_{v|\infty} V_{k, s}(\gamma_v)$$

with

$$V_{k, s}(\gamma) = \int_{\mathbf{R}} \frac{e^{-2\pi i \gamma x}}{(x^2 + 1)^{s-k/2} (x+i)^k} dx.$$

Proof. — We use the Bruhat decomposition

$$\mathbf{GL}_2(F) = B(F) \coprod B(F)wN(F)$$

with $w = \begin{pmatrix} & -1 \\ 1 & \end{pmatrix}$ and the unipotent subgroup $N(F) \cong F$ via $N(F) \ni \begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} \leftrightarrow x \in F$, to get

$$\begin{aligned} \varepsilon \phi(\gamma) \phi(\pi_{M/\delta}) c_s^\delta(\alpha, \gamma) &= L(2s, \varphi) D_F^{-1/2} \int_{\mathbf{A}_F/F} H_s \left(\begin{pmatrix} \gamma & x \\ & 1 \end{pmatrix} W_{M/\delta} \right) \mathbf{e}(-\alpha x) dx \\ &\quad + L(2s, \varphi) D_F^{-1/2} \int_{\mathbf{A}_F} H_s \left(w \begin{pmatrix} \gamma & x \\ & 1 \end{pmatrix} W_{M/\delta} \right) \mathbf{e}(-\alpha x) dx. \end{aligned}$$

At any place $v|M/\delta$, we have the decomposition

$$\begin{pmatrix} \gamma_v & x_v \\ & 1 \end{pmatrix} W_{M/\delta, v} = \begin{pmatrix} \gamma_v & \pi_v x_v \\ & \pi_v \end{pmatrix} \begin{pmatrix} & 1 \\ -1 & \end{pmatrix}$$

so that the first summand is always zero.

For the second integral, we use the identity

$$w \begin{pmatrix} y & x \\ & 1 \end{pmatrix} = \begin{pmatrix} 1 & \\ & y \end{pmatrix} \begin{pmatrix} & -1 \\ 1 & xy^{-1} \end{pmatrix}$$

and the substitution $x \rightarrow xy$ to get

$$\int_{\mathbb{A}_F} H_s \left(w \begin{pmatrix} y & x \\ & 1 \end{pmatrix} W_{M/\delta} \right) \mathbf{e}(-\alpha x) dx = |y|^{1-s} \prod_v V_s^{M/\delta}(\alpha_v y_v)$$

where for $y \in F_v$,

$$(3.2.1) \quad V_s^M(y) = \int_{F_v} H_s \left(\begin{pmatrix} & -1 \\ 1 & x \end{pmatrix} W_{M,v} \right) \mathbf{e}(-xy) dx.$$

Archimedean places. — As in [57, Proposition 3.5.2].

Nonarchimedean places $v \nmid M/\delta$. — If v is a finite place, we have $\begin{pmatrix} & -1 \\ 1 & x \end{pmatrix} \in \mathbf{GL}_2(\mathcal{O}_{F,v})$ if $x \in \mathcal{O}_{F,v}$, and otherwise we have the decomposition

$$\begin{pmatrix} & -1 \\ 1 & x \end{pmatrix} = \begin{pmatrix} x^{-1} & -1 \\ & x \end{pmatrix} \begin{pmatrix} 1 & \\ & x^{-1} \end{pmatrix}.$$

Therefore

$$(3.2.2) \quad H_{s,v} \left(\begin{pmatrix} & -1 \\ 1 & x \end{pmatrix} \right) = \begin{cases} \bar{\varphi}_v(x) |x|^{-2s} & \text{if } v(x) \leq -1; \\ 1 & \text{if } v \nmid M, v(x) \geq 0; \\ 0 & \text{if } v|\delta, v(x) \geq 0. \end{cases}$$

The case $v \nmid M$. — We deduce that

$$\begin{aligned} V_s^{M/\delta}(y) &= \int_{\mathcal{O}_{F,v}} \mathbf{e}(-xy) dx + \sum_{n \geq 1} \int_{\mathcal{O}_{F,v}^\times} \bar{\varphi}_v(x \pi_v^{-n}) |x \pi_v^{-n}|^{-2s} \mathbf{e}(-xy \pi_v^{-n}) d(\pi_v^{-n} x) \\ &= \mathbf{1}[y \in d_F^{-1}] + \sum_{n \geq 1} \varphi_v(\pi_v)^n |\pi_v|^{n(2s-1)} \int_{\mathcal{O}_{F,v}^\times} \mathbf{e}(-xy \pi_v^{-n}) dx. \end{aligned}$$

The integral evaluates to $1 - |\pi_v|$ if $v(y d_F) \geq n$, to $-|\pi_v|$ if $v(y d_F) = n - 1$, and to zero otherwise. Therefore we have $V_s^M(y) = 0$ unless $v(y d_F) \geq 0$ in which case if $y \neq 0$

$$\begin{aligned} V_s^{M/\delta}(y) &= 1 + (1 - |\pi_v|) \sum_{n=1}^{v(y d_F)} (\varphi_v(\pi_v) |\pi_v|^{2s-1})^n - |\pi_v| (\varphi_v(\pi_v) |\pi_v|^{2s-1})^{v(y d_F)+1} \\ &= (1 - \varphi_v(\pi_v) |\pi_v|^{2s}) \sum_{n=0}^{v(y d_F)} \varphi_v(\pi_v)^n |\pi_v|^{n(2s-1)} \\ &= L_v(2s, \varphi)^{-1} \sum_{n=0}^{v(y d_F)} \varphi_v(\pi_v)^n |\pi_v|^{n(2s-1)}; \end{aligned}$$

whereas for $y = 0$, we have

$$V_s^{M/\delta}(0) = 1 + (1 - |\pi_v|) \sum_{n=1}^{\infty} (\varphi_v(\pi_v) |\pi_v|^{2s-1})^n$$

$$\begin{aligned} &= 1 + (1 - |\pi_v|)(1 - \varphi_v(\pi_v)|\pi_v|^{2s-1})^{-1}(1 - \varphi_v(\pi_v)|\pi_v|^{2s}) \\ &= L_v(2s, \varphi)^{-1}L_v(2s-1, \varphi) \end{aligned}$$

The case $v|\delta$. — Again by (3.2.2) we find

$$V_s^{M/\delta}(y) = \sum_{n \geq 1} \int_{\mathcal{O}_{F,v}^\times} \overline{\varphi}_v(x\pi_v^{-n})|x\pi_v^{-n}|^{-(2s-1)} \mathbf{e}(-xy\pi_v^{-n}) dx.$$

All the integrals vanish except the one with $n = v(yd_F) + 1$ which gives

$$\varepsilon_v(y\pi_v^n)\phi_v(y\pi_{d_F,v}\pi_v)|y\pi_{d_F,v}\pi_v|^{2s-1}|\pi_v|^{1/2}\chi(v);$$

therefore we have⁽¹⁹⁾

$$V_s^{M/\delta}(y) = \varepsilon_v(y)\phi_v(y\pi_{d_F,v}\pi_v)|y\pi_{d_F,v}\pi_v|^{2s-1}|\pi_v|^{1/2}\chi(v)$$

if $y \neq 0$ and $v(yd_F) \geq 0$ and $V_s(y) = 0$ otherwise. In particular, we see that if $\delta \neq 1$ then $V_s(0) = c_s(0, y) = 0$.

Places $v|M/\delta$. — For $w \begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} \begin{pmatrix} & 1 \\ -\pi_v^{v(M)} & \end{pmatrix} = \begin{pmatrix} \pi_v^{v(M)} & \\ -x\pi_v^{v(M)} & 1 \end{pmatrix}$ we have the decompositions

$$\begin{aligned} \begin{pmatrix} \pi_v^{v(M)} & \\ -x\pi_v^{v(M)} & 1 \end{pmatrix} &= \begin{pmatrix} -\pi_v^{v(M)} & \\ & 1 \end{pmatrix} \begin{pmatrix} -1 & \\ -x\pi_v^{v(M)} & 1 \end{pmatrix} \\ &= \begin{pmatrix} x^{-1} & -\pi_v^{v(M)} \\ & x\pi_v^{v(M)} \end{pmatrix} \begin{pmatrix} & 1 \\ -1 & x^{-1}\pi_v^{-v(M)} \end{pmatrix} : \end{aligned}$$

for $v(x) \geq 0$ we use the first one to find

$$H_s \left(\begin{pmatrix} -\pi_v^{v(M)} & \\ x\pi_v^{v(M)} & -1 \end{pmatrix} \right) = \varphi_v(\pi_v)^{v(M)}|\pi_v^{v(M)}|^s;$$

for $v(x) < 0$ the second decomposition shows that the integrand vanishes. We conclude that

$$V_s^{M/\delta}(y) = \begin{cases} \varphi_v(\pi_v)^{v(M)}|\pi_v^{v(M)}|^s & \text{if } v(yd_F) \geq 0 \\ 0 & \text{otherwise.} \end{cases}$$

The final formula follows from these computations. □

We specialise to the case $s = 1/2$, and consider the rescaled holomorphic Eisenstein series:⁽²⁰⁾

$$(3.2.3) \quad \begin{aligned} \mathbf{E}_{k,\varepsilon\phi}^{\Delta P}(g) &= \frac{D_F^{1/2}\mathbf{N}(\Delta P)^{1/2}}{(-2\pi i)^{[F:\mathbb{Q}]}} E_k^{\Delta P}(g, 1/2; \varepsilon\phi), \\ \tilde{\mathbf{E}}_{k,\varepsilon\phi}^{\Delta P}(g) &= \frac{D_F^{1/2}\mathbf{N}(\Delta P)^{1/2}}{(-2\pi i)^{[F:\mathbb{Q}]}} \tilde{E}_k^{\Delta P}(g, 1/2; \varepsilon\phi). \end{aligned}$$

We further specialise to the case $k = 1$.

⁽¹⁹⁾Recall that we always choose π_v so that $\varepsilon_v(\pi_v) = 1$.

⁽²⁰⁾Notice that these series do not depend on the ideal P but only on its support.

Corollary 3.2.2. — *The Eisenstein series $\tilde{\mathbf{E}}_{1,\varepsilon\phi}^{\Delta P}$ belongs to $M_1(K_1(\Delta P), \varepsilon\phi^{-1})$. The Whittaker-Fourier coefficients of $\tilde{\mathbf{E}}_{\varepsilon\phi}^{(\delta)} = \tilde{\mathbf{E}}_{1,\varepsilon\phi}^{\Delta P,(\delta)}$ for $\delta|\Delta$ are given by*

$$\tilde{a}^0(\tilde{\mathbf{E}}_{\varepsilon\phi}^{(\delta)}, y) = \varepsilon\phi(y)|y|^{1/2} \frac{L^{(P)}(0, \varepsilon\phi)}{2^g}$$

if $\delta = 1$ and $\tilde{a}^0(\tilde{\mathbf{E}}_{\varepsilon\phi}^{(\delta)}, y) = 0$ otherwise; and

$$c^\delta(y) = \tilde{a}(\tilde{\mathbf{E}}_{\varepsilon\phi}^{(\delta)}, y) = \varepsilon\phi(y)|y|^{1/2} \chi(\delta)\phi(\delta)\varepsilon_\delta\phi_\delta(y^\infty d_F)\sigma_{\varepsilon\phi}(y^\infty d_F),$$

where for any integral ideal m of $\mathcal{O}_F[\Delta^{-1}P^{-1}]$,

$$\sigma_{\varepsilon\phi}(m) = \sum_{d|m} \varepsilon\phi(d),$$

the sum likewise running over integral ideals of $\mathcal{O}_F[\Delta^{-1}P^{-1}]$.

(If m is an integral ideal of \mathcal{O}_F prime to P , then $\sigma_{\varepsilon_1}(m) = r(m)$.)

Proof. — This follows from Proposition 3.2.1 together with the evaluation

$$V_{1,1/2}(t) = \begin{cases} 0 & \text{if } t < 0 \\ -\pi i & \text{if } t = 0 \\ -2\pi i e^{-2\pi t} & \text{if } t > 0. \end{cases}$$

which can be found in [20, Proposition IV.3.3 (a), (d)] (for the case $t = 0$, this is deduced from (a) of *loc. cit.* using $\lim_{s \rightarrow 0} \frac{\Gamma(2s)}{\Gamma(s)} = 1/2$). \square

Definition 3.2.3. — Let F'_∞ be the maximal abelian extension of F unramified outside p , and let $\mathcal{G}'_F = \text{Gal}(F'_\infty/F)$. We define the **Eisenstein (pseudo-) measure**⁽²¹⁾ $\tilde{\mathbf{E}}_\varepsilon$ on \mathcal{G}'_F by

$$\tilde{\mathbf{E}}_\varepsilon(\phi) = \tilde{\mathbf{E}}_{\varepsilon\phi}^{\Delta P} = \frac{D_F^{1/2} \mathbf{N}(\Delta P)^{1/2}}{(-2\pi i)^g} \tilde{E}_{\varepsilon\phi}^{\Delta P}$$

for any character ϕ of \mathcal{G}'_F of conductor dividing P (it does not depend on the choice of P once we require P to satisfy $v|P \leftrightarrow v|p$). We denote with the same name the distribution induced on the group \mathcal{G}' of §2.3 by

$$\tilde{\mathbf{E}}_\varepsilon(\mathcal{W}) = \tilde{\mathbf{E}}_\varepsilon(\mathcal{W}|_{F_\Lambda^\times}).$$

It has values in $\mathbf{M}_1(K_1(N\Delta), \varepsilon)$ and is defined over \mathbf{Q}_p .

To prove the soundness of the definition, it is easy to see that the nonzero Fourier coefficients interpolate to a measure on \mathcal{G}'_F , that is an element of $\mathbf{Z}_p[[\mathcal{G}'_F]]$. The L -values giving the constant term interpolate to the Deligne–Ribet p -adic L -function [16]; it is a pseudo-measure in the sense of Serre [46], that is an element of the total quotient ring of $\mathbf{Z}_p[[\mathcal{G}'_F]]$ with denominators of a particularly simple form.

⁽²¹⁾We do not need to assume that Δ is squarefree when making the definition. See after the definition for the meaning of the term pseudo-measure.

4. The p -adic L -function

4.1. Rankin–Selberg convolution. — Let f, g be modular forms of common level M , weights k_f, k_g , and characters ψ_f, ψ_g respectively. We define a normalised Dirichlet series

$$D^M(f, g, s) = L^M(2s - 1, \psi_f \psi_g) \sum_m a(f, m) a(g, m) \mathbf{N} m^{-s},$$

where the imprimitive L -function $L^M(s, \varphi)$ of a Hecke character φ of conductor dividing M is as in (3.1.1).

When f and g are primitive forms of level N_f, N_g (that is, normalised new eigenforms at those levels), for a prime $\varphi \nmid N_f$ denote by $\gamma_\varphi^{(1)}(f), \gamma_\varphi^{(2)}(f)$ the two roots of the φ^{th} Hecke polynomial of f

$$P_{\varphi, f}(X) = X^2 - a(f, \varphi)X + \psi_f(\varphi) \mathbf{N} \varphi^{k_f - 1},$$

and by $\gamma_\varphi^{(1)}(g), \gamma_\varphi^{(2)}(g)$ the analogous quantities for g . Then the degree four Rankin–Selberg L -function $L(f \times g, s)$ with unramified Euler factors at φ given by

$$\prod_{i, j=1}^2 \left(1 - \gamma_\varphi^{(i)}(f) \gamma_\varphi^{(j)}(g) \mathbf{N} \varphi^{-s} \right)^{-1}$$

equals the above Dirichlet series

$$L(f \times g, s) = D^{N_f N_g}(f, g, s)$$

if N_f and N_g are coprime.

Suppose now for simplicity that $k_f = 2, k_g = 1$, and f is a cusp form (not necessarily primitive). The Rankin–Selberg convolution method⁽²²⁾ gives

$$(4.1.1) \quad \langle f^\rho, g E_1^M(s; \psi_f \psi_g) \rangle_M = D_F^{s+1} \left[\frac{\Gamma(s+1/2)}{(4\pi)^{s+1/2}} \right]^{[F:\mathbf{Q}]} D^M(f \times g, s+1/2),$$

where $\langle \cdot, \cdot \rangle_M$ is the Petersson inner product (1.1.2).

4.2. Convoluted measure and the p -adic L -function in the ordinary case. — Consider the convolution pseudo-measure $\Theta * \tilde{\mathbf{E}}_{\varepsilon, N}$ on \mathcal{G}' defined by $\Theta * \tilde{\mathbf{E}}_{\varepsilon, N}(\mathcal{W}) = \Theta(\mathcal{W}) \tilde{\mathbf{E}}_{\varepsilon, N}(\overline{\mathcal{W}})$ for any character $\mathcal{W} : \mathcal{G} \rightarrow \mathbf{Z}_p^\times$, where $\tilde{\mathbf{E}}_{\varepsilon, N} = [N] \tilde{\mathbf{E}}_\varepsilon$. We deduce from it the (pseudo-)measure

$$(4.2.1) \quad \Phi(\mathcal{W}) = \text{Tr}_\Delta[\Theta * \tilde{\mathbf{E}}_{\varepsilon, N}(\overline{\mathcal{W}})] = \text{Tr}_\Delta[\Theta(\mathcal{W}) \cdot [N] \tilde{\mathbf{E}}_\varepsilon(\overline{\mathcal{W}})]$$

on \mathcal{G}' , which is a kind of p -adic kernel of the Rankin–Selberg L -function as will be made precise below. It is valued in $\mathbf{S}_2(K_0(N), \mathbf{C}_p)$. Notice that while $\Phi(\mathcal{W})$, like $\tilde{\mathbf{E}}_{\varepsilon, N}$, is not a measure, we can see that, for any $\varphi | p$,

$$U_\varphi \Phi(\mathcal{W})$$

is. Indeed its Fourier coefficients are the Fourier coefficients of $\Phi(\mathcal{W})$ at ideals m divisible by φ , hence sums of coefficients of the theta and Eisenstein series at pairs of ideals ($m_1 = nm, m_2 = (1-n)m$) for some $n \in F$; since the coefficients of the theta series are zero at ideals m_1 divisible by

⁽²²⁾See [47] or [28, Ch. V] for general treatments; our setting and normalisations are the same as in [57, Lemma 6.1.3] (where g is a specific form, but the same calculation works in general to prove (4.1.1)).

\wp , only those pairs (m_1, m_2) with m_1, m_2 both prime to \wp contribute. In particular, the constant term of the Eisenstein series does not contribute to the Fourier expansion of $U_\wp \Phi(\mathcal{W})$, which therefore belongs to $\mathbf{Z}_p[[\mathcal{G}']]$.

Thanks to this discussion and the identity $l_{f_\alpha} = \alpha^{-1} l_{f_\alpha} \circ U_\wp$, the following definition makes sense.

Definition 4.2.1. — The p -adic Rankin–Selberg L -function is the element of $\mathcal{O}_L[[\mathcal{G}]] \otimes L$ defined by

$$L_{p,\alpha}(f_E, \mathcal{W}) = D_F^{-2} H_p(f) l_{f_\alpha}(\Phi(\mathcal{W}))$$

for any character $\mathcal{W} : \mathcal{G} \rightarrow \mathcal{O}_L^\times$, where

$$(4.2.2) \quad H_p(f) = \prod_{\wp|p} \left(1 - \frac{1}{\alpha_\wp(f)^2} \right) \left(1 - \frac{\mathbf{N}_\wp}{\alpha_\wp(f)^2} \right).$$

Functional equation. — The p -adic L -function admits a functional equation; we prove it in the case of anticyclotomic characters which is the only one we shall need.

Proposition 4.2.2. — Suppose that \mathcal{W} is an anticyclotomic character of \mathcal{G} , i.e., $\mathcal{W}|_{F_{A^\times}} = 1$. Then there are functional equations for the p -adic L -function

$$(4.2.3) \quad L_{p,\alpha}(f_E)(\mathcal{W}) = (-1)^g \varepsilon(N) L_{p,\alpha}(\mathcal{W})$$

and for the analytic kernel

$$(4.2.4) \quad \Phi(\mathcal{W}) = (-1)^g \varepsilon(N) \Phi(\mathcal{W}).$$

In particular, if $\varepsilon(N) = (-1)^{g-1}$, we have

$$\Phi(\mathcal{W}) = L_{p,\alpha}(f_E)(\mathcal{W}) = 0.$$

Proof. — The functional equation for $L_{p,\alpha}$ is implied by the functional equation for Φ . We prove the latter by comparing the coefficients on both sides. From (4.5.1) below,⁽²³⁾ the coefficients of $\Phi(\mathcal{W})$ are given by

$$b(m) = \sum_{\delta|\Delta} \sum_{\substack{n \in F \\ 0 < n < 1}} \varepsilon_\delta((n-1)n) r_{\mathcal{W}^-}((1-n)m\delta) \sigma_{\varepsilon_1}(nm/N).$$

⁽²³⁾Which does not use the present result. The formula (4.5.1) is stated in the case when the anticyclotomic part $\mathcal{W}^- = 1$ but the very same calculation gives the result in general.

(We use the notation $\mathbf{1}$ for the character of ideals defined by $\mathbf{1}(m) = 1$ if $(m, p) = 1$ and $\mathbf{1}(m) = 0$ otherwise.) We rewrite this as $b(m) = \sum_{\delta, n} b_{\delta, n}(m)$ with, using $\varepsilon_\delta(x) = \varepsilon^\delta(x)$ for $x \in F^\times$:

$$\begin{aligned}
 b_{\delta, n}(m) &= \varepsilon_\delta(-1) \\
 &\quad \cdot \varepsilon_{\Delta/\delta}((1-n)m) \varepsilon_{\Delta/\delta}(nm) \\
 &\quad \cdot \varepsilon^\Delta((1-n)m) r_{\mathcal{W}}((1-n)m\delta) \\
 &\quad \cdot \varepsilon(N) \\
 &\quad \cdot \varepsilon^\Delta(nm/N) \sigma_{\varepsilon_1}(nm/N) \\
 &= (-1)^g \varepsilon_{\Delta/\delta}(-1) \\
 &\quad \cdot \varepsilon_{\Delta/\delta}((1-n)m) \varepsilon_{\Delta/\delta}(nm) \\
 &\quad \cdot r_{\mathcal{W}^-}((1-n)m\Delta/\delta) \\
 &\quad \cdot \varepsilon(N) \\
 &\quad \cdot \sigma_{\varepsilon_1}(nm/N) \\
 &= (-1)^g \varepsilon(N) b_{\Delta/\delta, n}.
 \end{aligned}$$

Here we have used the following facts. In the first line, $\varepsilon_\Delta(-1) = \varepsilon_\infty(-1) = (-1)^g$. In the third line, we have that $r_{\mathcal{W}}(m) = 1$ if m is divisible only by ramified primes in E , since in that case $m = m^2$ is a square and $\mathcal{W}(m)^2 = \mathcal{W}(m) = 1$ - this implies $\mathcal{W}(m) = \pm 1$, hence $\mathcal{W}(m) = 1$ since \mathcal{W} , which is a character of $\mathcal{G} \cong \mathbf{Z}_p^{1+g+\delta}$, has values in $1 + p\mathbf{Z}_p$. Finally, in the third and fifth line one can observe that if $q = \sigma_{\varepsilon_1}$ or $q = r$, then $\varepsilon^\Delta(m)q(m) = q(m)$; indeed this is trivial if $\varepsilon^\Delta(m) = 1$, while both sides are zero if $\varepsilon^\Delta(m) = -1$. \square

4.3. Interpolation property. — We manipulate the definition to show that the p -adic L -function $L_{p, \alpha}(f_E)(\mathcal{W})$ of Definition 4.2.1 interpolates the special values of the complex Rankin-Selberg L -function $L(f_E, \mathcal{W}, s) = L(f \times \Theta(\mathcal{W}), s)$ defined in the Introduction.

We will need a few technical lemmas.

Lemma 4.3.1. — *Let P be an ideal of \mathcal{O}_F such that $v|P$ if and only if $v|p$. We have*

$$\langle W_{NP} f_\alpha^\rho, f_\alpha \rangle_{NP} = \alpha_P(f) (-1)^g \tau(f) H_p(f) \langle f, f \rangle_N$$

with $H_p(f)$ as in (4.2.2) and

$$\alpha_P(f) = \prod_{\wp|P} \alpha_\wp(f)^{v_\wp(P)}$$

Proof. — When $P = P_0 := \prod_{\wp|p} \wp$, this is the direct generalisation of [40, Lemme 27], and it is proved in the same way. In general, we can write $P = P_0 P_1$, and then

$$W_{NP} f_\alpha^\rho = \mathbf{N}(P_1)[P_1] W_{NP_0} f_\alpha^\rho$$

Observing that $[\wp]$ is the adjoint of U_\wp for the Petersson inner product and that $\mathbf{N}(P_1)[P_1] = [K_0(NP) : K_0(NP_0)]$, we deduce

$$\langle W_{NP} f_\alpha^\rho \rangle_{NP} = [K_0(NP) : K_0(NP_0)] \langle W_{NP_0} f_\alpha^\rho, U(P_1) f_\alpha \rangle_{NP} = \alpha_{P_1} \langle W_{NP_0} f_\alpha^\rho, f_\alpha \rangle_{NP_0}.$$

The lemma then follows from this and the special case $P = P_0$. \square

Lemma 4.3.2. — For a character φ of conductor dividing M and an ideal N prime to M , let $E_\varphi^M = E_1^M(g, 1/2; \varphi)$, $\tilde{E}_\varphi^M = W_M E_\varphi^M$. We have

$$W_M[N]\tilde{E}_\varphi^M = E_\varphi^{MN} + E^{\text{old}}$$

where the form E^{old} is old at N (in particular, E^{old} is orthogonal to newforms of level N and so is its product with any other N -old form).

Proof. — It is easy to see that $W_M[N]\tilde{E}_\varphi^M = [N]E_\varphi^M$. Then we are reduced to showing that

$$[N]E_\varphi^M = E_\varphi^{MN} + E^{\text{old}}.$$

We show more generally and more precisely that

$$\mathbf{N}(N)^{s-} E_\varphi^M(g \begin{pmatrix} 1 & \\ & \pi_M \end{pmatrix}, s) = \sum_{d|N} \frac{\varphi(d)}{\mathbf{N}(d)^{2s}} E_\varphi^{MN/d}(g, s)$$

which is [57, Lemma 6.1.4] with ε replaced by φ . The lemma then holds with

$$E^{\text{old}} = \sum_{d|N, d \neq 1} \frac{\varphi(d)}{\mathbf{N}(d)} E_\varphi^{MN/d}.$$

□

Lemma 4.3.3. — With notation as in §4.1, we have

$$D([\Delta]f, \Theta(\mathcal{W}), 1) = \mathcal{W}(\mathfrak{D})D(f, \Theta(\mathcal{W}), 1).$$

The proof is an easy calculation, see [37, §I.5.9].

Theorem 4.3.4. — Let $\mathcal{W}: \mathcal{G}' \rightarrow \overline{\mathbf{Q}}^\times$ be a finite order character of conductor \mathfrak{f} divisible only by primes above p . Then we have

$$L_{p,\alpha}(f_E)(\mathcal{W}) = \frac{\tau(\overline{\mathcal{W}})\mathbf{N}(\Delta(\mathcal{W}))^{1/2}V_p(f, \mathcal{W})\overline{\mathcal{W}}(\Delta)}{\alpha_{\mathfrak{N}(\mathfrak{f}(\mathcal{W}))}(f)\Omega_f} L(f_E, \overline{\mathcal{W}}, 1),$$

where $\Omega_f = (8\pi^2)^g \langle f, f \rangle_N$, $\tau(\overline{\mathcal{W}})$ is as in Lemma 2.4.3, and

$$(4.3.1) \quad V_p(f, \overline{\mathcal{W}}) = \prod_{\mathfrak{p}|p} \prod_{\mathfrak{p}|\mathfrak{p}} \left(1 - \frac{\overline{\mathcal{W}}(\mathfrak{p})}{\alpha_\varphi(f)} \right).$$

Proof. — Denote $P = \mathfrak{N}(\mathfrak{f}(\mathcal{W}))$, $\Delta(\mathcal{W}) = \Delta P$, $\phi = \mathcal{W}|_{F_A^\times}$. We suppose that \mathcal{W} is ramified at all places $v|p$ (in this case, we have $V_p(f, \mathcal{W}) = 1$). Then the result follows from the definition and

the following calculation.

$$\begin{aligned}
 L_f(\Phi(\mathcal{W})) &= \frac{\langle W_{NP} f_\alpha^\rho, \text{Tr}_\Delta[\Theta(\mathcal{W}) \tilde{\mathbf{E}}_{\varepsilon, N}(\overline{\mathcal{W}})] \rangle_{NP}}{\langle W_{NP} f_\alpha^\rho, f_\alpha \rangle_{NP}} \\
 \text{(L. 4.3.1)} &= \frac{\langle W_{N\Delta} f_\alpha^\rho, W_{\Delta(\mathcal{W})} \Theta(\mathcal{W}) W_{\Delta(\mathcal{W})} \tilde{\mathbf{E}}_{\varepsilon \phi^{-1}, N}^{\Delta(\mathcal{W})} \rangle_{N\Delta(\mathcal{W})}}{\alpha_p(f) (-1)^g \tau(f) H_p(f) \Omega_f} \\
 \text{(L. 4.3.2, L. 2.4.3)} &= \frac{(-i)^g \tau(\overline{\mathcal{W}}) \overline{\mathcal{W}}(\mathfrak{D}) D_E}{\alpha_p(f) (-1)^g \tau(f) H_p(f) \Omega_f} \langle W_N[\Delta] f_\alpha^\rho, \Theta(\overline{\mathcal{W}}) \mathbf{E}_{\varepsilon \phi^{-1}}^{N\Delta(\mathcal{W})} \rangle_{N\Delta(\mathcal{W})} \\
 &= \frac{(-i)^g \tau(\overline{\mathcal{W}})}{\alpha_p(f) H_p(f) \Omega_f} \langle [\Delta] f_\alpha^\rho, \Theta(\overline{\mathcal{W}}) \mathbf{E}_{\varepsilon \phi^{-1}}^{N\Delta(\mathcal{W})} \rangle_{N\Delta(\mathcal{W})} \\
 \text{(eq. (4.1.1))} &= \frac{\tau(\overline{\mathcal{W}}) D_F^2 \mathbf{N}(\Delta(\mathcal{W}))^{1/2}}{\alpha_p(f) H_p(f) \Omega_f} D^{N\Delta(\mathcal{W})}([\Delta] f_\alpha, \Theta(\overline{\mathcal{W}}), 1) \\
 \text{(L. 4.3.3)} &= \frac{\tau(\overline{\mathcal{W}}) D_F^2 \mathbf{N}(\Delta(\mathcal{W}))^{1/2} \overline{\mathcal{W}}(\Delta)}{\alpha_p(f) H_p(f) \Omega_f} L(f_E, \overline{\mathcal{W}}, 1)
 \end{aligned}$$

where we have used various results from §1.4, and the fact that in our case $f^\rho = f$ as f has trivial character.

The previous calculation goes through in general with $\Delta(\mathcal{W})$ replaced by $\Delta(\mathcal{W})' = \text{l.c.m.}(\Delta(\mathcal{W}), \prod_{\varphi|p} \delta^\rho)$; then one further needs to compare the imprimitive Dirichlet series $D^{\Delta(\mathcal{W})'}(f_\alpha, \Theta(\overline{\mathcal{W}}), 1)$ with the L -value $L(f_E, \overline{\mathcal{W}}, 1)$. This is done in the same way as in the case of elliptic modular forms, see [40, Lemme 2.3 (i) and §4.4 (III)].⁽²⁴⁾ We omit the details since no new phenomena appear in our context and strictly speaking we do not need to use the precise form of the interpolation result except in the ramified case, which already determines $L_{p,\alpha}(f_E)$ uniquely. \square

4.4. Factorisation. — The p -adic analogue of the standard L -function of f has been studied by several authors (Manin, Dabrowski, Dimitrov, ...). Let $\mathcal{G}_F = \text{Gal}(F_\infty/F)$ where F_∞ is the maximal abelian extension of F unramified outside p .

Theorem 4.4.1. — *There is a p -adic L -function $L_{p,\alpha}(f) \in \mathcal{O}_L[[\mathcal{G}_F]] \otimes_{\mathcal{O}_L} L$ uniquely determined by the following property: for each finite order character $\chi: \mathcal{G}_F \rightarrow \overline{\mathbf{Q}}^\times$ of conductor $\mathfrak{f}(\chi)$ divisible by all the primes $\varphi|p$, we have*

$$L_{p,\alpha}(f, \chi) = \frac{\tau(\overline{\chi}) \mathbf{N}(\mathfrak{f}(\chi))^{1/2}}{\alpha_{\mathfrak{f}(\chi)}} \frac{L(f, \overline{\chi}, 1)}{\Omega_f^+}$$

⁽²⁴⁾Notice that, as in *op. cit.*, our $\Theta(\mathcal{W})$ is not the primitive theta series when \mathcal{W} is unramified at some $\varphi|p$; in general we have

$$\Theta(\mathcal{W}) = \prod_{\mathfrak{p}|\varphi|p} (1 - \mathbf{N}(\mathfrak{p})^{1/2} \mathcal{W}(\mathfrak{p})[\varphi]) \Theta(\mathcal{W})^{\text{prim}}$$

if $\Theta(\mathcal{W})^{\text{prim}}$ is the primitive theta series (i.e. the normalised newform in its representation). This replaces the second-last equation on [40, p. 21], whose $\Theta(\mathcal{W})$ (respectively $\Theta(\mathcal{W}'')$) is our $\Theta(\mathcal{W})^{\text{prim}}$ (respectively our $\Theta(\mathcal{W})$). The factor $V_p(f, \overline{\mathcal{W}})$ comes from the analogue of [40, Lemme 23].

where Ω_f is a normalised period and $\tau(\overline{\chi}) = \prod_{v|p} \tau(\overline{\chi}_v)$ with

$$\tau(\overline{\chi}_v) = |\pi_v|^{-c/2} \int_{F_v^\times} \overline{\mathcal{W}}_v(x_v) \mathbf{e}_v(-x_v) dx_v$$

if $c = v(\mathfrak{f}(\chi))$.

See [15] for the proof. Then we have

$$(4.4.1) \quad L_{p,\alpha}(f_E, \chi \circ \mathfrak{N}) = \frac{\Omega_f^+ \Omega_{f_\varepsilon}^+}{D_E^{-1/2} \Omega_f} L_{p,\alpha}(f, \chi) L_{p,\alpha}(f_\varepsilon, \chi),$$

where f_ε is the form with coefficients $a(f_\varepsilon, m) = \varepsilon(m)a(f, m)$ and $D_E = \mathbf{N}(\Delta)$.

4.5. Fourier expansion of the analytic kernel. — Consider the restriction of Φ to \mathcal{G} , the Galois group of the maximal \mathbf{Z}_p -extension of E unramified outside p . Any character \mathcal{W} of \mathcal{G} decomposes uniquely as $\mathcal{W} = \mathcal{W}^+ \mathcal{W}^-$ with $(\mathcal{W}^+)^c = \mathcal{W}$, $(\mathcal{W}^-)^c = \mathcal{W}^{-1}$ (we say that \mathcal{W}^+ is cyclotomic and \mathcal{W}^- is anticyclotomic or dihedral). Since we are interested in the case $\varepsilon(N) = (-1)^{s-1}$ in which Φ is zero on the anticyclotomic characters, we study the restriction of Φ to the cyclotomic characters. We can write $\mathcal{W}^+ = \chi \circ \mathfrak{N}$ for a Hecke character $\chi : F^\times \backslash F_{\mathbf{A}}^\times \rightarrow 1 + p\mathbf{Z}_p$, and we denote

$$\Theta_\chi = \Theta(\chi \circ \mathfrak{N}), \quad \Phi_\chi = \Phi(\chi \circ \mathfrak{N}).$$

From now on we assume that $(\Delta, 2) = 1$ and all primes $\wp|p$ are split in E .

Proposition 4.5.1. — *The Fourier coefficients $b(m) = a_p(\Phi_\chi, m)$ of the p -adic modular form Φ_χ are given by*

$$b(m) = \sum_{\substack{n \in F \\ 0 < n < 1 \\ n \in N m^{-1} \Delta^{-1}}} \chi((1-n)m) \prod_{v|\Delta} \left[1[v(nm) = 0] + \varepsilon_v((n-1)n) \chi_v^{-2}(nm \wp_v / N) \right] \\ \cdot r((1-n)m\Delta) \sigma_{\varepsilon \chi^{-2}}(nm/N).$$

Proof. — By (1.4.6), the Fourier coefficient $b(m)$ of $\Phi_\chi = \text{Tr}_\Delta[\Theta_\chi \tilde{\mathbf{E}}_{\varepsilon \chi^2, N}]$ is given by

$$b(m) = \sum_{\delta|\Delta} b^\delta(m\delta)$$

with

$$b^\delta(m) = a(\Phi_\chi^{(\delta)}, m) = |y|^{-1} \tilde{a}(\Phi_\chi^{(\delta)}, y) = |y|^{-1} \sum_{n \in F} \tilde{a}(\Theta_\chi^{(\delta)}, (1-n)y) \tilde{a}(\tilde{\mathbf{E}}_{\varepsilon \chi^{-2}, N}^{(\delta)}, ny) \\ = |y|^{-1} \sum_{n \in F} \tilde{a}(\Theta_\chi^{(\delta)}, (1-n)y) \tilde{a}(\tilde{\mathbf{E}}_{\varepsilon \chi^{-2}, N}^{(\delta)}, ny/\pi_N)$$

if $y \in F_{\mathbf{A}}^\times$ satisfies $y_\infty > 0$ and $y^\infty d_F = m$.

Then by Proposition 2.5.1 and Corollary 3.2.2, we have:⁽²⁵⁾

$$(4.5.1) \quad b(m) = \sum_{\delta|\Delta} \sum_{\substack{n \in F \\ 0 < n < 1}} \varepsilon_\delta((n-1)n) \chi^{-1}(\delta) \chi((1-n)m\delta) \chi_\delta^{-2}(nm\delta/N) \\ \cdot r((1-n)m\delta) \sigma_{\varepsilon_\delta \chi^{-2}}(nm/N).$$

We interchange the two sums and notice that the term corresponding to δ and n is nonzero only if $n \in Nm^{-1}\Delta^{-1}$ and $\delta_0|\delta$, where

$$\delta_0 = \delta_0(n) = \prod_{\substack{v|\Delta \\ v(nm)=-1}} \wp_v$$

(\wp_v being the prime corresponding to v). Now for each n we can rewrite the sum over δ (omitting the factor $\chi((1-n)m)$ and those on the second line of (4.5.1), which do not actually depend on δ) as

$$\varepsilon_{\delta_0}((n-1)n) \chi_{\delta_0}^{-2}(nm\delta_0/N) \sum_{\delta'|\Delta/\delta_0} \varepsilon_{\delta'}((n-1)n) \chi_{\delta'}^{-2}(nm\delta'/N) \\ = \prod_{v|\delta_0} \varepsilon_{\delta'}((n-1)n) \chi_v^{-2}(nm\wp_v) \prod_{v|\Delta/\delta_0} [1 + \varepsilon_v((n-1)n) \chi_v^{-2}(nm\wp_v)].$$

The asserted formula follows. \square

Remark 4.5.2. — If $v(nm) = -1$ then $(n-1)\pi_m\pi_v \equiv n\pi_m\pi_v$ in $(\mathcal{O}_{F,v}/\pi_v\mathcal{O}_{F,v})^\times$, so that we actually have

$$\varepsilon_v((n-1)n) = \varepsilon_v((n-1)\pi_m\pi_v) \varepsilon_v(n\pi_m\pi_v) = 1.$$

We can now compute the Fourier coefficients of the analytic kernel giving the central derivative of the p -adic L -function in the cyclotomic direction. To this end, let

$$\nu: \text{Gal}(\overline{\mathbf{Q}}/F) \rightarrow 1 + p\mathbf{Z}_p \subset \mathbf{Q}_p^\times$$

be a character of F which is ramified exactly at the primes dividing p , and for $s \in \mathbf{Z}_p$ denote⁽²⁶⁾ $\Phi(s) = \Phi_{\nu^s}$. By Remark ??, $\Phi(s)$ is “almost” overconvergent in the sense that $\prod_{\wp} (\alpha_\wp^{-1} U_\wp) \Phi(s)$ is, and we have $L_{p,\alpha}(f_E, \nu^s \circ \mathfrak{N}) = l_{f_\alpha}(\Phi(s))$. By Lemma 1.6.1.??, l_{f_α} is a continuous operator on overconvergent forms, hence

$$\frac{d}{ds} L_{p,\alpha}(f_E, \nu^s \circ \mathfrak{N}) = \frac{d}{ds} l_{f_\alpha}(\Phi(s)) = l_{f_\alpha}\left(\frac{d}{ds} \Phi(s)\right).$$

In particular $L'_{p,\alpha;\nu \circ \mathfrak{N}}(f_E, \mathbf{1}) = l_{f_\alpha}(\Phi'(0))$.

Let $\ell_F = \frac{d}{ds}|_{s=0} \nu^s: F^\times \setminus F_{\mathbf{A}^\infty}^\times \rightarrow \mathbf{Q}_p$ be the p -adic logarithm associated to ν .

Proposition 4.5.3. — Assume that $\varepsilon(N) = (-1)^{g-1}$. Then $\Phi(0) = 0$ and the Fourier coefficients $b'(m)$ of

$$\Phi'_\nu = \Phi'(0) = \frac{d}{ds} \Phi_{\nu^s}|_{s=0}$$

⁽²⁵⁾Recall that $\chi(v)^2 = \varepsilon_v(-1)$.

⁽²⁶⁾No confusion should arise from our recycling the letter s for this p -adic variable, the complex variable having now exited the scene.

are nonzero only for m integral and nonzero, in which case

$$b'(m) = \sum_v b'_v(m)$$

with the sum running over all finite places v of F and $b'_v(m)$ given for $(\prod_{\wp|p} \wp) | m$ by:

1. If $v = \wp$ is inert in E , then

$$b'_v(m) = \sum_{\substack{n \in N m^{-1} \Delta^{-1} \\ (p, nm) = 1 \\ \varepsilon_v((n-1)n) = 1 \forall v | \Delta \\ 0 < n < 1}} 2^{\omega_\Delta(n)} r((1-n)m\Delta) r(nm\Delta/N_\wp) (v(nm/N) + 1) \ell_{F,v}(\pi_v),$$

where

$$\omega_\Delta(n) = \#\{v | (\Delta, nm\Delta)\}.$$

2. If $v = \wp | \Delta$ is ramified in E , then

$$b'_v(m) = \sum_{\substack{n \in N m^{-1} \Delta^{-1} \\ (p, nm) = 1 \\ \varepsilon_v((n-1)n) = -1 \\ \varepsilon_w((n-1)n) = 1 \forall w \neq v | \Delta \\ 0 < n < 1}} 2^{\omega_\Delta(n)} r((1-n)m\Delta) r(nm\Delta/N) (v(nm) + 1) \ell_{F,v}(\pi_v).$$

3. If v is split in E , then

$$b'_v(m) = 0.$$

Proof. — The vanishing of $\Phi(0) = \Phi_1$ follows from the functional equation (4.2.4) and the sign assumption.

By Proposition 4.5.1, the Fourier coefficient $b_s(m)$ of $\Phi(s) = \Phi_v$ can be expressed as $b_s(m) = \sum_{n \in F} b_{n,s}(m)$ with

$$b_{n,s}(m) = v^s((1-n)m) r((1-n)m\Delta) \prod_{\wp \nmid p\infty} \sigma_{s,v}^n(m/N)$$

where, using Remark 4.5.2:

$$\sigma_{s,v}^n(m) = \begin{cases} \frac{1 - \varepsilon(nm\wp)v(nm\wp)^{-2s}}{1 - \varepsilon(\wp)v(\wp)^{-2s}} & \text{if } v = \wp \nmid \Delta; \\ 1 + \varepsilon_v(n(n-1))v(nm\wp)^{-2s} & \text{if } v = \wp | \Delta \text{ and } v(nm) = 0; \\ v(nm\wp)^{-2s} & \text{if } v = \wp | \Delta \text{ and } v(nm) = -1. \end{cases}$$

Then $b'(m) = \sum_n b'_n(m) = \sum_n \sum_v b'_{n,v}(m)$ with $\sum_n b'_{n,v}(m) = b'_v(m)$, and $b'_n(m)$ can be nonzero only if exactly one of the factors $\sigma_{s,v}^n$ vanishes at $s = 0$. If this happens for the place v_0 , then the set over which n ranges accounts for the positivity and integrality conditions and the non-vanishing conditions at other places, whereas the condition $(p, nm) = 1$ results from observing that $\lim_{s \rightarrow 0} v^s(a) = 1[(p, a) = 1]$.

The values of $b'_{n,v}$ can then be determined in each case from the above expressions: for v ramified this is straightforward. For $v = \wp$ inert, notice that if $v(nm/N)$ is odd then $r(nm\Delta/N_\wp) = r((nm\Delta/N)^{(\wp)})$, where the superscript denotes prime-to- \wp part; whereas if $v(nm/N)$ is even then $\sigma_{0,v}^n(m/N)$ does not vanish so (n, v) does not contribute to $b'(m)$ and indeed $r(nm/N_\wp) = 0$. \square

PART II HEIGHTS

5. p -adic heights and Arakelov theory

By the work of many authors (Schneider, Perrin-Riou, Mazur–Tate, Coleman–Gross, Zarhin, Nekovář, ...) there are p -adic height pairings on the Mordell–Weil group of an abelian variety defined over a number field. In this section, we first recall (§§5.1–5.2) a definition of the height pairing as a sum of local symbols following Zarhin [56] and Nekovář [36], and explain how it induces a pairing on degree zero divisors on curves. Then we describe a useful construction of the local symbols at places dividing p using norm systems (§??). In §§5.3–5.4 we explain how p -adic Arakelov theory allows to extend the height pairing for curves to a pairing on divisors of any degree.

5.1. Local symbols. — Let A be an abelian variety of dimension g over a local field E_v , A^\vee its dual abelian variety, and let $V = V_p A = T_p A \otimes_{\mathbf{Z}_p} \mathbf{Q}_p$ be the rational Tate module of A , a continuous $\text{Gal}(\bar{E}/E)$ -representation.⁽²⁷⁾ Let $\ell_v : E_v^\times \rightarrow \mathbf{Q}_p$ be a homomorphism; we call ℓ a *local p -adic logarithm* and assume that it is ramified, that is, $\ell_v : E_v^\times \rightarrow \mathbf{Q}_p$ does not vanish identically on $\mathcal{O}_{E,v}^\times$. Let $D_{\text{dR}}(V_v)$ be the filtered \mathbf{Q}_p -vector spaces associated to V by the theory of Fontaine. The comparison theorem identifies $D_{\text{dR}}(V_v)$ with $H_{\text{dR}}^1(A^\vee/E_v)$, equipped with the Hodge filtration; it is also identified with the filtered Dieudonné module of the special fibre of the p -divisible group of A (after an extension of scalars if E_v is ramified over \mathbf{Q}_p ; see [18]). Let L be a finite extension of the coefficient field \mathbf{Q}_p , and if $v|p$, let $W_v \subset D_{\text{dR}}(V) \otimes L$ be a splitting of the Hodge filtration, that is, a complementary subspace to $\Omega^1(A^\vee \otimes E_v/E_v) \otimes L \subset D_{\text{dR}}(V) \otimes L$, which is isotropic⁽²⁸⁾ for the cup product. When V is ordinary, there is a canonical choice for W_v , the “unit root” subspace (see e.g. [27] for a nice discussion); in general, the unit root subspace can be of any dimension between 0 and g , and we always require W_v to contain it.

We proceed to define pairings, called *local Néron symbols*⁽²⁹⁾

$$\langle \cdot, \cdot \rangle_{v,W} : (\mathcal{D}_0(A)(E_v) \times Z_0(A)^0(E_v))_e \rightarrow L$$

on the subset of pairs with disjoint supports in the product of the group $\mathcal{D}_0(A)(E_v)$ of divisors algebraically equivalent to zero defined over E_v and the group $Z_0(A)^0(E_v)$ of zero-cycles of degree zero defined over E_v .

Let $\mathcal{A}/\mathcal{O}_{E,v}$, $\mathcal{A}^\vee/\mathcal{O}_{E,v}$ be the Néron models of A and A^\vee , and let \mathcal{A}^0 be the identity component of \mathcal{A} . The rational equivalence class $[D]$ of $D \in \mathcal{D}_0(A)(E_v)$ defines a point in $A^\vee(E_v) = \mathcal{A}^\vee(\mathcal{O}_{E,v}) = \text{Ext}_{\text{fppf}}^1(\mathcal{A}^0, \mathbf{G}_m)$, hence an extension

$$1 \rightarrow \mathbf{G}_m \rightarrow \mathcal{Y}_{[D]} \rightarrow \mathcal{A}^0 \rightarrow 1$$

⁽²⁷⁾Nekovář [36] defines height pairings for Galois representations in much greater generality than described here.

⁽²⁸⁾The isotropy condition ensures that the resulting height pairing is symmetric [36, Theorem 4.1.1 (4)]

⁽²⁹⁾The notation is a bit abusive: the subscript W is meant to recall that the local pairing depends on the choice of W_v when $v|p$; when $v \nmid p$, it has no meaning. Although the symbol also depends on ℓ , we will usually omit it from the notation.

of abelian *fppf* sheaves on $\mathcal{O}_{E,v}$, and $\mathcal{Y}_{[D]}$ is represented by a smooth commutative group scheme. On the generic fibre, $\mathcal{Y}_{[D]} \otimes E_v$ can be identified with the complement Y_D of the zero section in the total space of the line bundle $\mathcal{O}(D)$ on A , and thus the extension admits a section

$$s_D: A \setminus |D| \rightarrow Y_D$$

which is canonical up to scaling.

Suppose given a morphism $\ell_{v,D,W}$ which makes the following diagram commute:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_{E,v}^\times \hat{\otimes} L & \longrightarrow & \mathcal{Y}_{[D]}(\mathcal{O}_{E,v}) \hat{\otimes} L & \longrightarrow & \mathcal{A}^0(\mathcal{O}_{E,v}) \hat{\otimes} L \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & E_v^\times \hat{\otimes} L & \longrightarrow & Y_D(E_v) \hat{\otimes} L & \longrightarrow & A(E_v) \hat{\otimes} L \longrightarrow 0 \\ & & \downarrow \ell_v & & \downarrow \ell_{v,D,W} & & \\ & & L & \xlongequal{\quad} & L & & \end{array}$$

Then we can define the local pairing by

$$(5.1.1) \quad \langle D, z \rangle_{v,W} = \ell_{v,D,W}(s_D(z)),$$

where s_D is extended to the divisor z in the obvious way. Notice that since z has degree zero, this is well-defined independently of the scaling ambiguity in s_D .

When $v \nmid p$, the logarithm ℓ_v vanishes on $\mathcal{O}_{E,v}^\times$ for topological reasons and we can uniquely extend it to an $\ell_{v,D}$ as in the above diagram by requiring its restriction to $\mathcal{Y}_{[D]}(\mathcal{O}_{E,v})$ to be trivial. When $v|p$, given the splitting W_v one can construct a section

$$s_{v,D,W}: A(E_v) \hat{\otimes} L \rightarrow Y_D(E_v) \hat{\otimes} L$$

and define the extension $\ell_{v,D,W}$ by requiring it to be trivial on the image of $s_{v,D,W}$. The standard construction is explained e.g. in [31, §3.2]. In the ordinary case, when W_v is chosen to be the unit root subspace, the crucial properties of the (canonical) local symbol are the last two in Proposition 5.1.2 below; in this case the construction rests on the following result (see [45] or [36, §6.9]).

Lemma 5.1.1. — *Let $E_{v,\infty}$ be a totally ramified \mathbf{Z}_p -extension of E_v , and denote by $E_{v,n}$ its n^{th} layer. Let $e \in \text{End}(A) \otimes \overline{\mathbf{Q}}$ be an idempotent. Assume that eV is ordinary as a Galois representation. Then the module of universal norms*

$$U(eA(E_v)) = \bigcap_n \text{Im} \left[\text{Tr}_{E_{v,n}/E_v} : eA(E_{v,n}) \rightarrow eA(E_v) \right]$$

has finite index in $eA(E_v)$.

Proposition 5.1.2. — *The p -adic local symbol*

$$\langle \cdot, \cdot \rangle_v = \langle \cdot, \cdot \rangle_{v,W}: (\mathcal{D}_0(A)(E_v) \times Z_0(A)^0(E_v))_e \rightarrow L$$

defined by (5.1.1) has the following properties (valid whenever they make sense).

1. It is bilinear.
2. If $h \in E_v(A)$ is a rational function, we have

$$\langle (h), z \rangle_v = \ell_v(h(z))$$

where if $z = \sum n_p P$, $h(z) = \prod h(P)^{n_p}$.

3. If $\phi: A \rightarrow A$ is an isogeny, we have

$$\langle \phi^* D, z \rangle_v = \langle D, \phi_* z \rangle_v.$$

4. For any $D \in \mathcal{D}_0(A)(E_v)$, $x_0 \in A(E_v) \setminus |D|$ the map from $A(E_v) \setminus |D| \rightarrow L$ defined by

$$x \mapsto \langle D, x - x_0 \rangle_v$$

is continuous.

5. (Compatibility.) Let E'_w/E_v be a finite extension, and let. If $D \in \mathcal{D}_0(A)(E'_w)$, $z \in Z_0(A)^\circ(E_v)$, we have

$$\langle \text{Tr}_{E'_w/E_v}(D'), z \rangle_v = \langle D', z \rangle_w$$

where $\langle \cdot, \cdot \rangle_w$ is the local pairing associated to $\ell_w = \ell_v \circ N_{E'_w/E_v}$ and (if $v|p$) the splitting W_w is induced from W_v .

6. (Boundedness.) Let $E_{v,\infty}^\ell = \cup_n E_{v,n}^\ell$ be the ramified⁽³⁰⁾ \mathbf{Z}_p -extension of E_v determined by the isomorphism

$$E_v^\times \supset \text{Ker}(\ell_v) \cong \text{Gal}(E_{v,\infty}^\ell/E) \subset \text{Gal}(E_v^{\text{ab}}/E).$$

induced from class field theory. In the ordinary situation of Lemma 5.1.1, if $e W_v$ is the unit root subspace of eV , there is a nonzero constant $c \in \mathbf{Z}_p$ such that

$$\langle D, z \rangle_{v,n} \in c^{-1} \ell_w(E_{v,n}^{\ell,\times})$$

if $D \in e\mathcal{D}_0(A)(E_{v,n})$, $z \in eZ_0(A)^\circ(E_{v,n})$ and $\langle \cdot, \cdot \rangle_{v,n}$ is the local pairing associated to the extension $E_{v,n}^\ell/E_v$ as in 5.

If $v \nmid p$, the local symbol is characterised by properties 1–4.

We refer to [37, II.1], [30, §4.2] and references therein for the proof and more details on the construction. See also Proposition 5.4.1 below.

5.2. The p -adic height pairing. — Let A be an abelian variety over a number field E . Let $\ell: E^\times \setminus E_A^\times \rightarrow \mathbf{Q}_p$ be a homomorphism (which we call a *global p -adic logarithm*), whose restrictions $\ell_v = \ell|_{E_v^\times}$ are ramified for all $v|p$. Let W_v be Hodge splittings at the places $v|p$ as in §5.1. Then we can define a height pairing

$$\langle \cdot, \cdot \rangle: A^\vee(E) \times A(E) \rightarrow L$$

as the sum of local height pairings

$$\langle x, y \rangle = \sum_v \langle \tilde{x}, \tilde{y} \rangle_v,$$

where \tilde{x} is a divisor on A whose class in $A^\vee(E) \cong \text{Pic}^0(A)$ is x , and $\tilde{y} = \sum n_p [P]$ is a zero-cycle of degree zero on A with support disjoint from the support of \tilde{x} , which satisfies $\sum n_p P = y$. The result is independent of the choices of \tilde{x} and \tilde{y} .

Let X be a (proper, smooth) curve over E of genus $g \geq 1$, together with a degree 1 divisor class defined over E inducing an embedding

$$\iota: X \hookrightarrow J(X)$$

⁽³⁰⁾Recall that we choose ℓ_w to be ramified.

into its Albanese variety $J(X)$.⁽³¹⁾ Let $\text{Div}(X)$ be the group of divisors on X , $\text{Div}^0(X)$ the subgroup of degree zero divisors, and similarly $\text{CH}(X) = \text{Div}^0(X)/\sim$, $\text{CH}(X)_0 = \text{Div}^0(X)/\sim$ the Chow group of zero-cycles modulo rational equivalence and its subgroup of degree zero elements. Then, given a p -adic logarithm and Hodge splittings for $V_p J(X)$, we can define local and global pairings on degree zero divisors on X (denoted with a subscript X) from the above pairings on $J(X)$ (here denoted with a subscript $J(X)$). Let D_1, D_2 be divisors of degree zero on X defined over E and with disjoint support. The morphism ι induces an isomorphism $\iota^*: \text{Pic}^0 J(X) \cong \text{Pic}^0(X)$, hence we can pick an algebraically trivial divisor D'_1 on $J(X)$ satisfying $D_1 = \iota^* D'_1 + (b)$ for some rational function $b \in E(X)$. If D'_1 is chosen so that its support is disjoint from the support of $\iota_* D_2$ and the support of (b) is disjoint from the support of D_2 , we can define

$$\langle D_1, D_2 \rangle_{v,X} = -\langle D_1, \iota_* D_2 \rangle_{v,J(X)} - \ell_v(b(D_2)),$$

and

$$\langle D_1, D_2 \rangle_X = \sum_v \langle D_1, D_2 \rangle_{v,X}.$$

The latter pairing descends to a **height pairing** on divisor classes

$$\langle \cdot, \cdot \rangle: \text{CH}(X)_0 \times \text{CH}(X)_0 \rightarrow L.$$

There are various conventions in the literature for the normalisation of the signs of height pairings. Our choices are the same as those of Kobayashi [30, §4.3], whose discussion we have followed and to which we refer for a comparison with other authors' choices.

5.3. p -adic Arakelov theory – local aspects. — Here and in §5.4 we summarise the main results of Besser [4], who develops the p -adic analogue of classical Arakelov theory.

Metrised line bundles. — Let X_v be a proper smooth variety over the finite extension E_v of \mathbf{Q}_p , and fix a ramified local p -adic logarithm $\ell_v: E_v^\times \rightarrow \mathbf{Q}_p$ which we extend to $\overline{\mathbf{Q}_p}^\times$ by $\ell_v|_{E_v'^\times} = \ell_v \circ N_{E_v'/E_v}$ for any finite extension E_v'/E_v .

A *metrised line bundle* $\widehat{\mathcal{L}} = (\mathcal{L}, \log_{\mathcal{L}})$ on X_v is a line bundle on X_v together with a choice of a *log function* $\log_{\mathcal{L},v}$ on the total space of $\mathcal{L}_v = \mathcal{L}|_{X \otimes E_v}$ minus the zero section (which will also be viewed as a function on the nonzero sections of \mathcal{L}_v). A log function is the analogue in the p -adic theory of the logarithm of a metric on the sections of \mathcal{L} on an infinite fibre of \mathcal{X} . It is a Coleman function having a certain analytic property⁽³²⁾ and the following algebraic property. If the p -adic logarithm ℓ_v factors as

$$(5.3.1) \quad \ell_v = t_v \circ \log_v$$

for some $\log_v: E_v^\times \rightarrow E_v$ and some \mathbf{Q}_p -linear $t_v: E_v \rightarrow \mathbf{Q}_p$, then for any nonzero section s of \mathcal{L}_v and rational function $f \in E(X_v)$ we have

$$(5.3.2) \quad \log_{\mathcal{L},v}(fs) = \log_v(f) + \log_{\mathcal{L},v}(s).$$

Adding a constant to a log function produces a new log function; this operation is called *scaling*.

⁽³¹⁾In our applications, we only have a rational divisor class, inducing a compatible system of maps $\iota_{E'}: X(E') \otimes \mathbf{Q} \rightarrow J(X)(E') \otimes \mathbf{Q}$ for E' a finite extension of E , such that for some integer n , $(n \iota_{E'})_{E'}$ is induced from an E -morphism. This causes no extra difficulties.

⁽³²⁾For which we refer to [4, Definition 4.1].

One can define a notion of $\bar{\partial}\partial$ -operator on Coleman functions, and attach to any log function $\log_{\mathcal{L}}$ on \mathcal{L} its *curvature* $\bar{\partial}\partial \log_{\mathcal{L}} \in H_{\text{dR}}^1(X_v) \otimes \Omega^1(X_v)$; its cup product is the first Chern class of \mathcal{L} .

Log functions on a pair of line bundles induce in the obvious way a log function on their tensor product, and similarly for the dual of a line bundle. If $\pi : X_v \rightarrow Y_v$ is a morphism, then a log function on a line bundle on Y_v induces in the obvious way a log function on the pullback line bundle on X . If moreover π is a finite Galois cover with Galois group G , and \mathcal{L} is a line bundle on X_v with log function $\log_{\mathcal{L}}$ and associated curvature β , then the norm line bundle $N_{\pi}\mathcal{L}$ on Y_v with stalks

$$(N_{\pi}\mathcal{L})_y = \bigotimes_{x \mapsto y} \mathcal{L}_x^{\otimes e(x|y)}$$

has an obvious candidate log function $N_{\pi}\log_{\mathcal{L}}$ obtained by tensor product. A delicate point is that it is not automatic that the latter is a genuine log function (i.e. it satisfies the analytic property alluded to above); cf. [4, Proposition 4.8] for a sufficient condition.

The canonical Green function. — Now let X_v/E_v be a curve of genus $g \geq 1$ with good reduction above p . Choose a splitting $W_v \subset H_{\text{dR}}^1(X_v) \otimes L$ of the Hodge filtration as in §5.1, which we use to identify $W_v \cong \Omega^1(X_v)^{\vee}$; we then define a canonical element

$$\mu_{X_v} = \frac{1}{g} \text{id} \in \text{End} \Omega^1(X_v) \cong W_v \otimes \Omega^1(X_v)$$

and similarly for the self-product $X_v \times X_v$ (denoting π_1, π_2 the projections)

$$\Phi = \begin{pmatrix} \frac{1}{g} & -1 \\ -1 & \frac{1}{g} \end{pmatrix} \in \text{End}(\pi_1^* \Omega^1(X_v) \oplus \pi_2^* \Omega^1(X_v)) \hookrightarrow H_{\text{dR}}^1(X_v \times X_v) \otimes \Omega^1(X_v \times X_v).$$

The first Chern class of Φ is the class of the diagonal $\Delta \subset X_v \times X_v$.

Let s_{Δ} denote the canonical section of the line bundle $\mathcal{O}(\Delta)$ on $X_v \times X_v$. Given any log function $\log_{\mathcal{O}(\Delta)}$ on $\mathcal{O}(\Delta)$ with curvature Φ , we can consider the function G on $X_v \times X_v$ given by

$$G(P, Q) = \log_{\mathcal{O}(\Delta)}(s_{\Delta})(P, Q).$$

It is a Coleman function with singularities along Δ ; we call G a *Green function* for X_v .

A Green function G induces a log function on any line bundle $\mathcal{O}(D)$ on X_v by

$$\log_{\mathcal{O}(D)}(s_D)(Q) = \sum n_i G(P_i, Q)$$

if $D = \sum n_i P_i$ and s_D is the canonical section of $\mathcal{O}(D)$. A log function $\log_{\mathcal{L}}$ on the line bundle \mathcal{L} and the resulting metrised line bundle $(\mathcal{L}, \log_{\mathcal{L}})$ are called *admissible* with respect to G if for one (equivalently, any) nonzero rational section s of \mathcal{L} , the difference $\log_{\mathcal{L}}(s) - \log_{\text{div}(s)}$ is a constant. Such a constant is denoted by $\iota_{\log_{\mathcal{L}}}(s)$, or $\iota_{\log_v}(s)$ in the case of the trivial line bundle with the log function \log_v . It is the analogue of the integral of the norm of s . It follows easily from the definitions that any isomorphism of admissible metrised line bundles is an isometry up to scaling.

Let ω_{X_v} be the canonical sheaf on X_v . The canonical isomorphism $\omega_{X_v} \cong \Delta^* \mathcal{O}(-\Delta)$ gives another way to induce from G a log function $\log_{\omega_{X_v}}^G$ on ω_{X_v} , namely by pullback (and the resulting metrised line bundle has curvature $(2g - 2)\mu_{X_v}$). The requirement that this log function be admissible, together with a symmetry condition, leads to an almost unique choice of G .

Proposition 5.3.1 ([4, Theorem 5.10]). — *There exists a unique up to constant symmetric Green function G with associated curvature Φ such that $(\omega_{X_v}, \log_{\omega_{X_v}}^G)$ is an admissible metrised line bundle with respect to G .*

In the following we will arbitrarily fix the constant implied by the Proposition. In our context, the canonical Green function thus determined is, in a suitable sense, defined over E_v [4, Proposition 8.1].

5.4. p -adic Arakelov theory – global aspects. — Let E be a number field with ring of integers \mathcal{O}_E . Let $\mathcal{X}/\mathcal{O}_E$ be an arithmetic surface with generic fibre X , that is, $\mathcal{X} \rightarrow \mathcal{O}_E$ is a proper regular relative curve and $\mathcal{X} \otimes_{\mathcal{O}_E} E = X$. We assume that \mathcal{X} has *good reduction* at all place $v|p$, and denote $X_v = \mathcal{X} \otimes E_v$. Fix choices of a ramified p -adic logarithm ℓ and Hodge splittings W_v as in §5.2.

Arakelov line bundles and divisors. — An *Arakelov line bundle* on \mathcal{X} is a pair

$$\widehat{\mathcal{L}} = (\mathcal{L}, (\log_{\mathcal{L}_v})_{v|p})$$

consisting of a line bundle \mathcal{L} on \mathcal{X} together with admissible (with respect to the Green functions of Proposition 5.3.1) log functions $\log_{\mathcal{L}_v}$ on $\mathcal{L}_v = \mathcal{L}|_{X_v}$. We denote by $\text{Pic}^{\text{Ar}}(\mathcal{X})$ the group of isometry classes of Arakelov line bundles on \mathcal{X} .

The group $\text{Div}^{\text{Ar}}(\mathcal{X})$ of *Arakelov divisors* on \mathcal{X} is the group of formal combinations

$$D = D_{\text{fin}} + D_{\infty}$$

where D_{fin} is a divisor on \mathcal{X} and $D_{\infty} = \sum_{v|p} \lambda_v X_v$ is a sum with coefficients $\lambda_v \in E_v$ of formal symbols X_v for each place $v|p$ of E . To an Arakelov line bundle $\widehat{\mathcal{L}}$ and a nonzero rational section s of \mathcal{L} we associate the Arakelov divisor

$$\widehat{\text{div}}(s) = (s)_{\text{fin}} + (s)_{\infty}$$

where $(s)_{\text{fin}}$ is the usual divisor of s and $(s)_{\infty} = \sum_{v|p} \ell_{\log_{\mathcal{L}_v}}(s_v) X_v$. The group $\text{Prin}^{\text{Ar}}(\mathcal{X})$ of *principal* Arakelov divisors on \mathcal{X} is the group generated by the $\widehat{\text{div}}(b)$ for $b \in E(\mathcal{X})^{\times}$. The Arakelov Chow group of \mathcal{X} is

$$\text{CH}^{\text{Ar}}(\mathcal{X}) = \text{Div}^{\text{Ar}}(\mathcal{X}) / \text{Prin}^{\text{Ar}}(\mathcal{X}),$$

and we have an isomorphism

$$\text{Pic}^{\text{Ar}}(\mathcal{X}) \cong \text{CH}^{\text{Ar}}(\mathcal{X})$$

given by $\widehat{\mathcal{L}} \rightarrow [\widehat{\text{div}}(s)]$ for any rational section s of \mathcal{L} .

The p -adic Arakelov pairing. — Most important for us is the existence of a pairing on $\text{CH}^{\text{Ar}}(\mathcal{X})$, extending the p -adic height pairing of divisors of §5.2. Let $(,)_v$ denote the (\mathbf{Z} -valued) intersection pairing of cycles on \mathcal{X}_v with disjoint support.

Proposition 5.4.1 (Besser [4]). — *Let $\mathcal{X}/\mathcal{O}_E$ be an arithmetic surface with good reduction above p . For any choice of ramified p -adic logarithm $\ell : E_{\mathbf{A}}^{\times}/E^{\times} \rightarrow \mathbf{Q}_p$ and Hodge splittings $(W_v)_{v|p}$ as above, there is a symmetric bilinear paring⁽³³⁾*

$$\langle , \rangle^{\text{Ar}} : \text{CH}^{\text{Ar}}(\mathcal{X}) \times \text{CH}^{\text{Ar}}(\mathcal{X}) \rightarrow L$$

⁽³³⁾The notation of [4] is $D_1 \cdot D_2$ for $\langle D_1, D_2 \rangle^{\text{Ar}}$.

satisfying:

1. If D_1 and D_2 are finite and of degree zero on the generic fibre, and one of them has degree zero on each special fibre of \mathcal{X} , then

$$\langle D_1, D_2 \rangle^{\text{Ar}} = \langle D_{1,E}, D_{2,E} \rangle,$$

where $D_{i,E} \in \text{Div}^0(X)$ is the generic fibre of D_i and $\langle \cdot, \cdot \rangle$ denotes the height pairing of Proposition 5.1.2 associated with the same choices of ℓ and W_v .

2. If $D_{1,\text{fin}}, D_{2,\text{fin}}$ have disjoint supports on the generic fibre, then

$$\langle D_1, D_2 \rangle^{\text{Ar}} = \sum_v \langle D_1, D_2 \rangle_v^{\text{Ar}}$$

where the sum runs over all finite places of E , and the local Arakelov pairings are defined by

$$\langle D_1, D_2 \rangle_v^{\text{Ar}} = (D_1, D_2)_v \ell_v(\pi_v)$$

for $v \nmid p$ and below for $v|p$.

If moreover we are in the situation of 1, then for each place v we have

$$\langle D_1, D_2 \rangle_v^{\text{Ar}} = \langle D_{1,E}, D_{2,E} \rangle_v.$$

3. In the situation of 2, if moreover $D_1 = \widehat{\text{div}}(h)$ is the Arakelov divisor of a rational function h , then

$$\langle D_1, D_2 \rangle_v^{\text{Ar}} = \ell_v(h(D_{2,\text{fin}}))$$

for all places v .

For completeness, we give the description of the local pairing at $v|p$ of divisors with disjoint supports. If $\ell_v = t_v \circ \log_v$ as in (5.3.1) and G_v is the Green function on $X_v \times X_v$, we have $\langle D, X_w \rangle_v^{\text{Ar}} = 0$ if $v \neq w$, $\langle X_v, X_v \rangle_v^{\text{Ar}} = 0$, $\langle D, \lambda_v X_v \rangle_v^{\text{Ar}} = (\deg D_E) t_v(\lambda_v)$ and if D_1, D_2 are finite divisors with images $D_{1,v} = \sum n_i P_i, D_{2,v} = \sum m_j Q_j$ in X_v then

$$\langle D_1, D_2 \rangle_v^{\text{Ar}} = \sum_{i,j} n_i m_j t_v(G_v(P_i, Q_j)).$$

In fact, in [4] it is proved directly that the global Arakelov pairing and its local components at p coincide with the global and local height pairings of Coleman–Gross [11]. The latter coincide with the Zarhin–Nekovář pairings by [3].

6. Heegner points on Shimura curves

In this section we describe our Shimura curve and construct Heegner points on it, following [57, §§1-2], to which we refer for the details (see also [58, §5], and [9] for the original source of many results on Shimura curves). We go back to our usual notation, so F is a totally real number field of degree g , N is an ideal of \mathcal{O}_F , E is a CM extension of F of discriminant Δ coprime to $2Np$, and ε is its associated Hecke character.

6.1. Shimura curves. — Let B be a quaternion algebra over F which is ramified at all but one infinite place. Then we can choose an isomorphism $B \otimes \mathbf{R} \cong M_2(\mathbf{R}) \oplus \mathbf{H}^{g-1}$, where \mathbf{H} is the division algebra of Hamilton quaternions. There is an action of B^\times on $\mathfrak{H}^\pm = \mathbf{C} - \mathbf{R}$ by Möbius

transformations via the map $B^\times \rightarrow \mathbf{GL}_2(\mathbf{R})$ induced from the above isomorphism. For each open subgroup K of $\widehat{B}^\times = (B \otimes_F \widehat{F})^\times$ which is compact modulo \widehat{F}^\times we then have a **Shimura curve**

$$M_K(\mathbf{C}) = B^\times \backslash \mathfrak{H}^\pm \times \widehat{B}^\times / K,$$

where $\mathfrak{H}^\pm = \mathbf{C} \setminus \mathbf{R}$. Unlike modular curves, the curves M_K do not have a natural moduli interpretation. However, by the work of Carayol [9], $M_K(\mathbf{C})$ has a finite map⁽³⁴⁾ to another (unitary) Shimura curve $M'_{K'}(\mathbf{C})$ which, if the level K' is small enough, has an interpretation as the moduli space of certain quaternionic abelian varieties. Namely, $M'_{K'}$ parametrises isomorphism classes of abelian varieties of dimension $4[F : \mathbf{Q}]$ with multiplication by the ring of integers $\mathcal{O}_{B'}$ of $B \otimes_F F'$ and some extra structure (a polarisation and a K' -level structure, compatible with the quaternionic multiplication) [57, Proposition 1.1.5].

We will usually denote a point of $M'_{K'}$ simply by $[A]$ where A is the underlying abelian variety. If K' has maximal components at places dividing m , one can define a notion of an *admissible submodule* D of level m (see [57, §1.4.3]): it is an $\mathcal{O}_{B'}$ -submodules of $A[m]$ satisfying a certain condition, which ensures that the quotient A/D can be naturally endowed with the extra structure required by the functor $M'_{K'}$. We denote by $[A_D]$ the object whose underlying abelian variety is A/D , with the induced extra structure.

As a consequence of the moduli interpretation, the curve $M_K(\mathbf{C})$ has a canonical model M_K defined over F (it is connected but not, in general, geometrically connected), and a proper regular integral model⁽³⁵⁾ \mathcal{M}_K over \mathcal{O}_F ; if v is a finite place where B is split, then \mathcal{M}_K is smooth over $\mathcal{O}_{F,v}$ if K_v is a maximal compact subgroup of B_v and K^v is sufficiently small. We denote $\mathcal{M}_{K,v} = \mathcal{M}_K \otimes \mathcal{O}_{F,v}$.

Universal formal group and ordinary points. — Assume that the level structure K is maximal at \wp . The curve $\mathcal{M}_{K,\wp}$ carries a universal \wp -divisible $\mathcal{O}_{B,\wp}$ -module \mathcal{G} obtained from the \wp -divisible group $\mathcal{A}[\wp^\infty]$ of the universal abelian scheme \mathcal{A} over $M'_{K',\wp}$. More precisely, choosing an auxiliary quadratic field F' which is split at \wp and an isomorphism $j : \mathcal{O}_{F',\wp} \cong \mathcal{O}_{F,\wp} \oplus \mathcal{O}_{F,\wp}$, we have

$$\mathcal{G} = \mathcal{A}[\wp^\infty]^{(2)} = e_2 \mathcal{A}[\wp^\infty],$$

where e_2 is the idempotent in $\mathcal{O}_{F',\wp}$ corresponding to $(0, 1)$ under j .

Assume that B is split at \wp . Then we denote by $\mathcal{G}^1, \mathcal{G}^2$ the images under the projectors corresponding to $\begin{pmatrix} 1 & \\ & 0 \end{pmatrix}$ and $\begin{pmatrix} 0 & \\ & 1 \end{pmatrix}$ under a fixed isomorphism $B \cong M_2(F_\wp)$; they are isomorphic via the element $\begin{pmatrix} & 1 \\ -1 & \end{pmatrix}$.

Let x be a geometric point of the special fibre $\mathcal{M}_{K,\wp}$. Then the \mathcal{G}_x^i are divisible \mathcal{O}_\wp -modules of dimension 1 and height 2, hence isomorphic to either of:

- the direct sum $\Sigma_1 \oplus F_\wp / \mathcal{O}_{F,\wp}$, where Σ_1 is the unique formal $\mathcal{O}_{F,\wp}$ -module of height 1 – in this case x is called *ordinary*;
- the unique formal \mathcal{O}_\wp -module of dimension 1 and height 2 – in this case x is called *supersingular*.

⁽³⁴⁾Which is an embedding if $K \supset \widehat{F}^\times$.

⁽³⁵⁾In the modular curve case $F = \mathbf{Q}$, $\varepsilon(v) = 1$ for all $v|N$, M_K and \mathcal{M}_K are proper only after the addition of finitely many cusps. (We caution the reader that Carayol [9] uses the notation \mathcal{M}_K to denote instead the set of geometrically connected components of M_K .)

Let $M_K(\overline{\mathbf{Q}}_\varphi)^{\text{ord}} \subset M_K(\overline{\mathbf{Q}}_\varphi)$ be the set of points with ordinary reduction. Then the Frobenius map Frob_φ admits a lift

$$(6.1.1) \quad \varphi: M_K(\overline{F}_\varphi)^{\text{ord}} \rightarrow M_K(\overline{F}_\varphi)^{\text{ord}}$$

given in the moduli interpretation by $[A] \mapsto [A_{\text{can}(A)}]$, where $\text{can}(A)$ is the *canonical submodule* of A , that is the sub- $\mathcal{O}_{F,\varphi}$ -module of $A[\varphi]$ in the kernel of multiplication by φ in the formal group of A .

The order R and the curve X . — Assume that $\varepsilon(N) = (-1)^{g-1}$. Then the quaternion algebra \mathbf{B} over \mathbf{A}_F ramified exactly at all the infinite places and the finite places $v|N$ such that $\varepsilon(v) = -1$ is *incoherent*, that is, it does not arise via extension of scalars from a quaternion algebra over F . On the other hand, for any embedding $\tau: F \hookrightarrow \mathbf{R}$, there is a *nearby* quaternion algebra $B(\tau)$ defined over F and ramified at τ and the places where \mathbf{B} is ramified. Fix any embedding $\rho: E \rightarrow B(\tau)$, and let R be an order of $\widehat{B} = \widehat{B}(\tau)$ which contains $\rho(\mathcal{O}_E)$ and has discriminant N (this is constructed in [57, §1.5.1]). Then the curve X over F of interest to us is the (compactification of) the curve M_K defined above for the subgroup $K = \widehat{F}^\times \widehat{R}^\times \subset \widehat{B}$; that is, for each embedding $\tau: F \rightarrow \mathbf{C}$, we have

$$(6.1.2) \quad X(\mathbf{C}) = B(\tau)^\times \backslash \mathfrak{H}^\pm \times \widehat{B}^\times / \widehat{F}^\times \widehat{R}^\times \cup \{\text{cusps}\}.$$

The finite set of cusps is nonempty only in the classical case where $F = \mathbf{Q}$, $\varepsilon(v) = 1$ for all $v|N$ so that $X = X_0(N)$. In what follows we will not burden the notation with the details of this particular case, which poses no additional difficulties and is already treated in the original work of Perrin-Riou [41].

We denote by \mathcal{X} the canonical model of X over \mathcal{O}_F , and by \mathcal{X}_v its base change to $\mathcal{O}_{F,v}$. We also denote by $J(X)$ the Albanese variety of X and by \mathcal{J}_v its Néron model over $\mathcal{O}_{F,v}$.

Hecke correspondences. — Let m be an ideal of \mathcal{O}_F which is coprime to the ramification set of B . Let $\gamma_m \in \widehat{\mathcal{O}}_B$ be an element with components 1 away from m and such that $\det \gamma_m$ generates m at the places dividing m . Then the Hecke operator $T(m)$ on X is defined by

$$T(m)[(z, g)] = \sum_{\gamma \in K\gamma_m K/K} [(z, g\gamma)]$$

under the complex description (6.1.2). When m divides N we often denote the operator $T(m)$ by $U(m)$ or U_m .

Let \mathbf{T}'_N be the algebra generated by the $T(m)$ for m prime to N . Then by [57, Theorem 3.2.1], the algebra \mathbf{T}'_N is a quotient of the Hecke algebra on Hilbert modular forms \mathbf{T}_N (hence the names $T(m)$ are justified). It acts by correspondences on $X \times X$, and taking Zariski closures of cycles on $\mathcal{X} \times \mathcal{X}$ extends the action to \mathcal{X} .

As in the classical case, the Hecke operators $T(m)$ admit a moduli interpretation, after base change to a suitable quadratic extension F' and passing to the curve X' . Namely we have

$$T(m)[A] = \sum_D [A_D],$$

where the sum runs over the admissible submodules of A of level m .

6.2. Heegner points. — The curve X defined above has a distinguished collection of points defined over abelian extensions of E : we briefly describe it, referring the reader to [57, §2] for more details.

A point y of X is called a **CM point** with multiplication by E if it can be represented by $(x_0, g) \in \mathfrak{H}^\pm \times \widehat{B}^\times$ via (6.1.2), where $x_0 \in \mathfrak{h}^+$ is the unique point fixed by E^\times . The order

$$\text{End}(y) = g \widehat{R} g^{-1} \cap \rho(E)$$

in $E = \rho(E)$ is defined independently of the choice of g , and

$$\text{End}(y) = \mathcal{O}_E[c] = \mathcal{O}_F + c \mathcal{O}_E$$

for a unique ideal c of \mathcal{O}_F called the **conductor** of y . We say that the point $y = [(x_0, g)]$ has the *positive orientation* if for every finite place v the morphism $t \mapsto g^{-1} \rho(t) g$ is R_v^\times -conjugate to ρ in $\text{Hom}(\mathcal{O}_{E,v}, R_v)/R_v^\times$.⁽³⁶⁾ Let Y_c be the set of positively oriented CM points of conductor c . By the work of Shimura and Taniyama, it is a finite subscheme of X defined over E , and the action of $\text{Gal}(\overline{\mathbf{Q}}/E)$ is given by

$$\sigma([(x_0, g)]) = [(x_0, \text{rec}_E(\sigma)g)],$$

where $\text{rec}_E: \text{Gal}(\overline{E}/E) \rightarrow \text{Gal}(\overline{E}/E)^{\text{ab}} \xrightarrow{\sim} \overline{E}^\times \backslash \widehat{E}^\times$ is the reciprocity map of class field theory. If $y = [(x_0, g)]$ has conductor c , then the action factors through

$$\text{Gal}(H[c]/E) \cong E^\times \backslash \widehat{E}^\times / \widehat{F}^\times \widehat{\mathcal{O}}_E[c]^\times$$

where $H[c]$ is the ring class field of E of conductor c ; the action of this group on Y_c is simply transitive.

For each nonzero ideal c of \mathcal{O}_F , let $u(c) = [\mathcal{O}_E[c]^\times : \mathcal{O}_F^\times]$ and define the divisor

$$(6.2.1) \quad \eta_c = u(c)^{-1} \sum_{y \in Y_c} y.$$

Let $\eta = \eta_1$. By the above description of the Galois action on CM points, each divisor η_c is defined over E .

A **Heegner point** $y \in X(H)$ is a positively oriented CM point with conductor 1. We can use the embedding $\iota: X \rightarrow J(X) \otimes \mathbf{Q}$ to define the point

$$[z] = \iota(\eta) = [\eta] - b[\xi] \in J(X)(E) \otimes \mathbf{Q}$$

where b is a number such that $[z]$ has degree zero in each geometrically connected component of X , and $[\xi]$ is the Hodge class of the Introduction (see below for more on the Hodge class).

Arakelov Heegner divisors. — The Heegner divisor on X can be refined to an Arakelov divisor \hat{z} having degree zero on each irreducible component of each special fibre. On a suitable Shimura curve $\tilde{X} \xrightarrow{\pi} X$ of deeper level away from $N\Delta_{E/F}$, we can give an explicit description of the pullback $\tilde{\hat{z}}$ of \hat{z} and of the Hodge class as follows.

As outlined in in §6.1, after base change to a suitable quadratic extension F' of F , we have an embedding $\tilde{X} \hookrightarrow \tilde{X}'$ of $\tilde{X} = M_{\tilde{K}}$ into the unitary Shimura curve $\tilde{X}' = M'_{\tilde{K}'}$ parametrising

⁽³⁶⁾This set has two elements only if $v|N$ (the other element is called the negative orientation at v); otherwise it has one element and the condition at v is empty. There is a group of Atkin-Lehner involutions acting transitively on orientation classes.

abelian varieties of dimension $4g$ with multiplication by $\mathcal{O}_{B'}$ and some extra structure. Then by the Kodaira–Spencer map, we have an isomorphism $\omega_{\tilde{X}'} \cong \det \text{Lie } \mathcal{A}^\vee|_{\tilde{X}'}$, where $\mathcal{A} \rightarrow \tilde{X}'$ is the universal abelian scheme and the determinant is that of an $\mathcal{O}_{F'}$ -module of rank 4 (the structure of $\mathcal{O}_{F'}$ -module coming from the multiplication by $\mathcal{O}_{B'}$ on \mathcal{A}). This gives a way⁽³⁷⁾ of extending the line bundle $\omega_{\tilde{X}'}$ to the integral model $\tilde{\mathcal{X}}'$ and to a line bundle \mathcal{L} on $\tilde{\mathcal{X}}$. For each finite place $v|p$ we endow $\mathcal{L}|_{\tilde{\mathcal{X}}_v}$ with the canonical log functions $\log_{\mathcal{L},v}$ coming from the description $\mathcal{L}|_{\tilde{\mathcal{X}}_v} = \omega_{\tilde{\mathcal{X}}_v}$ and a fixed choice of Hodge splittings on $\tilde{\mathcal{X}}$. We define $[\tilde{\xi}] \in \text{CH}^{\text{Ar}}(\tilde{\mathcal{X}}) \otimes \mathbb{Q}$ to be the class of $(\mathcal{L}, (\log_{\mathcal{L}})_v|_p)$ divided by its degree, $[\tilde{\xi}]$ to be its finite part, and $\tilde{\xi}$ to be any Arakelov divisor in its class.

Then the Arakelov Heegner divisor $\hat{z} \in \text{Div}^{\text{Ar}}(\mathcal{X} \otimes \mathcal{O}_E)$ is described by

$$(6.2.2) \quad \hat{z} = \hat{\eta} - h\tilde{\xi} + Z,$$

where $\hat{\eta}$ is the Zariski closure in $\mathcal{X} \otimes \mathcal{O}_E$ of the pullback of η to $\tilde{\mathcal{X}}$, and Z is a finite vertical divisor uniquely determined by the requirement that \hat{z} should have degree zero on each irreducible component of each special fibre.

6.3. Hecke action on Heegner points. — Recall from §1.5 the spaces of Fourier coefficients $\mathcal{D}_{\mathcal{N}} \subset \mathcal{S}$, the arithmetic functions $\sigma_1, r \in \mathcal{D}_{\mathcal{N}}$, and the space $\overline{\mathcal{S}} = \mathcal{S}/\mathcal{D}_{\mathcal{N}}$. The action of Hecke operators on the Arakelov Heegner divisor is described as follows.

Proposition 6.3.1. — *Let m be an ideal of \mathcal{O}_F coprime to N . We have*

1. $T(m)\eta = \sum_{c|m} r(m/c)\eta_c$.
2. Let $\eta_c^\circ = \sum_{\mathcal{O}_F \neq d|c} \eta_d$, and let $T^0(m)\eta = \sum_{c|m} \varepsilon(c)\eta_{m/c}^\circ$. Then η and $T^0(m)\eta$ have disjoint support and if m is prime to $N\Delta$ then $T(m)\eta = T^0(m)\eta + r(m)\eta$.
3. $T(m)[\xi] = \sigma_1(m)[\xi]$ and $m \mapsto T(m)\tilde{\xi}$ is zero in $\overline{\mathcal{S}} \otimes \text{Div}^{\text{Ar}}(\tilde{\mathcal{X}})$.
4. The arithmetic function $m \mapsto T(m)Z$ is zero in $\overline{\mathcal{S}} \otimes \text{Div}^{\text{Ar}}(\mathcal{X})$.

Proof. — Parts 1, 2 and 4 are proved in [57, §4]. For part 3, we switch to the curve $\tilde{\mathcal{X}}$. By definition $[\tilde{\xi}]$ is a multiple of the class of the Arakelov line bundle $\mathcal{L} = \det \text{Lie } \mathcal{A}^\vee$ on $\tilde{\mathcal{X}}$ with the canonical log functions on $\mathcal{L}_v \cong \omega_{\tilde{\mathcal{X}}_v}$, where $\mathcal{A} \rightarrow \tilde{\mathcal{X}}$ is the universal abelian scheme. We view $T(m)$ as a finite algebraic correspondence of degree $\sigma_1(m)$ induced by the subscheme $\tilde{\mathcal{X}}_m \subset \tilde{\mathcal{X}} \times \tilde{\mathcal{X}}$ of pairs $(A, A/D)$ where D is an admissible submodule of A of level m . If $p_1, p_2: \tilde{\mathcal{X}}_m \rightarrow \tilde{\mathcal{X}}$ are the two projections, then we have

$$T(m)\mathcal{L} = N_{p_1} p_2^* \mathcal{L},$$

and the log functions $\log_{T(m)\mathcal{L}_v}$ on $T(m)\mathcal{L}|_{\tilde{\mathcal{X}}_v}$ are the ones induced by this description. (That these are genuine log functions – cf. the caveat in §5.3 – will be shown in the course of proving Proposition 6.3.1.3 below.)

⁽³⁷⁾See [57, §4.1.3, §1] for more details on this construction.

Let $\pi: \mathcal{A}_1 \rightarrow \mathcal{A}_2$ be the universal isogeny over $\widetilde{\mathcal{X}}_m$. As $p_i^* \mathcal{L} = \det \text{Lie } \mathcal{A}_i^{\vee}$, we have an induced map

$$\psi_m = N_{p_2} \pi^*: T(m)\mathcal{L} \rightarrow N_{p_2} p_2^* \mathcal{L} = \mathcal{L}^{\sigma_1(m)},$$

and [57, §4.3] shows that $\psi_m(T(m)\mathcal{L}) = c_m \mathcal{L}^{\sigma_1(m)}$ where $c_m \subset \mathcal{O}_F$ is an ideal with divisor $[c_m]$ on $\text{Spec } \mathcal{O}_F$ such that $m \rightarrow [c_m]$ is a σ_1 -derivative (§1.5), hence zero in $\overline{\mathcal{S}} \otimes \text{Div}(\text{Spec } \mathcal{O}_F) \subset \overline{\mathcal{S}} \otimes \text{Div}^{\text{Ar}}(\mathcal{X})$. In fact if the finite divisor $\tilde{\xi}_{\text{fin}} = \text{div}(s)$ for a rational section s of \mathcal{L} , the same argument shows that $T(m)\tilde{\xi}_{\text{fin}} = \text{div}(T(m)s) = \sigma_1(m)\text{div}(s) + \text{div}(c_m)$, hence $m \mapsto \tilde{\xi}_{\text{fin}}$ is zero in $\overline{\mathcal{S}} \otimes \text{Div}^{\text{Ar}}(\mathcal{X})$.

We complete the proof by showing that, for each $v|p$, the difference of log functions

$$(6.3.1) \quad \psi_m^* \log_{\mathcal{L}_v^{\sigma_1(m)}} - \log_{T(m)\mathcal{L}_v}$$

on the line bundle $T(m)\mathcal{L}_v$ on $\widetilde{\mathcal{X}}_v$ is a constant on the total space of \mathcal{L}_v , and it is a σ_1 -derivative when viewed as a function of m . (In particular this shows that $\log_{T(m)\mathcal{L}_v} = \sigma_1(m)\psi_m^* \log_{\mathcal{L}_v} + \text{constant}$ is a genuine log function.)

It is enough to show this after pullback via p_1 on $\widetilde{\mathcal{X}}_m$, where (denoting pulled back objects with a prime) the map ψ'_m decomposes as

$$\psi'_m = \otimes_D \pi_D^*: \otimes_D \det \text{Lie}(\mathcal{A}'/D)^{\vee} \rightarrow (\det \text{Lie } \mathcal{A}'^{\vee})^{\otimes \sigma_1(m)}$$

where the tensor product runs over admissible submodule schemes of level m of \mathcal{A}' (since base change via p_1 splits the cover p_1 , there are exactly $\sigma_1(m)$ of those). Now the difference (6.3.1) is the sum of the $\sigma_1(m)$ differences

$$(\pi_D^*)^* \log_{\mathcal{L}} - \log_{\mathcal{L}},$$

which are all the same since they are permuted by the Galois group of p_1 . As π_D^* acts by multiplication by $(\deg \pi_D)^{1/2} = \mathbf{N}(m)^2$, by (5.3.2) each of these differences is $2 \log_v \mathbf{N}(m)$ so that (6.3.1) equals

$$2\sigma_1(m) \log_v \mathbf{N}(m)$$

which is indeed a σ_1 -derivative. \square

7. Heights of Heegner points

Let Ψ be the modular form of level N with Fourier coefficients given by the p -adic height pairing $\langle z, T(m)z \rangle$ (it is a modular form because of Lemma 1.4.1 and the fact that the quaternionic Hecke algebra \mathbf{T}'_N is a quotient of \mathbf{T}_N , as explained at the end of §6.1). We will compute the heights of Heegner points, with the goal of showing (in §8) that $l_{f_\alpha}(\Phi')$ and $l_{f_\alpha}(\Psi)$ are equal up to the action of some Hecke operators. The main theorem will follow.

The strategy is close to that of Perrin-Riou, namely we separate the local contributions to Ψ from primes above p , writing $\Psi \sim \Psi_{\text{fin}} + \Psi_p$; using the computations of [57, 58] we find an explicit expression for Ψ_{fin} , which in §8 we will show to be “almost” equal to the expression for Φ' , while the contribution of Ψ_p is shown to vanish. We circumvent the difficulties posed by the absence of cusps through the use of p -adic Arakelov theory.

It will be crucial to work in the quotient spaces $\overline{\mathcal{S}}, \overline{\mathcal{S}}^{\text{ord}}$ introduced in §1.6; the diagram (1.6.1) summarises the relations among them. We abuse notation by using the same name for a modular form and its image in $\overline{\mathcal{S}}_N^{\text{ord}}$.

The height pairings $\langle \cdot, \cdot \rangle$ (and the accompanying Arakelov pairings) on the base change of X to E that will be considered are the ones associated to a “cyclotomic” p -adic logarithm given by $\ell = \ell_F \circ \mathfrak{N}: E^\times \setminus E_{A^\infty}^\times \rightarrow \mathbf{Q}_p$ for some⁽³⁸⁾

$$\ell_F: F^\times \setminus F_{A^\infty}^\times \rightarrow \mathbf{Q}_p,$$

and to choices of Hodge splittings on $V_{v,L} = H_{\text{dR}}^1(X_v/E_v) \otimes L (v|p)$ such that on $e_f V_{v,L} \cong e_f M_{g,L}$, the induced Hodge splitting is the unit root splitting.

As mentioned before, the Shimura curve X and its integral model \mathcal{X} may not be fine enough for the needs of Arakelov and intersection theory, so that we may need to pass to a Shimura curve $\widetilde{\mathcal{X}} \xrightarrow{\pi} \mathcal{X}$ of deeper level away from p and consider the pullbacks $\tilde{\eta}$ of the divisors η , etc. Then notation such as $\langle \hat{\eta}, T^0(m)\hat{\eta} \rangle^{\text{Ar}}$ is to be properly understood as $\langle \tilde{\eta}, T^0(m)\tilde{\eta} \rangle^{\text{Ar}} / \deg \pi$.

7.1. Local heights at places not dividing p . — The next two results will be used to show the main identity.

Lemma 7.1.1. — *In the space $\overline{\mathcal{S}}$ we have*

$$\langle z, T(m)z \rangle = \langle \hat{z}, T(m)\hat{z} \rangle^{\text{Ar}} \sim \langle \hat{\eta}, T^0(m)\hat{\eta} \rangle^{\text{Ar}}.$$

Proof. — First observe that by Lemma 1.4.1, the first member is a modular form of level N , so it does indeed belong to $\overline{\mathcal{S}}_N$. The first equality is a consequence of Proposition 5.4.1.1 and the construction of \hat{z} . The second part follows from expanding the second term for m prime to $N\Delta$ according to (6.2.2) and observing that the omitted terms are zero in $\overline{\mathcal{S}}$ by Proposition 6.3.1. \square

We can therefore write

$$(7.1.1) \quad \Psi \sim \sum_w \Psi_w = \sum_v \Psi_v = \Psi_{\text{fin}} + \Psi_p$$

in $\overline{\mathcal{S}}$, with the first sum running over the finite places w of E , the second sum running over the finite places v of F , and

$$\Psi_w(m) = \langle \hat{\eta}, T^0(m)\hat{\eta} \rangle_w^{\text{Ar}}, \quad \Psi_v = \sum_{w|v} \Psi_w, \quad \Psi_{\text{fin}} = \sum_{v \nmid p} \Psi_v, \quad \Psi_p = \sum_{v|p} \Psi_v.$$

(We are exploiting the fact that for m prime to $N\Delta$ the divisors $\hat{\eta}$ and $T^0(m)\hat{\eta}$ have disjoint supports so that we can apply Proposition 5.4.1.2.)

For each prime \wp of F above p , we define an operator⁽³⁹⁾ on \mathcal{S}

$$\mathcal{R}_\wp = U_\wp - 1, \quad \mathcal{R}_p = \prod_{\wp|p} \mathcal{R}_\wp$$

⁽³⁸⁾In our application, we will take $\ell_F = \frac{d}{ds}|_{s=0} \nu^s$ for a character $\nu: \mathcal{G}_F \rightarrow 1 + p\mathbf{Z}_p$.

⁽³⁹⁾This is different from the operator bearing the same name in [41].

Proposition 7.1.2. — *In the space $\overline{\mathcal{S}}$ we have*

$$\Psi_{\text{fin}} \sim \sum_{v \nmid p} \Psi_v + b,$$

where b is a modular form which is killed by l_f ; the sum runs over the finite places of F and the summands are given by:

1. If $v = \wp$ is inert in E , then

$$\Psi_v(m) = \sum_{\substack{n \in N m^{-1} \Delta^{-1} \\ \varepsilon_v((n-1)n)=1 \forall v|\Delta \\ 0 < n < 1}} 2^{\omega_\Delta(n)} r((1-n)m\Delta) r(nm\Delta/N_{\wp})(v(nm/N) + 1) \ell_{F,v}(\pi_v).$$

2. If $v = \wp|\Delta$ is ramified in E , then

$$\Psi_v(m) = \sum_{\substack{n \in N m^{-1} \Delta^{-1} \\ \varepsilon_v((n-1)n)=-1 \\ \varepsilon_w((n-1)n)=1 \forall w \neq v|\Delta \\ 0 < n < 1}} 2^{\omega_\Delta(n)} r((1-n)m\Delta) r(nm\Delta/N)(v(nm) + 1) \ell_v(\pi_v).$$

3. If v is split in E , then

$$\Psi_v(m) = 0.$$

Proof. — For m prime to $N\Delta$ we have $\Psi_{\text{fin}}(m) = \sum_{w \nmid p} \langle \hat{\eta}, T^0(m)\hat{\eta} \rangle_w^{\Delta r}$ (the sum running over all finite places w of E). By Proposition 5.4.1.2, up to the factor $\ell_{F,v}(\pi_v)$ (which equals $\ell_w(\pi_w)$ or its half for each place w of E above v), each term is given by an intersection multiplicity $(\hat{\eta}, T(m)\hat{\eta})_w$, which is computed by Zhang.

When $v(N) \leq 1$ for all v which are not split in E , the result is summarised in [57, Proposition 5.4.8]; in this case, the values obtained there are equivalent to the asserted ones by [57, Proposition 7.1.1 and Proposition 6.4.5], and there is no extra term b . In fact (and with no restriction on N), these values also appear as the local components ${}^C\Phi'_v$ at finite places of a form ${}^C\Phi'$ of level N which is a kernel of the Rankin–Selberg convolution for the central derivative $L'(f_E, 1)$ of the complex L -function.

In general, [58, Lemma 6.4.3] proves that⁽⁴⁰⁾

$$(7.1.2) \quad \frac{\Psi_v}{\ell_{F,v}(\pi_v)} \sim \frac{{}^C\Phi_v^\#}{\log \mathbf{N}(\wp_v)} + {}_v b,$$

where ${}_v b$ is a modular form with zero projection onto the f -eigenspace (see the discussion at the very end of [58]; the forms ${}_v b$ come from intersections at bad places), and ${}^C\Phi^\#$ is a form of level $N\Delta$ which is a kernel for the complex Rankin–Selberg convolution in level $N\Delta$ (in particular, it is modular and $\text{Tr}_\Delta({}^C\Phi^\#) = {}^C\Phi' + b'$ where b' is a modular form of level N which is orthogonal to f). Applying the operator Tr_Δ in (7.1.2) we recover the asserted formula. \square

7.2. Local heights at p / \mathbf{I} . — The following is the key result concerning the local heights at places dividing p . We assume that all primes \wp of F dividing p are split in E .

⁽⁴⁰⁾We are adapting the notation to our case. In [58], the form f is denoted by ϕ , the functions ${}_v b$ are denoted by ${}_v f$.

Proposition 7.2.1. — *The arithmetic function $\mathcal{R}_p^4 \Psi_p$ belongs to $\overline{\mathcal{S}}_N^{\text{ord}} \subset \overline{\mathcal{S}}^{\text{ord}}$, and we have*

$$l_{f_\alpha}(\mathcal{R}_p^4 \Psi_p) = 0.$$

The modularity assertion follows by difference from the modularity of Ψ (hence of $\mathcal{R}_p^4 \Psi$) and the modularity of $\mathcal{R}_p^4 \Psi_{\text{fin}}$ proved in Proposition 8.1.1 below. (In fact one could likely show the stronger result that $\mathcal{R}_\varphi^3 \Psi_v$ is modular for each $v = \varphi | p$, with a method similar to that of [41, Lemme 5.8].)

The proof of the vanishing of the f_α -component will be completed in §7.3 using the results of the rest of this subsection.

We start by fixing for the rest of this section a prime φ of F dividing p . Fix an isomorphism $B_\varphi = B \otimes_F F_\varphi \cong M_2(F_\varphi)$ identifying the local order R_φ with $M_2(\mathcal{O}_{F,\varphi})$, and the field $E \subset B$ with the diagonal matrices in $M_2(F_\varphi)$. Let the divisors η_c be as in (6.2.1), and denote

$$H_s = H[\varphi^s], \quad u_s = u(\varphi^s).$$

Let $\gamma_s \in X(H_s)$ be the CM point of conductor φ^s defined by

$$\gamma_s = \left[\left(x_0, \iota_\varphi \left(\begin{pmatrix} \pi^s & 1 \\ & 1 \end{pmatrix} \right) \right) \right],$$

where $\iota_\varphi: \text{GL}_2(F_\varphi) \rightarrow \widehat{B}^\times$ is the natural inclusion, and π is a uniformiser at φ .

Fix a place w of H above φ ; we still denote by w the induced place on each H_s , and by \mathfrak{p} the prime of E lying below w . Since φ splits in E , by [57, §2.2] the CM points $\gamma_s = [A_s]$ are ordinary, and their canonical submodules with respect to the reduction modulo w are given by $A_s[\mathfrak{p}]$.

Proposition 7.2.2 (Norm relations). — *Let γ_s be the system of CM points defined above.*

1. *Let $m = m_\varphi \varphi^n$ be an ideal of F with m_φ prime to φN . We have*

$$[T(m\varphi^{r+2}) - 2T(m\varphi^{r+1}) + T(m\varphi^r)](\eta) = u_{n+r+2}^{-1} T(m_\varphi) \text{Tr}_{H_{n+r+2}/E}(\gamma_{n+r+2})$$

as divisors on X .

2. *For all $s \geq 1$, we have*

$$T(\varphi)\gamma_s = \text{Tr}_{H_{s+1,w}/H_{s,w}}(\gamma_{s+1}) + \gamma_{s-1}.$$

3. *For all $s \geq 1$, we have*

$$\varphi(\gamma_s) = \gamma_{s-1},$$

where φ is the lift (6.1.1) of Frobenius with respect to the reduction modulo w .

Proof. — By the multiplicativity of Hecke operators it is enough to prove the statement of part 1 for $m_\varphi = 1$. A simple computation based on Proposition 6.3.1 shows that the left-hand side is equal to $\eta_{\varphi^{n+r+2}}$. Since the Galois action of $\text{Gal}(H_{n+r+2}/E)$ is simply transitive on $Y_{\varphi^{n+r+2}}$, the right-hand side is also equal to $\eta_{\varphi^{n+r+2}}$.

For part 2, use the notation $[g]$ to denote $[(x_0, \iota_\varphi(g))]$. Then we have

$$T(\varphi)\gamma_s = \sum_{j \in \mathcal{O}_{F,\varphi}/\varphi} \left[\left(\begin{pmatrix} \pi^s & 1 \\ & 1 \end{pmatrix} \right) \left(\begin{pmatrix} \pi & j \\ & 1 \end{pmatrix} \right) \right] + \left[\left(\begin{pmatrix} \pi^s & 1 \\ & 1 \end{pmatrix} \right) \left(\begin{pmatrix} 1 & \\ & \pi \end{pmatrix} \right) \right].$$

The last term is identified as y_{s-1} after acting by the diagonal matrix $\pi^{-1}\text{id}$ (whose action is trivial on X). On the other hand, by local class field theory and the description of the Galois action on CM points of §6.2, we have

$$\text{Tr}_{H_{s+1,w}/H_{s,w}}(y_{s+1}) = \sum_{j \in \mathcal{O}_{F,\wp}/\wp} \left[\begin{pmatrix} 1 + j\pi^s & \\ & 1 \end{pmatrix} \begin{pmatrix} \pi^{s+1} & 1 \\ & 1 \end{pmatrix} \right],$$

which is the same as the above sum.

For part 3 we switch to the moduli description,⁽⁴¹⁾ so $y_s = [A_s] = [A_{D_s}]$ for an increasing sequence of admissible submodules D_s of level \wp^s (this follows from part 2, together with a variant for $s = 0$ that we omit, and the moduli description of Hecke correspondences). Now D_1 is different from $\text{can}(A) = A[\mathfrak{p}]$ since $[A_{A[\mathfrak{p}}]]$ has conductor 1; and it in fact each D_s does not contain $A[\mathfrak{p}]$, since if it did then $[A_s]$ would be in the support of $T(\wp)^{s-1}[A_{A[\mathfrak{p}}]]$ which is easily seen⁽⁴²⁾ to consist of CM points of conductor dividing \wp^{s-1} . It follows that the point $\varphi([A_s]) = [A_{D_s + \text{can}(A)}] = [A_{D_s + A[\mathfrak{p}]}]$ is in the support of $T(\wp)[A_s]$, but it is not one of the Galois conjugates of y_{s+1} since as just seen it has lower conductor; by part 2, it must then be y_{s-1} . \square

Lemma 7.2.3. — *Let w a place of E dividing \wp , and let $h \in E_w(X)$ be a rational function whose reduction at v is defined and nonzero. Then the arithmetic functions*

$$\mathcal{R}_\wp^3 \langle \widehat{\text{div}}(h), T^0(m)\hat{\eta} \rangle_w^{\text{Ar}}, \quad \mathcal{R}_\wp^3 \langle \text{div}(h), T(m)z \rangle_w$$

belong to the kernel of the \wp -partial ordinary projection e_\wp

Proof. — We show more precisely that

$$v(U_\wp^s \mathcal{R}_\wp^3 \langle \widehat{\text{div}}(h), T^0(m)\hat{\eta} \rangle_w^{\text{Ar}}) \geq v(\mathbf{N}_\wp^s) - C$$

for a uniform constant C , where v is the p -adic valuation. The proof of the same result for the second expression is similar using in addition the fact that $T(m)\xi = \sigma_1(m)\xi$, cf. [41, Lemme 5.4]. We may assume m prime to $\wp N\Delta$. As $\mathcal{R}_\wp^2 r(m) = 0$, Proposition 7.2.2.1 gives

$$U_\wp^s \mathcal{R}_\wp^2 \eta = u_{s+2}^{-1} \text{Tr}_{H_{s+2}/E} y_{s+2}$$

where $y_{s+2} \in Y_{\wp^{s+2}}$; we make a compatible choice of y_s such as the one described above Proposition 7.2.2.

For s large enough the divisor of h is supported away from y_s and its conjugates. Then by Proposition 5.4.1.3 we have

$$\begin{aligned} U_\wp^s \mathcal{R}_\wp^2 \langle \widehat{\text{div}}(h), T^0(m)\hat{\eta} \rangle_w^{\text{Ar}} &= u_{s+2}^{-1} \ell_w(h(T^0(m)y_{s+2})) \\ &= u_{s+2}^{-1} \sum_{w'|w} \ell_w(N_{H_{s+2,w'}/E_w} h(y_{s+2})), \end{aligned}$$

where w' runs over the places of H above w (which are identified with the places of H_{s+2} above w , since H_{s+2}/H is totally ramified above \wp).

⁽⁴¹⁾As usual, after base change to a suitable quadratic extension F' .

⁽⁴²⁾By the following observation: if y is a CM point of conductor c , then the support of $T(m)y$ consists of CM points of conductors dividing cm .

For any $w'|w$ we have

$$\mathcal{R}_\wp \ell_w(N_{H_{s+2,w'}/E_w} b(y_{s+2})) = \ell_w \circ N_{H_{w'}/E_w}(N_{H_{s+3,w'}/H_{w'}} b(y_{s+3})/N_{H_{s+2,w'}/H_{w'}} b(y_{s+2})).$$

Suppose that (for s large enough)

$$(7.2.1) \quad \text{the } w'\text{-adic valuation of } N_{H_{s,w'}/H_{w'}}(b(y_s)) \text{ is independent of } s.$$

Then each w' -summand in the expression of interest is the product of μ_{s+2}^{-1} (which is eventually constant in s) and the p -adic logarithm of a unit which is a norm from an extension of E_w whose ramification degree is a constant multiple of \mathbf{N}_\wp^s ; hence its p -adic valuation is also at least a constant multiple of the valuation of \mathbf{N}_\wp^s , which proves the Lemma.

It remains to prove (7.2.1). We have

$$(7.2.2) \quad w'(N_{H_{s,w'}/H_{w'}}(b(y_s))) = [H_{s,w'} : H_{w'}](\underline{(b)}, \underline{y_s}),$$

where the right-hand side denotes the intersection of the Zariski closures in the integral model. Now as in [41, Lemme 5.5], if π_s denotes a uniformiser of $H_{s,w'}$ we can show that we have

$$(7.2.3) \quad \underline{y_s} \equiv \underline{y_{s-1}} \pmod{\pi_s}, \quad \underline{y_s} \not\equiv \underline{y_{s-1}} \pmod{\pi_s^2} :$$

by Honda–Tate theory, it suffices to check this on the \wp -divisible groups $\mathcal{A}_{\underline{y_s}}[\wp^\infty]$ or equivalently on $\mathcal{G}_s = \mathcal{G}_{\underline{y_s}}$, or on \mathcal{G}_s^1 (with the notation of §6.1). That $\underline{y_s}$ is ordinary means that \mathcal{G}_s^1 is an extension of $F_\wp/\mathcal{O}_{F_\wp}$ by the Lubin–Tate formal module Σ_1 . Now \mathcal{G}_s^1 is defined precisely over the ring of integers of $H_{s,w'}$, and so it is a quasi-canonical lifting of level s of its reduction, as studied by Gross [19, §6]. Then \mathcal{G}_s^1 is congruent to the canonical lifting modulo π_s but not modulo π_s^2 , whereas \mathcal{G}_{s-1}^1 is congruent to the canonical lifting modulo $\pi_{s-1} = \pi_s^{\mathbf{N}_\wp}$; this implies (7.2.3). Then for each irreducible component \underline{a} in the support of $\underline{(b)}$, the sequence $[H_{s,w'} : H_{w'}](\underline{a}, \underline{y_s})$ stabilises to either 0 or 1, so that the expression (7.2.2) is indeed eventually constant. \square

Lemma 7.2.4. — *For each divisor $D \in \text{Div}^0(X)(E_w)$ (respectively, $\widehat{D} \in \text{Div}^{\text{Ar}}(X)$), the element of $\overline{\mathcal{F}}^{\text{ord}}$ given by*

$$m \mapsto \mathcal{R}_\wp^3 \langle D, T(m)z \rangle_w \quad (\text{respectively } m \mapsto \mathcal{R}_\wp^3 \langle \widehat{D}, T^0(m)\hat{\eta} \rangle_w^{\text{Ar}})$$

is well-defined independently of the choice of D in its class $[D]$ (of \widehat{D} in its class $[\widehat{D}]$); it will be denoted by

$$\mathcal{R}_\wp^3 \langle [D], T(m)z \rangle_w \quad (\text{respectively } \mathcal{R}_\wp^3 \langle [\widehat{D}], T^0(m)\hat{\eta} \rangle_w^{\text{Ar}}).$$

If $\widehat{D} = D$, then the two elements coincide as elements of $\overline{\mathcal{F}}^{\text{ord}}$; moreover, for the arithmetic function $\Psi_w \in \overline{\mathcal{F}}$ with $\Psi_w(m) = \langle \hat{\eta}, T^0(m)\hat{\eta} \rangle_w^{\text{Ar}}$, we have

$$(7.2.4) \quad \mathcal{R}_\wp^3 \Psi_w \sim \mathcal{R}_\wp^3 \langle [z], T(m)z \rangle_w$$

in $\overline{\mathcal{F}}^{\text{ord}}$.

Proof. — The first part follows from Lemma 7.2.3. For the second part we may argue as in the proof of Lemma 7.1.1: for example, in $\overline{\mathcal{F}}$ we have

$$\mathcal{R}_\wp^3 \langle \hat{\eta}, T^0(m)\hat{\eta} \rangle_w^{\text{Ar}} \sim \mathcal{R}_\wp^3 \langle \hat{\eta} + \widehat{\text{div}}(b), T^0(m)\hat{\eta} \rangle_w^{\text{Ar}}$$

$$\sim \mathcal{R}_\varphi^3 \langle \widehat{z} + \widehat{\operatorname{div}}(b), T(m)\widehat{z} \rangle_w^{\text{Ar}} = \mathcal{R}_\varphi^3 \langle z + \operatorname{div}(b), T(m)z \rangle_w.$$

□

7.3. Local heights at p / \mathbf{II} . — Here we prove the vanishing statement for p -adic local symbols asserted in Proposition 7.2.1.

Let $e_f \in \mathbf{T}_{Np} \otimes M_f$ be the maximal idempotent satisfying $T(m) \circ e_f = a(f, m)e_f$ for all m prime to Np ; ⁽⁴³⁾ viewed as an endomorphism of $S_{N\prod_{\varphi|p} \varphi}$, it is the projector onto the subspace generated by f and $[\varphi]f$ for all the primes φ of F dividing p . With $z_f = e_f[z]$, we have by (7.2.4)

$$e_f e \mathcal{R}_p^4 \Psi_p = \mathcal{R}_p^4 \langle z_f, T(m)z \rangle_p$$

in $\overline{\mathcal{S}}_N^{\text{ord}}$, where the left-hand side makes sense by the modularity part of Proposition 7.2.1 proved in §7.2 and the right-hand side makes sense by Lemma 7.2.4. We also denote, for w a place of E above the F -prime $\varphi|p$, and $i = 3$ or 4 ,

$$(7.3.1) \quad e_f e \mathcal{R}_\varphi^i \Psi_w := \mathcal{R}_\varphi^i \langle z_f, T(m)z \rangle_w$$

where the right-hand side makes sense as an element of $\overline{\mathcal{S}}^{\text{ord}}$ by Lemma 7.2.4. (As we have not shown that $\mathcal{R}_\varphi^3 \Psi_w$ is modular, the left-hand side is not otherwise defined.) Then by definition we have

$$(7.3.2) \quad e_f e \mathcal{R}_p^4 \Psi_p = \sum_{w|p} e_f e \mathcal{R}_p^4 \Psi_w.$$

Recall that we want to show that

$$l_{f_x}(\mathcal{R}_p^4 \Psi_p) = 0.$$

Since $l_{f_x} = l_{f_x} \circ e_f = l_{f_x} \circ e_f \circ e$, by (7.3.2) the desired result is implied by the following Lemma for all $\varphi|p$.

Lemma 7.3.1. — *Suppose that f is ordinary at φ . For each place w of E above $\varphi|p$, the element $e_f \mathcal{R}_\varphi^3 \Psi_w$ is zero in $\overline{\mathcal{S}}^{\text{ord}}$.*

Proof. — The ordinarity assumption and Lemma 5.1.1 (cf. [41, Exemple 4.12]) imply that z_f is “almost” a universal norm in the totally ramified \mathbf{Z}_p -extension $E_{w,\infty}^\ell$ of E_w : that is, after perhaps replacing z_f by an integer multiple, for each layer $E_{w,n}^\ell$ we have

$$z_f = \operatorname{Tr}_n(z_n)$$

for some $z_n \in e_f J(X)(E_{v,n}^\ell)$, where $\operatorname{Tr}_n = \operatorname{Tr}_{E_{w,n}^\ell/E_w}$. Then we have

$$e_f \mathcal{R}_\varphi^3 \Psi_w(m) = \mathcal{R}_\varphi^3 \langle \operatorname{Tr}_n(z_n), T(m)z \rangle_w = \mathcal{R}_\varphi^3 \langle z_n, T(m)z \rangle_{w,n}.$$

where $\langle \cdot, \cdot \rangle_{w,n}$ is the local height pairing on $\operatorname{Div}^0(X)(E_{w,n}^\ell)$ associated to the logarithm $\ell_{n,v} = \ell_w \circ N_{E_{w,n}^\ell/E_w}$. By Proposition 5.1.2.5-6, the right-hand side above has image in $c^{-1} \operatorname{Im}(\ell_n) \subset \mathbf{Z}_p$ for a uniform nonzero constant $c \in \mathbf{Z}_p$. As the extension $E_{w,n}^\ell/E_w$ has ramification degree p^n , we

⁽⁴³⁾Recall that M_f is the number field generated by the Fourier coefficients $a(m, f)$.

have for some nonzero $c' \in \mathbf{Z}_p$

$$e_f \mathcal{R}_\varphi^3 \Psi_v(m) \in c'^{-1} \text{Im}(\ell_n) \subset c'^{-1} p^n \mathbf{Z}_p$$

for all n ; therefore $e_f \mathcal{R}_\varphi^3 \Psi_w = 0$. □

PART III MAIN THEOREM AND CONSEQUENCES

8. Proof of the main theorem

In this section we prove Theorem B. We resume the notation $\langle \cdot, \cdot \rangle_{\mathcal{W}, \alpha}$ for our height pairing (see §7 for the dependence on α). First, notice that when \mathcal{W} is anticyclotomic, both sides of the formula are zero: indeed, $L_{p, \alpha}(f_E)(\mathcal{W}^s)$ vanishes identically by the functional equation (4.2.3), and

$$\langle z_f, z_f \rangle_{\mathcal{W}, \alpha} = \langle z_f^c, z_f^c \rangle_{\mathcal{W}^c, \alpha} = -\langle z_f, z_f \rangle_{\mathcal{W}, \alpha}$$

in view of the following lemma (recall that $\tau(f) = \pm 1$.)

Lemma 8.0.2. — *The action of complex conjugation on z_f is given by*

$$z_f^c = (-1)^g \tau(f) z_f.$$

Therefore it will suffice to prove the formula when

$$\mathcal{W} = \mathcal{W}^+ = \nu \circ \mathfrak{N}$$

for some Hecke character ν of F valued in $1 + p\mathbf{Z}_p$.

8.1. Basic case. — First we prove the formula when $\Delta_{E/F}$ is totally odd and each prime φ of F dividing p splits in E .

Let $\Psi_{\mathcal{W}, \alpha} \in \overline{\mathcal{S}}_N$ denote the modular form with coefficients $([z], T(m)[z])_{\mathcal{W}, \alpha}$, where $\langle \cdot, \cdot \rangle_{\mathcal{W}, \alpha}$ is the height pairing on $J(X)(E)$ associated to the p -adic logarithm $\ell_F \circ \mathfrak{N}$, with

$$\ell_F = \frac{d}{ds} \nu^s |_{s=0} : F^\times \backslash F_{\mathbb{A}^\infty}^\times \rightarrow \mathbf{Q}_p.$$

Recall that l_{f_α} is a continuous functional, so that it commutes with limits and

$$L'_{p, \mathcal{W}}(f_E)(1) = l_{f_\alpha} \left(\frac{d}{ds} \Phi(\mathcal{W}^s) |_{s=0} \right) = l_{f_\alpha}(\Phi'_{\mathcal{W}}).$$

We compare the Fourier coefficients of $\Phi'_{\mathcal{W}}$ and $\Psi_{\mathcal{W}, \alpha} = \Psi_{\mathcal{W}, \text{fin}} + \Psi_{\mathcal{W}, \alpha, p}$.

Proposition 8.1.1. — *Suppose that all of the prime ideals φ of F dividing p are principal. Then we have*

$$\left(\prod_{\varphi|p} U_\varphi^4 - U_\varphi^2 \right) \Phi'_{\mathcal{W}} \sim \left(\prod_{\varphi|p} (U_\varphi - 1)^4 \right) \Psi_{\mathcal{W}, \alpha, \text{fin}}$$

in the quotient space $\overline{\mathcal{F}}_N / \text{Ker}(l_{f_\alpha})$.

Proof. — The coefficients of $\Psi_{\text{fin}} = \Psi_{\mathcal{W}, \alpha, \text{fin}}$ are computed in Proposition 7.1.2. To lighten the notation, we write the explicit expression for $\Psi(m) = \sum_{v \text{ non-split}} \Psi_v(m)$ as

$$\Psi_v(m) = \sum_{\substack{n \in S_v([m]) \\ v_\varphi(nm) \geq 0 \forall \varphi|p}} c_v([nm]) r((1-n)m\Delta) r(nm\Delta/N \varphi^{\epsilon(v)}),$$

where the value $c_v([nm])$ only depends on the prime-to- p part of the fractional ideal nm , and the set $S_v([m])$ only depends on v and the prime-to- p part of m ; here $\epsilon(v) = 1$ if v is inert and $\epsilon(v) = 0$ if v is ramified.

The coefficients of Φ' are computed in Proposition 4.5.3. They look “almost” the same, in that, up to the modular form h of Proposition 7.1.2, which is in $\text{Ker}(l_{f_\alpha})$, we have, when m is divisible by every $\varphi|p$,

$$\Phi'_{\mathcal{W}}(m) = \sum_{v \text{ non-split}} \Psi_v^{[P]}(m),$$

where for a product P of some of the primes $\varphi|p$, we denote

$$\Psi_v^{[P]}(m) = \sum_{\substack{n \in S_v([m]) \\ v_\varphi(nm) \geq 0 \forall \varphi|p \\ v_\varphi(nm) = 0 \forall \varphi|P}} c_v([nm]) r((1-n)m\Delta) r(nm\Delta/N).$$

Then it is enough to show that for each $v \nmid p$, each $\varphi|p$, and each $\varphi \nmid P$ with P as above, we have

$$(U_\varphi^4 - U_\varphi^2) \Psi_v^{[P, \varphi]} = (U_\varphi - 1)^4 \Psi_v^{[P]}.$$

For the sake of notation we write the computation when v is ramified in E and $P = \prod_{\varphi' \neq \varphi} \varphi'$ (for more general P one just needs more notation to keep track of $v_{\varphi'}(nm)$ for the primes $\varphi' \neq \varphi$).

The right-hand side equals

$$(8.1.1) \quad \sum_{i=0}^4 (-1)^i \binom{4}{i} \sum_{\substack{n_i \in S_v([m]) \\ v_{\varphi'}(n_i m) = 0 \forall \varphi' \neq \varphi, \varphi|p \\ v_\varphi(n_i m \varphi^i) \geq 0}} c_v([n_i m]) r((1-n_i)m\varphi^i\Delta) r(n_i m \varphi^i \Delta/N).$$

From the relation $r(m_0 \varphi^t) = (t+1)r(m_0)$, valid for $\varphi \nmid m_0$, we deduce the relations

$$\begin{aligned} 2r(m) &= r(m\varphi) + r(m\varphi^{-1}), \\ 2r(m) &= r(m\varphi^2) + r(m\varphi^{-2}) \quad \text{if } \varphi|m, \\ 2r(m) &= r(m\varphi^2) - r(m) \quad \text{if } \varphi \nmid m, \end{aligned}$$

where we recall that $r(m) = 0$ if m is not an integral ideal. Then we can pick a totally positive generator in F for the ideal φ , which abusing notation we will still denote by φ , and make the substitution $n_i = \varphi^{t-i} n_0$ with $\varphi^t \nmid n_i m \varphi^i$ to write (8.1.1) as

$$\sum_{t \geq 0} \sum_{\substack{n_0 \in S_v(m) \\ v_{\varphi'}(n_0 m) = 0 \forall \varphi|p}} c_v([n_0 m]) r((n_0 m)^{(\varphi)}) (t+1) A_t$$

where we recall that for an ideal m we denote $m^{(\wp)} = m\wp^{-v_\wp(m)}$, and

$$\begin{aligned} A_t &= r(m\Delta\wp^4(1 - n_0\wp^{t-4})[t + 1 - 2t + 2(t - 1)] \\ &\quad + r(m\Delta\wp^2(1 - n_0\wp^{t-2}) \left[-2(t + 2) + \begin{cases} 4(t + 1) - 2t & \text{if } t \geq 1 \\ 3 & \text{if } t = 0 \end{cases} \right] \\ &\quad + r(m\Delta(1 - n_0\wp^t))[t + 3 - 2(t + 2) + t + 1]. \end{aligned}$$

The three expressions in square brackets vanish when $t > 0$ and yield, respectively, 1, 1, and 0 when $t = 0$. Substituting back $n_4 = \wp^{t-4}n_0$ in the first line and $n_2 = \wp^{t-2}n_0$ in the second line, we deduce that (8.1.1) equals

$$(U_\wp^4 - U_\wp^2)\Psi_v^{[P\wp]}$$

as desired.⁽⁴⁴⁾ □

Combining this Proposition with Proposition 7.2.1 which says

$$l_{f_z} \left(\prod_{\wp|p} (U_\wp - 1)^4 \Psi_{\mathcal{W}, \alpha, p} \right) = 0,$$

we find for $\mathcal{W} = \nu \circ \mathfrak{N}$:

$$\begin{aligned} D_F^2 \prod_{\wp} (\alpha_\wp^4 - \alpha_\wp^2) L'_{p, \alpha, \mathcal{W}}(f_E, 1) &= \prod_{\wp} (\alpha_\wp^4 - \alpha_\wp^2) \left(1 - \frac{1}{\alpha_\wp^2} \right) \left(1 - \frac{\mathbf{N}_\wp}{\alpha_\wp^2} \right) l_{f_z}(\Phi'_{\mathcal{W}}) \\ &= \prod_{\wp} (\alpha_\wp - 1)^4 \left(1 - \frac{1}{\alpha_\wp^2} \right) \left(1 - \frac{\mathbf{N}_\wp}{\alpha_\wp^2} \right) l_{f_z}(\Psi_{\mathcal{W}, \alpha}) \\ &= \prod_{\wp} (\alpha_\wp - 1)^4 \left(1 - \frac{1}{\alpha_\wp^2} \right) \langle z_f, z_f \rangle_{\mathcal{W}, \alpha}. \end{aligned}$$

Here, besides the definition of $L_{p, \alpha}(f_E)$ (Definition 4.2.1) we have used various properties of the functional l_{f_z} from Lemma 1.6.1 and the observation that the projection onto the f -component of the modular form $\Psi_{\mathcal{W}} \in S_2(K_0(N), \mathbf{Q}_p)$ is $\mathbf{1}_f(\Psi_{\mathcal{W}}) = \langle z_f, z_f \rangle_{\mathcal{W}}$.

This completes the proof of Theorem B when $(\Delta_{E/F}, 2) = 1$ and all primes $\wp|p$ split in E .

8.2. Reduction to the basic case. — The general case, where E is only assumed to satisfy $(\Delta_{E/F}, Np) = 1$, can be reduced to the previous one under the assumption

$$L'_{p, \mathcal{W}}(f_E, 1) \neq 0$$

by the following argument due to Kobayashi [30, proof of Theorem 5.9] using the complex Gross–Zagier formula (which is known with no restrictions on Δ) and the factorisation $L_{p, \alpha}(f_E, \chi \circ \mathfrak{N}) \sim L_{p, \alpha}(f, \chi)L_{p, \alpha}(f_\varepsilon, \chi)$.

⁽⁴⁴⁾Cf. [41, Proof of Proposition 3.20].

By the factorisation the orders of vanishing at the central point of the factors of $L_{p,\alpha}(f_E, \nu^s \circ \mathfrak{N})$ will be one (say for $L_{p,\alpha}(f)$) and zero (say for $L_{p,\alpha}(f_\varepsilon)$). Then, by the first part of Theorem C,⁽⁴⁵⁾ the orders of vanishing of $L(f, s)$ and $L(f_\varepsilon, s)$ at $s = 1$ will also be one and zero. Moreover the Heegner point $z_{f,E'}$ attached to f and any E' also satisfying $L(f_{\varepsilon_{E'/F}}, 1) \neq 0$ is non-torsion, and in fact its trace $z_{f,F} = \text{Tr}_{E'/F}(z_{f,E'})$ is non-torsion and $z_{f,E'}$ is up to torsion a multiple of $z_{f,F}$ in $J(X)(E') \otimes \overline{\mathbf{Q}}$. Therefore, by the complex and p -adic Gross–Zagier formulas for a suitable E' satisfying the assumptions of §8.1 and $L(f_{\varepsilon_{E'/F}}, 1) \neq 0$, we have

$$L'_{p,\nu}(f, 1) = \prod_{\wp|p} \left(1 - \frac{1}{\alpha_\wp}\right)^2 \frac{L'(f, 1)}{\Omega_f^+ \langle z_{f,F}, z_{f,F} \rangle} \langle z_{f,F}, z_{f,F} \rangle_\nu$$

where $\langle \cdot, \cdot \rangle_\nu$ is the p -adic height pairing on $J(X)(F)$ attached to ν , and $\langle \cdot, \cdot \rangle$ is the Néron–Tate height (the ratio appearing above belongs to M_f^\times by the Gross–Zagier formula). This allows us to conclude

$$\begin{aligned} L'_{p,\alpha,\mathcal{W}}(f_E, 1) &= \frac{\Omega_f^+ \Omega_{f_\varepsilon}^+}{D_E^{-1/2} \Omega_f} L'_{p,\alpha,\nu}(f, 1) L_{p,\alpha}(f_\varepsilon, 1) \\ &= D_E^{1/2} \prod_{\wp|p} \left(1 - \frac{1}{\alpha_\wp}\right)^2 \left(1 - \frac{\varepsilon(\wp)}{\alpha_\wp}\right)^2 \frac{L'(f, 1) L(f_\varepsilon, 1)}{\Omega_f \langle z_{f,F}, z_{f,F} \rangle} \langle z_{f,F}, z_{f,F} \rangle_\nu \\ &= D_F^{-2} \prod_{\wp|p} \left(1 - \frac{1}{\alpha_\wp}\right)^2 \left(1 - \frac{\varepsilon(\wp)}{\alpha_\wp}\right)^2 \frac{\langle z_{f,E}, z_{f,E} \rangle}{\langle z_{f,F}, z_{f,F} \rangle} \langle z_{f,F}, z_{f,F} \rangle_\nu \\ &= D_F^{-2} \prod_{\wp|p} \left(1 - \frac{1}{\alpha_\wp}\right)^2 \left(1 - \frac{\varepsilon(\wp)}{\alpha_\wp}\right)^2 \langle z_{f,E}, z_{f,E} \rangle_{\mathcal{W}}. \end{aligned}$$

Remark 8.2.1. — It is natural to conjecture that when $L'_{p,\mathcal{W}}(f_E, 1) = 0$ we should have $\langle z_{f,F}, z_{f,F} \rangle_{\mathcal{W}} = 0$. However in this case the above argument fails because, without knowledge of the nontriviality of the p -adic height pairing, the vanishing of $L_{p,\alpha}(f_E, \mathcal{W}^s)$ to order ≥ 2 does not imply a similar high-order vanishing for $L(f_E, s)$.

9. Periods and the Birch and Swinnerton-Dyer conjecture

As seen in the Introduction, the application of our result to the Birch and Swinnerton-Dyer formula rests on a conjectural relation among the periods of f and the associated abelian variety A . Here we would like to briefly elaborate on this conjecture and its arithmetic consequences. We retain the notation of the Introduction, and set $M = M_f$ and $\dim A = [M : \mathbf{Q}] = d$.

9.1. Real periods. — The conjecture on periods stated in the Introduction can be refined to a conjecture on rationality rather than algebraicity. First we need to precisely define the automorphic periods $\Omega_{f,\sigma}^+$, for $\sigma \in \text{Hom}(M, \mathbf{C})$; they are naturally defined as elements of $\mathbf{C}^\times / M^\times$ (see [44] for a modern exposition): one can choose them “covariantly” in order to have $\prod_\sigma \Omega_{f,\sigma}^+$ defined up

⁽⁴⁵⁾Which can be proved by using the p -adic Gross–Zagier formula attached to a field E' satisfying the assumptions of §8.1.

to \mathbf{Q}^\times , or define directly the product as follows. Let $\mathcal{H}_N = Z(\mathbf{A}) \backslash \mathbf{GL}_2(\mathbf{A}) / K_0(N) K_\infty$ be the open Hilbert modular variety of level N . Then the perfect pairing of \mathbf{Q} -vector spaces

$$(9.1.1) \quad H_g(\mathcal{H}_N, \mathbf{Q})^+ \times S_2(K_0(N), \mathbf{Q}) \rightarrow \mathbf{C}$$

(where “+” denotes the intersection of the +1-eigenspaces for the complex conjugations) decomposes under the diagonal action of \mathbf{T}_N into \mathbf{Q} -rational blocks parametrised by the Galois-conjugacy classes of eigenforms. Then

$$\prod_{\sigma} \Omega_{f^\sigma}^+ \in \mathbf{C}^\times / \mathbf{Q}^\times$$

is $(2\pi i)^{dg}$ times the discriminant of the pairing on the rational block corresponding to $\{f^\sigma\}_\sigma$. (The individual $\Omega_{f^\sigma}^+ \in \mathbf{C}^\times / M^\times$ are defined as the discriminants of (9.1.1) on $\overline{\mathbf{Q}}$ -rational \mathbf{T}_N -eigenblocks.)

Conjecture 9.1.1. — *We have*

$$\Omega_A \sim \prod_{\sigma} \Omega_{f^\sigma}^+$$

in $\mathbf{C}^\times / \mathbf{Q}^\times$.

The conjecture, which originates from the work of Shimura, is also made by Yoshida [53] up to algebraicity. When A has complex multiplication, it has been proved by Blasius [6]. It is also known when $F = \mathbf{Q}$; before discussing that, let us translate it into a language closer to conjectures of Shimura.

For each $\tau: F \rightarrow \mathbf{R}$, let $f_{B(\tau)}$ be the Jacquet-Langlands transfer of f to a rational form on the quaternion algebra $B(\tau)/F$ defined in the Introduction (recall that $B(\tau)$ is ramified at all infinite places except τ), and let X be our Shimura curve. Then A is (up to isogeny) a quotient ϕ of $J(X)$, and for each embedding τ we can write

$$\phi^* \omega_A = c_\tau \bigwedge_{\sigma} 2\pi i f_{B(\tau)}^\sigma(z) dz$$

as forms in $H^0(J(X)(\mathbf{C}_\tau), \Omega^d)$, for some $c_\tau \in F^\times$ (since both are generators of a rank one F -vector space). Then we have

$$\int_{A(\mathbf{R}_\tau)} |\omega_A|_\tau \sim \prod_{\sigma} \Omega_{f_{B(\tau)}^\sigma}^+ \text{ in } \mathbf{C}^\times / F^\times,$$

where $\Omega_{f_{B(\tau)}^\sigma}^+$ is $2\pi i$ times the discriminant of the $f_{B(\tau)}^\sigma$ -part of the analogue of the pairing (9.1.1) on $X(\mathbf{C}_\tau)$. When choices are made covariantly in τ , we then get $\Omega_A \sim \prod_{\sigma, \tau} \Omega_{f_{B(\tau)}^\sigma}^+$ in $\mathbf{C}^\times / \mathbf{Q}^\times$.

Our conjecture, decomposed into its (σ, τ) -constituents, can then be rewritten as

$$(9.1.2) \quad \Omega_f^+ \sim \prod_{\tau} \Omega_{f_{B(\tau)}^\sigma}^+ \text{ in } \mathbf{C}^\times / (MF)^\times.$$

In this form, this is a stronger version of Shimura’s conjecture [49] on the factorisation of periods of Hilbert modular forms up to algebraic factors in terms of P -invariants. The reader is referred to [53] for a discussion of this point.

Notice that (9.1.2) is nontrivial even when $F = \mathbf{Q}$: it asserts that the periods of the transfers of f to any indefinite quaternion algebra have the same transcendental (or irrational) parts. However, in this case the conjecture is known by the work of Shimura [48] (for the algebraicity) and Prasanna [43] (for the rationality).

For general F , Shimura’s conjecture on P -invariants is largely proved by Yoshida [54] under an assumption of non-vanishing of certain L -values.

Remark 9.1.2. — It is clear that our conjecture implies that the Birch and Swinnerton-Dyer conjectural formula is true up to a nonzero rational factor when A has analytic M -rank zero. By the complex (respectively, the p -adic) Gross–Zagier formula, the conjecture for f also implies the complex (respectively, the p -adic) Birch and Swinnerton-Dyer formulas up to a rational factor when A has (p -adic) analytic M -rank one.

9.2. Quadratic periods. — We can formulate a conjecture analogous to Conjecture 9.1.1 for the periods of the base-changed abelian variety $A_E = A \times_{\text{Spec } F} \text{Spec } E$.

Conjecture 9.2.1. — *We have*

$$\Omega_{A_E} \sim \prod_{\sigma} \Omega_{f^{\sigma}}$$

in $\mathbf{C}^{\times}/\mathbf{Q}^{\times}$.

Here the period of A_E is

$$\Omega_{A_E} = \prod_{\tau: E \rightarrow \mathbf{C}} \int_{A(\mathbf{C}_{\tau})} |\omega_{A_E}|_{\tau},$$

where for a differential form $\omega = h(z)dz_1 \wedge \cdots \wedge dz_k$ we have $|\omega|_{\tau} = |h(z)|_{\tau}^2 dz_1 \wedge d\bar{z}_1 \wedge \cdots \wedge dz_k \wedge d\bar{z}_k$.

As above, this conjecture can be “decomposed” into

$$(9.2.1) \quad \Omega_f \sim \prod_{\tau} \Omega_{f_{B(\tau)}} \text{ in } \mathbf{C}^{\times}/(MF)^{\times}.$$

where $\Omega_{f_{B(\tau)}}$ is π^2 times the Petersson inner product of $f_{B(\tau)}$. This is essentially Shimura’s conjecture on \mathbf{Q} -invariants (see [49]). Up to algebraicity it has been proved by Harris [21] under a local condition (a new proof of the same result should appear in forthcoming work of Ichino–Prasanna, yielding rationality and removing the local assumption). Since (9.2.1) is implied by (9.1.2) for f and f_{ε} , Harris’s result can be seen as evidence for the conjecture on real periods.

We take the opportunity to record an immediate consequence of the conjecture on quadratic periods and the Gross–Zagier formulas.

Theorem 9.2.2. — *If A_E has complex (respectively, p -adic) analytic M -rank ≤ 1 , then the complex (respectively, the p -adic) Birch and Swinnerton-Dyer formula for A_E is true up to a nonzero algebraic factor.*

References

- [1] F. Andreatta and E. Z. Goren, *Hilbert modular forms: mod p and p -adic aspects*, Mem. Amer. Math. Soc. 173 (2005), no. 819, vi+100, DOI 10.1090/memo/0819. MR2110225 (2006f:11049) ↑13

- [2] Daniel Bertrand, *Propriétés arithmétiques de fonctions thêta à plusieurs variables*, Number theory, Noordwijkerhout 1983 (Noordwijkerhout, 1983), Lecture Notes in Math., vol. 1068, Springer, Berlin, 1984, pp. 17–22, DOI 10.1007/BFb0099438 (French). MR756080 ↑8
- [3] Amnon Besser, *The p -adic height pairings of Coleman-Gross and of Nekovář*, Number theory, CRM Proc. Lecture Notes, vol. 36, Amer. Math. Soc., Providence, RI, 2004, pp. 13–25. MR2076563 (2005f:11130) ↑45
- [4] ———, *p -adic Arakelov theory*, J. Number Theory 111 (2005), no. 2, 318–371, DOI 10.1016/j.jnt.2004.11.010. MR2130113 (2006j:14029) ↑9, 42, 43, 44, 45
- [5] Daniel Bump, Solomon Friedberg, and Jeffrey Hoffstein, *Nonvanishing theorems for L -functions of modular forms and their derivatives*, Invent. Math. 102 (1990), no. 3, 543–618, DOI 10.1007/BF01233440. MR1074487 (92a:11058) ↑8
- [6] Don Blasius, *On the critical values of Hecke L -series*, Ann. of Math. (2) 124 (1986), no. 1, 23–63, DOI 10.2307/1971386. MR847951 (88i:11035) ↑6, 61
- [7] Daniel Bump, *Automorphic forms and representations*, Cambridge Studies in Advanced Mathematics, vol. 55, Cambridge University Press, Cambridge, 1997. MR1431508 (97k:11080) ↑19
- [8] Colin J. Bushnell and Guy Henniart, *The local Langlands conjecture for $GL(2)$* , Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 335, Springer-Verlag, Berlin, 2006. MR2234120 (2007m:22013) ↑19
- [9] Henri Carayol, *Sur la mauvaise réduction des courbes de Shimura*, Compositio Math. 59 (1986), no. 2, 151–230 (French). MR860139 (88a:11058) ↑45, 46
- [10] William Casselman, *On some results of Atkin and Lehner*, Math. Ann. 201 (1973), 301–314. MR0337789 (49 #2558) ↑15
- [11] Robert F. Coleman and Benedict H. Gross, *p -adic heights on curves*, Algebraic number theory, Adv. Stud. Pure Math., vol. 17, Academic Press, Boston, MA, 1989, pp. 73–81. MR1097610 (92d:11057) ↑45
- [12] Robert F. Coleman, *p -adic Banach spaces and families of modular forms*, Invent. Math. 127 (1997), no. 3, 417–479, DOI 10.1007/s002220050127. MR1431135 (98b:11047) ↑
- [13] Robert F. Coleman and Bas Edixhoven, *On the semi-simplicity of the U_p -operator on modular forms*, Math. Ann. 310 (1998), no. 1, 119–127, DOI 10.1007/s002080050140. MR1600034 (99b:11043) ↑
- [14] Andrzej Dabrowski, *p -adic L -functions of Hilbert modular forms*, Ann. Inst. Fourier (Grenoble) 44 (1994), no. 4, 1025–1041 (English, with English and French summaries). MR1306548 (96b:11065) ↑
- [15] Mladen Dimitrov, *Automorphic symbols, p -adic L -functions and ordinary cohomology of Hilbert modular varieties*, Amer. J. Math. 135 (2013), 1–39. ↑4, 36
- [16] Pierre Deligne and Kenneth A. Ribet, *Values of abelian L -functions at negative integers over totally real fields*, Invent. Math. 59 (1980), no. 3, 227–286, DOI 10.1007/BF01453237. MR579702 (81m:12019) ↑30
- [17] Gerd Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, Invent. Math. 73 (1983), no. 3, 349–366, DOI 10.1007/BF01388432 (German). MR718935 (85g:11026a) ↑5
- [18] Jean-Marc Fontaine, *Sur certains types de représentations p -adiques du groupe de Galois d’un corps local; construction d’un anneau de Barsotti-Tate*, Ann. of Math. (2) 115 (1982), no. 3, 529–577, DOI 10.2307/2007012 (French). MR657238 (84d:14010) ↑39
- [19] Benedict H. Gross, *On canonical and quasicanonical liftings*, Invent. Math. 84 (1986), no. 2, 321–326, DOI 10.1007/BF01388810. MR833193 (87g:14051) ↑55
- [20] Benedict H. Gross and Don B. Zagier, *Heegner points and derivatives of L -series*, Invent. Math. 84 (1986), no. 2, 225–320, DOI 10.1007/BF01388809. MR833192 (87j:11057) ↑4, 7, 10, 30
- [21] Michael Harris, *L -functions of 2×2 unitary groups and factorization of periods of Hilbert modular forms*, J. Amer. Math. Soc. 6 (1993), no. 3, 637–719, DOI 10.2307/2152780. MR1186960 (93m:11043) ↑62
- [22] Haruzo Hida, *A p -adic measure attached to the zeta functions associated with two elliptic modular forms. I*, Invent. Math. 79 (1985), no. 1, 159–195, DOI 10.1007/BF01388661. MR774534 (86m:11097) ↑18
- [23] ———, *On p -adic L -functions of $GL(2) \times GL(2)$ over totally real fields*, Ann. Inst. Fourier (Grenoble) 41 (1991), no. 2, 311–391 (English, with French summary). MR1137290 (93b:11052) ↑4, 16, 18
- [24] H. Hida and J. Tilouine, *Anti-cyclotomic Katz p -adic L -functions and congruence modules*, Ann. Sci. École Norm. Sup. (4) 26 (1993), no. 2, 189–259. MR1209708 (93m:11044) ↑23
- [25] Benjamin Howard, *Iwasawa theory of Heegner points on abelian varieties of GL_2 type*, Duke Math. J. 124 (2004), no. 1, 1–45, DOI 10.1215/S0012-7094-04-12411-X. MR2072210 (2005f:11117) ↑9
- [26] Ming-Lun Hsieh, *Eisenstein congruence on unitary groups and Iwasawa main conjecture for CM fields*, preprint available at <http://www.math.ntu.edu.tw/mlhsieh/research.html>. ↑9
- [27] Adrian Iovita, *Formal sections and de Rham cohomology of semistable abelian varieties*. part B, Proceedings of the Conference on p -adic Aspects of the Theory of Automorphic Representations (Jerusalem, 1998), 2000, pp. 429–447, DOI 10.1007/BF02834846. MR1809629 (2002g:14026) ↑39

- [28] Hervé Jacquet, *Automorphic forms on $GL(2)$. Part II*, Lecture Notes in Mathematics, Vol. 278, Springer-Verlag, Berlin, 1972. MR0562503 (58 #27778) ↑31
- [29] Nicholas M. Katz, *p -adic properties of modular schemes and modular forms*, Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), Springer, Berlin, 1973, pp. 69–190. Lecture Notes in Mathematics, Vol. 350. MR0447119 (56 #5434) ↑
- [30] Shinichi Kobayashi, *The p -adic Gross-Zagier formula for elliptic curves at supersingular primes*, Invent. Math. **191** (2013), no. 3, 527–629, DOI 10.1007/s00222-012-0400-9. MR3020170 ↑41, 42, 59
- [31] ———, *The p -adic height pairing on abelian varieties at non-ordinary primes*, preprint. ↑40
- [32] V. A. Kolyvagin, *Finiteness of $E(\mathbb{Q})$ and $SH(E, \mathbb{Q})$ for a subclass of Weil curves*, Izv. Akad. Nauk SSSR Ser. Mat. **52** (1988), no. 3, 522–540, 670–671 (Russian); English transl., Math. USSR-Izv. **32** (1989), no. 3, 523–541. MR954295 (89m:11056) ↑7
- [33] ———, *On the structure of Shafarevich-Tate groups*, Algebraic geometry (Chicago, IL, 1989), Lecture Notes in Math., vol. 1479, Springer, Berlin, 1991, pp. 94–121, DOI 10.1007/BFb0086267. MR1181210 (94b:11055) ↑9
- [34] V. A. Kolyvagin and D. Yu. Logachëv, *Finiteness of SH over totally real fields*, Izv. Akad. Nauk SSSR Ser. Mat. **55** (1991), no. 4, 851–876 (Russian); English transl., Math. USSR-Izv. **39** (1992), no. 1, 829–853. MR1137589 (93d:11063) ↑7
- [35] Yu. I. Manin, *Non-Archimedean integration and p -adic Jacquet-Langlands L -functions*, Uspehi Mat. Nauk **31** (1976), no. 1(187), 5–54 (Russian). MR0417134 (54 #5194) ↑4
- [36] Jan Nekovář, *On p -adic height pairings*, Séminaire de Théorie des Nombres, Paris, 1990–91, Progr. Math., vol. 108, Birkhäuser Boston, Boston, MA, 1993, pp. 127–202. MR1263527 (95j:11050) ↑39, 40
- [37] ———, *On the p -adic height of Heegner cycles*, Math. Ann. **302** (1995), no. 4, 609–686, DOI 10.1007/BF01444511. MR1343644 (96f:11073) ↑10, 34, 41
- [38] A. A. Panchishkin, *Convolutions of Hilbert modular forms and their non-Archimedean analogues*, Mat. Sb. (N.S.) **136(178)** (1988), no. 4, 574–587, 592 (Russian); English transl., Math. USSR-Sb. **64** (1989), no. 2, 571–584. MR965894 (89k:11033) ↑4
- [39] Alexei A. Panchishkin, *Motives over totally real fields and p -adic L -functions*, Ann. Inst. Fourier (Grenoble) **44** (1994), no. 4, 989–1023 (English, with English and French summaries). MR1306547 (96e:11087) ↑
- [40] Bernadette Perrin-Riou, *Fonctions L p -adiques associées à une forme modulaire et à un corps quadratique imaginaire*, J. London Math. Soc. (2) **38** (1988), no. 1, 1–32, DOI 10.1112/jlms/s2-38.1.1 (French). MR949078 (89m:11043) ↑18, 33, 35
- [41] ———, *Points de Heegner et dérivées de fonctions L p -adiques*, Invent. Math. **89** (1987), no. 3, 455–510, DOI 10.1007/BF01388982 (French). MR903381 (89d:11034) ↑2, 4, 7, 8, 9, 10, 14, 47, 51, 53, 54, 55, 56, 59
- [42] ———, *Théorie d’Iwasawa p -adique locale et globale*, Invent. Math. **99** (1990), no. 2, 247–292, DOI 10.1007/BF01234420 (French). MR1031902 (91b:11116) ↑
- [43] Kartik Prasanna, *Arithmetic properties of the Shimura-Shintani-Waldspurger correspondence*, Invent. Math. **176** (2009), no. 3, 521–600, DOI 10.1007/s00222-008-0169-z. With an appendix by Brian Conrad. MR2501296 (2011d:11102) ↑62
- [44] A. Raghuram and Naomi Tanabe, *Notes on the arithmetic of Hilbert modular forms*, J. Ramanujan Math. Soc. **26** (2011), no. 3, 261–319. MR2865819 (2012m:11060) ↑60
- [45] Peter Schneider, *p -adic height pairings. II*, Invent. Math. **79** (1985), no. 2, 329–374, DOI 10.1007/BF01388978. MR778132 (86j:11063) ↑7, 40
- [46] Jean-Pierre Serre, *Sur le résidu de la fonction zêta p -adique d’un corps de nombres*, C. R. Acad. Sci. Paris Sér. A-B **287** (1978), no. 4, A183–A188 (French, with English summary). MR0506177 (58 #22024) ↑30
- [47] Goro Shimura, *The special values of the zeta functions associated with Hilbert modular forms*, Duke Math. J. **45** (1978), no. 3, 637–679. MR507462 (80a:10043) ↑3, 6, 31
- [48] ———, *The periods of certain automorphic forms of arithmetic type*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **28** (1981), no. 3, 605–632 (1982). MR656039 (84f:10040) ↑6, 62
- [49] ———, *Algebraic relations between critical values of zeta functions and inner products*, Amer. J. Math. **105** (1983), no. 1, 253–285, DOI 10.2307/2374388. MR692113 (84j:10038) ↑61, 62
- [50] Christopher Skinner and Eric Urban, *The Iwasawa Main Conjectures for GL_2* , Invent. Math. **195** (2014), no. 1, 1–277, DOI 10.1007/s00222-013-0448-1. MR3148103 ↑8
- [51] Jeanine Van Order, *On the Dibedral Main Conjectures of Iwasawa Theory for Hilbert Modular Eigenforms*, Canad. J. Math., to appear. ↑9
- [52] J.-L. Waldspurger, *Sur les valeurs de certaines fonctions L automorphes en leur centre de symétrie*, Compositio Math. **54** (1985), no. 2, 173–242 (French). MR783511 (87g:11061b) ↑8, 19
- [53] Hiroyuki Yoshida, *On the zeta functions of Shimura varieties and periods of Hilbert modular forms*, Duke Math. J. **75** (1994), no. 1, 121–191, DOI 10.1215/S0012-7094-94-07505-4. MR1284818 (95d:11059) ↑61

- [54] ———, *On a conjecture of Shimura concerning periods of Hilbert modular forms*, Amer. J. Math. **117** (1995), no. 4, 1019–1038, DOI 10.2307/2374957. MR1342839 (96d:11056) ↑62
- [55] Xinyi Yuan, Shou-Wu Zhang, and Wei Zhang, *The Gross-Zagier Formula on Shimura Curves*, Annals of Mathematics Studies, vol. 184, Princeton University Press, Princeton, NJ, 2012. ↑6, 9, 19, 22
- [56] Yuri G. Zarhin, *p -adic heights on abelian varieties*, Séminaire de Théorie des Nombres, Paris 1987–88, Progr. Math., vol. 81, Birkhäuser Boston, Boston, MA, 1990, pp. 317–341. MR1042777 (91f:11043) ↑39
- [57] Shou-Wu Zhang, *Heights of Heegner points on Shimura curves*, Ann. of Math. (2) **153** (2001), no. 1, 27–147, DOI 10.2307/2661372. MR1826411 (2002g:11081) ↑2, 4, 6, 7, 8, 9, 10, 14, 16, 17, 19, 25, 26, 28, 31, 34, 45, 46, 47, 48, 49, 50, 52, 53
- [58] ———, *Gross-Zagier formula for GL_2* , Asian J. Math. **5** (2001), no. 2, 183–290. MR1868935 (2003k:11101) ↑2, 7, 10, 45, 50, 52
- [59] ———, *Gross-Zagier formula for $GL(2)$. II, Heegner points and Rankin L -series*, Math. Sci. Res. Inst. Publ., vol. 49, Cambridge Univ. Press, Cambridge, 2004, pp. 191–214, DOI 10.1017/CBO9780511756375.008. MR2083213 (2005k:11121) ↑2, 8, 10
- [60] Wei Zhang, *Selmer groups and the divisibility of Heegner points*, preprint. ↑8

DANIEL DISEGNI, Department of Mathematics and Statistics, McGill University, 805 Shebrooke St. West, Montreal, QC H3A 0B9, Canada • *E-mail*: daniel.disegni@mcgill.ca